

LDL Level in Ischaemic Stroke Patients at Dr. Soetomo General Hospital Surabaya

by Mayori Rachmawati

Submission date: 29-Aug-2020 11:49PM (UTC+0800)

Submission ID: 1375979271

File name: mic_Stroke_Patients_at_Dr._Soetomo_General_Hospital_Surabaya.pdf (243.21K)

Word count: 2323

Character count: 11684

ORIGINAL ARTICLE

LDL Level in Ischaemic Stroke Patients at Dr. Soetomo General Hospital Surabaya

Mayori Rachmawati¹, Rr. Indrayuni Lukitra Wardhani², Paulus Sugianto^{3*}

¹Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia

²Department of Physical and Rehabilitation Medicine, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia - Dr. Soetomo General Hospital Surabaya, Indonesia

³Department of Neurology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia - Dr. Soetomo General Hospital Surabaya, Indonesia

ARTICLE INFO

Article history:
Received 11 May 2019
Received in revised form 27 May 2019
Accepted 24 June 2019

Keywords:
Ischemic stroke,
Total Cholesterol,
Low Density Lipoprotein,
High Density Lipoprotein,
Tryglicerides.

ABSTRACT

Introduction: Stroke is one of the top killers amongst the non-infectious diseases. Ischemic stroke dominates the rate by 87.0% amongst all type of strokes. One of the risk factors for stroke ischemic stroke is atherosclerosis which caused by lipid build-up in the arteries.

Methods: This research is a retrospective study on secondary data. Admitted Ischaemic stroke patients. Lipid profile, and Barthel Index were collected from medical record and presented descriptively.

Results: Of the 248 patients, 144 (58.1%) were men, mean age was 59.33. 61.7% ischemic stroke patients have their LDL level elevated, 59.7% HDL decreased, 39.1% total cholesterol elevated, and 33.9% TG elevated. This study also observe the dependency based on Barthel Index of the admitted stroke patients. 75% percents amongst subjects were categorized as "total dependece" are having their LDL level elevated.

Conclusion: Most of stroke patients have elevated LDL and HDL while the TG and total cholesterol are mostly optimized. Stroke patients whose admitted with "total dependence" most likely to have elevated LDL in this study.

© 2018 Biomolecular and Health Science Journal. All rights reserved

Introduction

Amongst the non-infectious diseases in the world, stroke is the second common cause of deaths of long-term disability with high incidence rate and deaths (21.2%) which percentage is higher than ischemic heart disease (8.9%) and diabetes mellitus (6.5%).¹

Based on trends of global burden data from 1990 - 2013, adults aged 20 to 64 years, the global prevalence of ischemic stroke was 7.3 million cases, with men having consistently greater incidence of ischemic stroke than women (133/100,000 and 99/100,000, respectively).^{2,3}

There are two types of major stroke, ischaemic and haemorrhagic. Ischaemic stroke leads with 87.0% amongst all strokes.⁴ Stroke is a clinical manifest of brain dysfunction (local or global) occurs when there are problems at brain's vessels such as something clogs blood supply to part of the brain (ischemic stroke) or when a blood vessel in the brain ruptures (hemorrhagic stroke).⁴ Regarding to American Heart Association, Ischemic stroke is an infarct in central nervous system (brain, vertebral spine, and retinal ganglion cells) which manifest with visible symptoms so that it will differentiate from silent stroke.⁵ Risk factors for strokes are divided into modifiable and non-modifiable. Age,

sex, genetics, and race are non-modifiable risk factors.⁶ Smoking, alcohol abuse, do no exercise, having history of diseases such as stroke, diabetes, dyslipidemia, hypertension, and other cardiovascular diseases are examples of modifiable risk factors.⁷

There are three major etiology in ischemic stroke which is cardioembolic, vasculopathy, and atherosclerosis.^{8,9} Atherosclerosis is generally caused by LDL accumulation inside vascular endothelial.^{1,10,11} Saposnik et al (2014) found that more than 45% patients with recurrent stroke attacks are presented with high LDL level (> 100mg/dL).¹² Thus, measuring lipid profile in ischemic stroke patients could be useful to help diagnostic and therapy. We conducted this study to know LDL level is presented and impact in ischemic stroke patients at Dr. Soetomo General Hospital.

Methods

This study was conducted at Dr. Soetomo General Hospital (Surabaya) from January to March 2018. This research had been approved by research committee of Dr. Soetomo General Hospital (0113/KEPK/III/2018). All admitted Ischemic stroke patients within inclusion criteria: had complete record and lipid profile. The

*Correspondence: paulus.sugianto@gmail.com

exclusion criteria were any data which was incomplete or data couldn't be read clearly. Age, sex, admission date, lipid profile, and Barthel Index were collected from medical record. The lipid profile which was used to conduct this research was the lipid profile measured when firstly admitted to Dr. Soetomo General Hospital Surabaya. Descriptive data was presented with tables. Barthel index is also used as an additional data to describe the outcome of stroke patients.

Results

From 309 stroke patients whom admitted from January to March 2018, 248 was included into the criteria. 144 patients (58.1%) were men with men to women ratio was 1.38:1 and average age was 59.61 (SD 13.162). The most common group was 55 – 64 years old. Those range of age was still productive age.

From table 2 we can see that in this study patients with not optimized LDL level was more common (71.9%). Patients with desirable level of HDL was lower than half of the total patients (40.3 %). The percentage of patients with normal triglycerides level and normal total cholesterol level were dominant.

Stroke patients' disability was measured by Barthel index. Table 3 explained that patients with total dependency and not optimized LDL level were 75%, patients with severe dependency and not optimized LDL level were 58.3 %, patients with moderate dependency and not optimized LDL level were 75%. Patients with low Barthel index score (slight dependent and independent) were only 10% of all patients whom Barthel index score were recorded.

Table 1. Age and Sex distribution of Dr. Soetomo General Hospital Ischemic Stroke Patients

Age	Men	Women	Total
Age (10 years group)			
<15	1	0	1
15 - 24	1	1	2
25 - 34	5	1	6
35 - 44	9	8	17
45 - 54	31	19	50
55 - 64	53	39	92
65 - 74	29	27	56
75 - 84	11	7	18
85 - 94	2	2	4
>95	2	0	2
Total	144	104	248

Table 2 Distributions of Dr. Soetomo General Hospital Ischemic Stroke Patients Based on Their Lipid Profile

Lipid Profile	Frequencies	%
Optimal (<100 mg/dl)	70	28.1
Near Optimal (100-129 mg/dl)	77	31.0
Low Density Lipoprotein		
Borderline High (130-159 mg/dl)	61	24.6 %
High (160-189 mg/dl)	30	12.1
Very High (\geq 190 mg/dl)	10	
High Density Lipoprotein		
Low (< 40 mg/dl)	130	59.7
Desirable (> 40 mg/dl)	118	40.3
Triglycerides		
Normal (< 150 mg/dl)	164	66.1
Borderline high (150-199 mg/dl)	37	14.9
High (200-499 mg/dl)	46	18.5
Very high (\geq 500 mg/dl)	1	0.4
Total Cholesterol		
Desirable (<200 mg/dl)	151	60.9
Borderline High (200-239 mg/dl)	61	24.6
High (240mg/dl)	36	14.5

Table 3. Cross table of Barthel index and LDL level in Dr. Soetomo General Hospital Ischemic Stroke Patients

Admitted Barthel Index	LDL Profile					Total
	Optimal	Near Optimal	Borderline High	High	Very High	
Total	5	6	6	2	1	20
Severe	5	3	4	0	0	12
Moderate	3	3	5	1	0	12
Slight	1	2	0	1	1	5
Independent	0	0	0	0	0	0
Total	14	14	15	4	2	49

Discussion

From result, male was dominant as stated in previous study that ischemic stroke is more common in men rather than women because men tend to smoke more often than women.¹³ Other than smoke habit, estrogen level which is higher in women also become a protective agent for blood vessels. Estrogen treatments have been researched for women who has already in menopause. The theory is still in controversy due to possible side effects of exogenic estrogen, but it is proven that estrogen level is associated with reduction of peripheral vascular resistance.¹⁴

In this study, patients with desirable total cholesterol (60.9%) were more than both the borderline high and high combined (39.1%). A study also found the same result as prior research.¹⁵ Triglycerides level in this study is normal in more than half of the patients (66.1%). Other study also found that more than half of the subjects are having normal triglycerides.¹⁶ According to AHA journal, triglycerides and total cholesterol level aren't independent risk for ischemic stroke, however there was a suggestion of increased risk for patients with high triglycerides and total cholesterol level.¹⁷ High HDL in blood is considered good.¹⁸ HDL is commonly considered as "good" lipoprotein for its role in carried cholesterol to the liver so that it could be flushed out from the body and lowering risk of atherogenesis process.¹¹ In this study, it is found that patients with low HDL level (59.7%) is more than patients with high HDL level.

The results of this study are 71.9 % of the patients have their LDL level elevated. Previous study in 2017 also found the same result.¹⁹ On the other hand, a study by Siddeswari in India found different result, which was patients with elevated LDL level were only 21.6%.²⁰ Previous studies found that unoptimized LDL has been strongly related with stroke patient's mortality, poor outcome, and disability.^{21,23} LDL is an atherogenic agents because of its tendency of getting built up inside the vascular endothelial and causing atherosclerosis, narrow the vascular and causing stroke.^{1,10,11} The high level of LDL in most of the patients in this study describes the indirect effect of high LDL as a risk factor of ischemic stroke. Patients with total dependency and not optimized LDL level were 75%, and patients with severe dependency and not optimized LDL level were 58.3%. This shows high level of LDL will most likely to have worse outcome. This study has several limitations such as that the use of statin or other dyslipidemia treatments are not being recorded in this study. Further study considering the treatment for dyslipidemia is considered for follow up also study regarding the effect of dyslipidemia drugs on stroke patients' outcome.

Conclusion

This study concluded that most subjects have high LDL level which might affect their functional outcome when discharge out of the hospital. This study also finds more than half of all patients have low HDL level.

Conflict of Interest

The author stated there is no conflict of interest

References

- Indonesia, B.P.d.P.K.K.K.R., Riset Kesehatan Dasar (RISKESDAS) 2013. 2013.
- Feigin, V.L., B. Norrving, and G.A. Mensah, Global Burden of Stroke. *Circ Res*, 2017. 120(3): p. 439-448.
- About Stroke. 2018 [cited 2018 7th August]; Available from: <https://www.cdc.gov/stroke/about.htm>.
- McCance, K. and S. Hueter, Pathophysiology: the Biologic Basis for Disease in Adults and Children. Seventh Ed ed. 2014, St. Louis, Missouri: ELSEVIER.
- Sacco, R.L., et al., An Updated Definition of Stroke for the 21st Century: a Statement for Healthcare Professionals from the American Heart Association/American Stroke Association. *Stroke*, 2013. 44(7): p. 2064-89.
- Reprint: 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk. *J Am Pharm Assoc* (2003), 2013: p. e4.
- Fahimfar, N., et al., Risk Factors for Ischemic Stroke; Results from 9 Years of Follow up in a Population Based Cohort of Iran. *BMC Neurol*, 2012. 12: p. 117.
- Kass, J. and E.M. Mizrahi, *Neurology Secrets: Sixth Edition*. 2016. 1-531.
- Tuomilehto, J., M. Mähönen, and C. Sarti, Basic epidemiology of stroke and risk assessment, in *Textbook of Stroke Medicine*, M. Brainin and W.-D. Heiss, Editors. 2009, Cambridge University Press: Cambridge. p. 77-88.
- Mertens, A. and P. Holvoet, Oxidized LDL and HDL: Antagonists in Atherothrombosis. *FASEB J*, 2001. 15(12): p. 2073-84.
- Sherwood, L., *Fisiologi Manusia dari Sel ke Sistem Edisi 9*. 2013, Jakarta: EGC.
- Saposnik, G., et al., Guideline-directed Low Density Lipoprotein Management in High Risk Patients with Ischemic Stroke: Findings from Get with the Guidelines-Stroke 2003 to 2012. *Stroke*, 2014. 45(11): p. 3343-51.
- Appelros, P., B. Stegmayr, and A. Terent, Sex Differences in Stroke Epidemiology: a Systematic Review. *Stroke*, 2009. 40(4): p. 1082-90.
- Nicholson, C.J., et al., Estrogenic Vascular Effects are Diminished by Chronological Aging. *Sci Rep*, 2017. 7(1): p. 12153.
- Olamoyegun, M.A., R. Oluoyombo, and S.O. Asaolu, Evaluation of Dyslipidemia, Lipid Ratios, and Atherogenic Index as Cardiovascular Risk Factors Among Semi Urban Dwellers in Nigeria. *Ann Afr Med*, 2016. 15(4): p. 194-199.
- Karim, M., et al., Association of Hypertriglyceridemia with Ischemic Stroke, Study in a Tertiary Care Hospital in Bangladesh. *Journal of Medicine*, 2016. 17(1): p. 21-6.
- Bowman, T.S., et al., Cholesterol and the Risk of Ischemic Stroke. *Stroke*, 2003. 34(12): p. 2930-4.
- NHLBI, NCEP-ATP III Guidelines At-A-Glance Quick Desk Reference, in *National Cholesterol Education Program, U.S.D.O.H.A.H. SERVICES*, et al., Editors. 2001.
- Rai, O.N. and A. Kumar, Study of Serum Lipid Profile in Stroke Patients in Northern India. 2017, 2017. 4(5): p. 4.
- Siddeswari, R., et al., Pattern of Dyslipidemia in Ischemic Stroke. *J Med Allied Sci*, 2015. 5(2): p. 26-9.
- Song, T.J., et al., Low Density Lipoprotein Particle Size Predicts a Poor Outcome in Patients with Atherothrombotic Stroke. *J Clin Neurol*, 2015. 11(1): p. 80-6.
- Tsai, N.W., et al., Statin Therapy Reduces Oxidized Low Density Lipoprotein Level, a Risk Factor for Stroke Outcome. *Crit Care*, 2014. 18(1): p. R16.
- Xing, Y., et al., Low Density Lipoprotein Cholesterol and the Outcome of Acute Ischemic Stroke: Results of a Large Hospital-Based Study. *Eur Neurol*, 2016. 76(5-6): p. 195-201.

LDL Level in Ischaemic Stroke Patients at Dr. Soetomo General Hospital Surabaya

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

12%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Airlangga Student Paper	2%
2	www.studyblue.com Internet Source	2%
3	academic.oup.com Internet Source	2%
4	Valery L. Feigin, Bo Norrving, George A. Mensah. "Global Burden of Stroke", Circulation Research, 2017 Publication	1%
5	salemhospital.com Internet Source	1%
6	Fernando Fonseca, Marzia Macedo, Marisa da Cunha Gonzaga, Carlos D'Aparecida Machado, Anna Batista. "Study of lipid profile in adult women with acne", Clinical, Cosmetic and Investigational Dermatology, 2015 Publication	1%

7	doi.org Internet Source	1%
8	www.scribd.com Internet Source	1%
9	Dirani Dirani, Made Sedana, Siti Wahyuni, Chrismawan Ardianto, Chris Alderman. "Evaluation to the chemotherapy use in patients with diffuse large B-cell lymphoma", Journal of Basic and Clinical Physiology and Pharmacology, 2020 Publication	1%
10	bcn.iums.ac.ir Internet Source	1%
11	Laila Mahmoud Mohamed Gad. "Chapter 4 Atherosclerosis and Antihyperlipidemic Agents", IntechOpen, 2012 Publication	1%
12	jurnal.unissula.ac.id Internet Source	1%
13	journals.plos.org Internet Source	1%
14	banglajol.info Internet Source	1%
15	Josef Yayan. "Effects of alteplase in the treatment of acute ischemic stroke",	<1%

16

scholarbank.nus.edu.sg

Internet Source

<1%

17

Ilona Kopyta, Mikołaj Zimny, Beata Sarecka-Hujar. "The role of biochemical risk factors in the etiology of AIS in children and adults", International Journal of Neuroscience, 2014

Publication

<1%

18

"1-A1: Lung Cancer 1 : Poster Sessions", Respirology, 2013.

Publication

<1%

19

Artono, Bakti Surarto, Nyilo Purnami, Fransiska Hutahaen, M. Reza Mahardhika. "The Association of IL-1 Alpha Level and TNF Alpha Expressions on Bone Destruction in Chronic Suppurative Otitis Media and Cholesteatoma", Indian Journal of Otolaryngology and Head & Neck Surgery, 2019

Publication

<1%

20

"Abstracts", Hepatology International, 2020

Publication

<1%

21

Kazumitsu Nawata. "Risk Factors Affecting Ischemic Stroke: A Potential Side Effect of Antihypertensive Drugs", Health, 2020

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

LDL Level in Ischaemic Stroke Patients at Dr. Soetomo General Hospital Surabaya

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
