

Morrey spaces and fractional integral operators

by Eridani Eridani

Submission date: 13-Dec-2018 11:06AM (UTC+0800)

Submission ID: 1056164004

File name: Morrey_spaces_and_fractional_integral_operators.pdf (345.1K)

Word count: 5265

Character count: 19412

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Morrey spaces and fractional integral operators

A. Eridani^{a, b}, Vakhtang Kokilashvili^c, Alexander Meskhi^{b, c, *}

^a*Department of Mathematics, Airlangga University, Campus C, Mulyorejo, Surabaya 60115, Indonesia*

^b*School of Mathematical Sciences, Government College University, 68-B New Muslim Town, Lahore, Pakistan*

^c*A. Razmadze Mathematical Institute, M. Aleksidze St., Tbilisi 0193, Georgia*

Received 4 July 2008; received in revised form 8 December 2008

Abstract

The present paper is devoted to the boundedness of fractional integral operators in Morrey spaces defined on quasimetric measure spaces. In particular, Sobolev, trace and weighted inequalities with power weights for potential operators are established. In the case when measure satisfies the doubling condition the derived conditions are simultaneously necessary and sufficient for appropriate inequalities.

© 2009 Elsevier GmbH. All rights reserved.

MSC 2000: primary 26A33; secondary 42B35; 47B38

Keywords: Fractional integrals; Morrey spaces; Non-homogeneous spaces; Trace inequality; Two-weight inequality

1. Introduction

The main purpose of this paper is to establish the boundedness of fractional integral operators in (weighted) Morrey spaces defined on quasimetric measure spaces. We derive Sobolev, trace and two-weight inequalities for fractional integrals. In particular, we

* Corresponding author at: School of Mathematical Sciences, Government College University, 68-B New Muslim Town, Lahore, Pakistan. Tel.: +995 32 32 62 47; fax: +995 32 36 40 86.

E-mail addresses: keumala_ikhsanti@yahoo.com (A. Eridani), kokil@rmi.acnet.ge (V. Kokilashvili), meskhi@rmi.acnet.ge (A. Meskhi).

generalize: (a) Adams [1] trace inequality; (b) the theorem by Stein and Weiss [18] regarding the two-weight inequality for the Riesz potentials; (c) Sobolev-type inequality. We emphasize that in the most cases the derived conditions are necessary and sufficient for appropriate inequalities.

In the paper [9] (see also [10, Chapter 2]) integral-type sufficient condition guaranteeing the two-weight weak-type inequality for integral operator with positive kernel defined on non-homogeneous spaces was established. In the same paper (see also [10, Chapter 2]) the authors solved the two-weight problem for kernel operators on spaces of homogeneous type.

In [12] (see also [5, Chapter 6]) a complete description of non-doubling measure μ guaranteeing the boundedness of fractional integral operator I_α (see the next section for the definition) from $L^p(\mu, X)$ to $L^q(\mu, X)$, $1 < p < q < \infty$, was given. We notice that this result was derived in [11] for potentials on Euclidean spaces. In [12], theorems of Sobolev and Adams type for fractional integrals defined on quasimetric measure spaces were established. For the boundedness of fractional integrals on metric measure spaces we refer also to [7]. Some two-weight norm inequalities for fractional operators on \mathbf{R}^n with non-doubling measure were studied in [8]. Further, in the paper [13] necessary and sufficient conditions on measure μ governing the inequality of Stein–Weiss type on non-homogeneous spaces were established. For some properties of fractional integrals defined on \mathbf{R}^n in weighted Lebesgue spaces with power type weights see e.g., [16, Chapter 5].

The boundedness of the Riesz potential in Morrey spaces defined on Euclidean spaces was studied in [15,2]. The same problem for fractional integrals on \mathbb{R}^n with non-doubling measure was investigated in [17].

Finally, we mention that necessary and sufficient conditions for the boundedness of maximal operators and Riesz potentials in the local Morrey-type spaces were derived in [3,4].

The main results of this paper were presented in [6].

It should be emphasized that the results of this work are new even for Euclidean spaces.

Constants (often different constants in the same series of inequalities) will generally be denoted by c or C .

2. Preliminaries

Throughout the paper we assume that $X := (X, \rho, \mu)$ is a topological space, endowed with a complete measure μ such that the space of compactly supported continuous functions is dense in $L^1(X, \mu)$ and there exists a function (quasimetric) $\rho : X \times X \rightarrow [0, \infty)$ satisfying the conditions:

- (1) $\rho(x, y) > 0$ for all $x \neq y$, and $\rho(x, x) = 0$ for all $x \in X$;
- (2) there exists a constant $a_0 \geq 1$, such that $\rho(x, y) \leq a_0 \rho(y, x)$ for all $x, y \in X$;
- (3) there exists a constant $a_1 \geq 1$, such that $\rho(x, y) \leq a_1(\rho(x, z) + \rho(z, y))$ for all $x, y, z \in X$.

We assume that the balls $B(a, r) := \{x \in X : \rho(a, x) < r\}$ are μ -measurable and $0 < \mu(B(a, r)) < \infty$ for $a \in X, r > 0$. For every neighborhood V of $x \in X$, there exists $r > 0$, such that $B(x, r) \subset V$. We also assume that $\mu(X) = \infty, \mu\{a\} = 0$, and $B(a, r_2) \setminus B(a, r_1) \neq \emptyset$, for all $a \in X, 0 < r_1 < r_2 < \infty$.

The triple (X, ρ, μ) will be called quasimetric measure space.

Let $0 < \alpha < 1$. We consider the fractional integral operators I_α , and K_α given by

$$I_\alpha f(x) := \int_X f(y) \rho(x, y)^{\alpha-1} d\mu(y),$$

$$K_\alpha f(x) := \int_X f(y) (\mu B(x, \rho(x, y)))^{\alpha-1} d\mu(y),$$

for suitable f on X .

Suppose that ν is another measure on X , $\lambda \geq 0$ and $1 \leq p < \infty$. We deal with the Morrey space $L^{p,\lambda}(X, \nu, \mu)$, which is the set of all functions $f \in L^p_{loc}(X, \nu)$ such that

$$\|f\|_{L^{p,\lambda}(X, \nu, \mu)} := \sup_B \left(\frac{1}{(\mu(B)^\lambda)} \int_B |f(y)|^p d\nu(y) \right)^{1/p} < \infty,$$

where the supremum is taken over all balls B .

If $\nu = \mu$, then we have the classical Morrey space $L^{p,\lambda}(X, \mu)$ with measure μ . When $\lambda = 0$, then $L^{p,\lambda}(X, \nu, \mu) = L^p(X, \nu)$ is the Lebesgue space with measure ν .

Further, suppose that $\beta \in \mathbf{R}$. We are also interested in weighted Morrey space $M^{p,\lambda}_\beta(X, \mu)$ which is the set of all μ -measurable functions f such that

$$\|f\|_{M^{p,\lambda}_\beta(X, \mu)} := \sup_{a \in X; r > 0} \left(\frac{1}{r^\lambda} \int_{B(a,r)} |f(y)|^p \rho(a, y)^\beta d\mu(y) \right)^{1/p} < \infty.$$

If $\beta = 0$, then we denote $M^{p,\lambda}_\beta(X, \mu) := M^{p,\lambda}(X, \mu)$.

We say that a measure μ satisfies the growth condition ($\mu \in (GC)$), if there exists $C_0 > 0$ such that $\mu(B(a, r)) \leq C_0 r$; further, μ satisfies the doubling condition ($\mu \in (DC)$) if $\mu(B(a, 2r)) \leq C_1 \mu(B(a, r))$ for some $C_1 > 1$. If $\mu \in (DC)$, then (X, ρ, μ) is called a space of homogeneous type (SHT). A quasimetric measure space (X, ρ, μ) , where the doubling condition is not assumed, is also called a non-homogeneous space.

The measure μ on X satisfies the reverse doubling condition ($\mu \in (RDC)$) if there are constants η_1 and η_2 with $\eta_1 > 1$ and $\eta_2 > 1$ such that

$$\mu B(x, \eta_1 r) \geq \eta_2 \mu B(x, r). \tag{1}$$

It is known (see e.g., [19, p. 11]) that if $\mu \in (DC)$, then $\mu \in (RDC)$.

The next statements are from [12] (see also [5, Theorem 6.1.1, Corollary 6.1.1] and [11] in the case of Euclidean spaces).

Theorem A. Let (X, ρ, μ) be a quasimetric measure space. Suppose that $1 < p < q < \infty$ and $0 < \alpha < 1$. Then I_α is bounded from $L^p(X)$ to $L^q(X)$ if and only if there exists a positive constant C such that

$$\mu(B(a, r)) \leq C r^s, \quad s = \frac{pq(1-\alpha)}{pq+p-q}, \tag{2}$$

for all $a \in X$ and $r > 0$.

Corollary B. Let (X, ρ, μ) be a quasimetric measure space, $1 < p < 1/\alpha$ and $1/q = 1/p - \alpha$. Then I_α is bounded from $L^p(X)$ to $L^q(X)$ if and only if $\mu \in (GC)$.

The latter statement by different proof was also derived in [7] for metric spaces.

To prove some of our statements we need the following Hardy-type transform:

$$H_a f(x) := \int_{\rho(a,y) \leq \rho(a,x)} f(y) d\mu(y),$$

where a is a fixed point of X and $f \in L_{loc}(X, \mu)$.

Theorem C. Suppose that (X, ρ, μ) is a quasimetric measure space and $1 < p \leq q < \infty$. Assume that ν is another measures on X . Let V (resp. W) be non-negative $\nu \times \nu$ -measurable (resp. non-negative $\mu \times \mu$ -measurable) function on $X \times X$. If there exists a positive constant C independent of $a \in X$ and $t > 0$ such that

$$\left(\int_{\rho(a,y) \geq t} V(a, y) d\nu(y) \right)^{1/q} \left(\int_{\rho(a,y) \leq t} W(a, y)^{1-p'} d\mu(y) \right)^{1/p'} \leq C < \infty,$$

then there exists a positive constant c such that for all μ -measurable non-negative f and $a \in X$ the inequality

$$\left(\int_{B(a,r)} (H_a f(x))^q V(a, x) d\nu(x) \right)^{1/q} \leq c \left(\int_{B(a,r)} (f(x))^p W(a, x) d\mu(x) \right)^{1/p}$$

holds.

This statement was proved in [5, Section 1.1] for Lebesgue spaces.

Proof of Theorem C. Let $f \geq 0$. We define $S(s) := \int_{\rho(a,y) < s} f(y) d\mu(y)$, for $s \in [0, r]$. Suppose $S(r) < \infty$, then $2^m < S(r) \leq 2^{m+1}$, for some $m \in \mathbb{Z}$. Let

$$s_j := \sup\{t : S(t) \leq 2^j\}, \quad j \leq m \quad \text{and} \quad s_{m+1} := r.$$

Then it is easy to see that (see also [5, pp. 5–8] for details) $(s_j)_{j=-\infty}^{m+1}$ is a non-decreasing sequence, $S(s_j) \leq 2^j$, $S(t) \geq 2^j$ for $t > s_j$, and

$$2^j \leq \int_{s_j \leq \rho(a,y) \leq s_{j+1}} f(y) d\mu(y).$$

If $\beta := \lim_{j \rightarrow -\infty} s_j$, then

$$\rho(a, x) < r \Leftrightarrow \rho(a, x) \in [0, \beta] \cup \bigcup_{j=-\infty}^m (s_j, s_{j+1}].$$

If $S(r) = \infty$, then we may put $m = \infty$. Since

$$0 \leq \int_{\rho(a,y) < \beta} f(y) d\mu(y) \leq S(s_j) \leq 2^j,$$

for every j , therefore $\int_{\rho(a,y) < \beta} f(y) d\mu(y) = 0$. From these observations, we have

$$\begin{aligned} & \int_{\rho(a,x) < r} (H_a f(x))^q V(a, x) dv(x) \\ & \leq \sum_{j=-\infty}^m \int_{s_j \leq \rho(a,x) \leq s_{j+1}} (H_a f(x))^q V(a, x) dv(x) \\ & \leq \sum_{j=-\infty}^m \int_{s_j \leq \rho(a,x) \leq s_{j+1}} V(a, x) \left(\int_{\rho(a,y) \leq s_{j+1}} (f(y)) d\mu(y) \right)^q dv(x). \end{aligned}$$

Notice that

$$\int_{\rho(a,y) \leq s_{j+1}} f d\mu \leq S(s_{j+2}) \leq 2^{j+2} \leq C \int_{s_{j-1} \leq \rho(a,y) \leq s_j} f d\mu.$$

Using Hölder's inequality, we find that

$$\begin{aligned} & \int_{\rho(a,x) < r} (H_a f(x))^q V(a, x) d\mu(x) \\ & \leq \sum_{j=-\infty}^m \int_{s_j \leq \rho(a,x) \leq s_{j+1}} V(a, x) \left(\int_{\rho(a,y) \leq s_{j+1}} (f(y)) d\mu(y) \right)^q dv(x) \\ & \leq C \sum_{j=-\infty}^m \int_{s_j \leq \rho(a,x) \leq s_{j+1}} V(a, x) \left(\int_{s_{j-1} \leq \rho(a,y) \leq s_j} (f(y)) d\mu(y) \right)^q dv(x) \\ & \leq C \sum_{j=-\infty}^m \int_{s_j \leq \rho(a,x) \leq s_{j+1}} V(a, x) dv(x) \\ & \quad \times \left(\int_{s_{j-1} \leq \rho(a,y) \leq s_j} (f(y))^p W(a, y) d\mu(y) \right)^{q/p} \\ & \quad \times \left(\int_{s_{j-1} \leq \rho(a,y) \leq s_j} W(a, y)^{1-p'} d\mu(y) \right)^{q/p'} \\ & \leq C \sum_{j=-\infty}^m \int_{s_j \leq \rho(a,y)} V(a, y) dv(y) \left(\int_{\rho(a,y) \leq s_j} W(a, y)^{1-p'} d\mu(y) \right)^{q/p'} \\ & \quad \left(\int_{s_{j-1} \leq \rho(a,y) \leq s_j} (f(y))^p W(a, y) d\mu(y) \right)^{q/p} \\ & \leq C \sum_{j=-\infty}^m \left(\int_{s_{j-1} \leq \rho(a,y) \leq s_j} (f(y))^p W(a, y) d\mu(y) \right)^{q/p} \\ & \leq C \left(\int_{\rho(a,y) \leq r} (f(y))^p W(a, y) d\mu(y) \right)^{q/p}. \end{aligned}$$

This completes the proof of the theorem. \square

For our purposes we also need the following lemma (see [14] for the case of \mathbf{R}^n).

Lemma D. Suppose that (X, ρ, μ) be an SHT. Let $0 < \lambda < 1 \leq p < \infty$. Then there exists a positive constant C such that for all balls B_0 ,

$$\|\chi_{B_0}\|_{L^{p,\lambda}(X,\mu)} \leq C \mu(B_0)^{(1-\lambda)/p}.$$

Proof. Let $B_0 := B(x_0, r_0)$ and $B := B(a, r)$. We have

$$\|\chi_{B_0}\|_{L^{p,\lambda}(X,\mu)} = \sup_B \left(\frac{\mu(B_0 \cap B)}{\mu(B)^\lambda} \right)^{1/p}.$$

Suppose that $B_0 \cap B \neq \emptyset$. Let us assume that $r_0 \leq r$. Then (see [19, Lemma 1] or [10, p. 9]) $B \subset B(x_0, br_0)$, where $b = a_1(1 + a_0)$. By the doubling condition it follows that

$$\begin{aligned} \frac{\mu(B \cap B_0)}{\mu(B)^\lambda} &\leq \frac{\mu(B)}{\mu(B)^\lambda} = \mu(B)^{1-\lambda} \leq \mu(B(x_0, br_0))^{1-\lambda} \\ &\leq C \mu(B_0)^{1-\lambda}. \end{aligned}$$

Let now $r_0 < r$. Then $\mu B_0 \leq c \mu B$, where the constant c depends only on a_1 and a_0 . Then

$$\frac{\mu(B \cap B_0)}{\mu(B)^\lambda} \leq c \frac{\mu(B_0)}{\mu(B_0)^\lambda} = c \mu(B_0)^{1-\lambda}. \quad \square$$

The next lemma may be well known but we prove it for the completeness.

Lemma E. Let (X, ρ, μ) be a non-homogeneous space with the growth condition. Suppose that $\sigma > -1$. Then there exists a positive constant c such that for all $a \in X$ and $r > 0$, the inequality

$$I(a, r, \sigma) := \int_{B(a,r)} \rho(a, x)^\sigma d\mu \leq cr^{\sigma+1}$$

holds.

Proof. Let $\sigma \geq 0$. Then the result is obvious because of the growth condition for μ . Further, assume that $-1 < \sigma < 0$. We have

$$\begin{aligned} I(a, r, \sigma) &= \int_0^\infty \mu\{x \in B(a, r) : \rho(a, x)^\sigma > \lambda\} d\lambda \\ &= \int_0^\infty \mu(B(a, r) \cap B(a, \lambda^{1/\sigma})) d\lambda = \int_0^{r^\sigma} + \int_{r^\sigma}^\infty := I^{(1)}(a, r, \sigma) + I^{(2)}(a, r, \sigma). \end{aligned}$$

By the growth condition for μ we have

$$I^{(1)}(a, r, \sigma) \leq r^\sigma \mu(B(a, r)) \leq cr^{\sigma+1},$$

while for $I^{(2)}(a, r, \sigma)$ we find that

$$I^{(2)}(a, r, \sigma) \leq c \int_{r^\sigma}^\infty \lambda^{1/\sigma} d\lambda = \frac{-c(\sigma+1)}{\sigma} r^{\sigma+1} = c_1 r^{\sigma+1}$$

because $1/\sigma < -1$. \square

The following statement is the trace inequality for the operator K_α (see [1] for the case of Euclidean spaces and, e.g., [10] or [5, Theorem 6.2.1] for an SHT).

Theorem F. Let (X, ρ, μ) be an SHT. Suppose that $1 < p < q < \infty$ and $0 < \alpha < 1/p$. Assume that ν is another measure on X . Then K_α is bounded from $L^p(X, \mu)$ to $L^q(X, \nu)$ if and only if

$$\nu B \leq c(\mu B)^{q(1/p-\alpha)},$$

for all balls B in X .

3. Main results

In this section we formulate the main results of the paper. We begin with the case of an SHT.

Theorem 3.1. Let (X, ρ, μ) be an SHT and let $1 < p < q < \infty$. Suppose that $0 < \alpha < 1/p$, $0 < \lambda_1 < 1 - \alpha p$ and $\lambda_2/q = \lambda_1/p$. Then K_α is bounded from $L^{p, \lambda_1}(X, \mu)$ to $L^{q, \lambda_2}(X, \nu)$ if and only if there is a positive constant c such that

$$\nu(B) \leq c\mu(B)^{q(1/p-\alpha)}, \tag{3}$$

for all balls B .

The next statement is a consequence of Theorem 3.1.

Theorem 3.2. Let (X, ρ, μ) be an SHT and let $1 < p < q < \infty$. Suppose that $0 < \alpha < 1/p$, $0 < \lambda_1 < 1 - \alpha p$ and $\lambda_2/q = \lambda_1/p$. Then for the boundedness of K_α from $L^{p, \lambda_1}(X, \mu)$ to $L^{q, \lambda_2}(X, \mu)$ it is necessary and sufficient that $q = p/(1 - \alpha p)$.

For non-homogeneous spaces we have the following statements:

Theorem 3.3. Let (X, ρ, μ) be a non-homogeneous space with the growth condition. Suppose that $1 < p \leq q < \infty$, $1/p - 1/q \leq \alpha < 1$ and $\alpha \neq 1/p$. Suppose also that $p\alpha - 1 < \beta < p - 1$, $0 < \lambda_1 < \beta - \alpha p + 1$ and $\lambda_1 q = \lambda_2 p$. Then I_α is bounded from $M_\beta^{p, \lambda_1}(X, \mu)$ to $M_\gamma^{q, \lambda_2}(X, \mu)$, where $\gamma = q(1/p + \beta/p - \alpha) - 1$.

Theorem 3.4. Suppose that (X, ρ, μ) is a quasimetric measure space and μ satisfies condition (2). Let $1 < p < q < \infty$. Assume that $0 < \alpha < 1$, $0 < \lambda_1 < p/q$ and $s\lambda_1/p = \lambda_2/q$. Then the operator I_α is bounded from $M^{p, \lambda_1 s}(X, \mu)$ to $M^{q, \lambda_2}(X, \mu)$.

4. Proof of the main results

In this section we give the proofs of the main results.

Proof of Theorem 3.1. Necessity: Suppose K_α is bounded from $L^{p, \lambda_1}(\mu)$ to $L^{q, \lambda_2}(X, \nu, \mu)$. Fix $B_0 := B(x_0, r_0)$. For $x, y \in B_0$, we have that

$$B(x, \rho(x, y)) \subseteq B(x, a_1(a_0 + 1)r_0) \subseteq B(x_0, a_1(1 + a_1(a_0 + 1))r_0).$$

Hence using the doubling condition for μ , it is easy to see that

$$\mu(B_0)^\alpha \leq c K_\alpha \chi_{B_0}(x), \quad x \in B_0.$$

Consequently, using the condition $\lambda_2/q = \lambda_1/p$, the boundedness of K_α from $L^{p,\lambda}(X, \mu)$ to $L^{q,\lambda_2}(X, \nu, \mu)$ and Lemma D we find that

$$\begin{aligned} \mu(B_0)^{\alpha-\lambda_1/p} \nu(B_0)^{1/q} &\leq c \|K_\alpha \chi_{B_0}\|_{L^{q,\lambda_2}(X,\nu,\mu)} \\ &\leq c \|\chi_{B_0}\|_{L^{p,\lambda_1}(X,\mu)} \leq c \mu(B_0)^{(1-\lambda_1)/p}. \end{aligned}$$

Since c does not depend on B_0 we have condition (3).

Sufficiency: Let $B := B(a, r)$, $\tilde{B} := B(a, 2a_1r)$ and $f \geq 0$. Write $f \in L^{p,\lambda_1}(\mu)$ as $f = f_1 + f_2 := f \chi_{\tilde{B}} + f \chi_{\tilde{B}^c}$, where χ_B is a characteristic function of B . Then we have

$$\begin{aligned} S &:= \int_B (K_\alpha f(x))^q d\nu(x) \leq c \left(\int_B (K_\alpha f_1(x))^q d\nu(x) + \int_B (K_\alpha f_2(x))^q d\nu(x) \right) \\ &:= c(S_1 + S_2). \end{aligned}$$

Applying Theorem F and the fact $\mu \in (DC)$ we find that

$$S_1 \leq \int_X (K_\alpha f_1)^q(x) d\nu(x) \leq c \left(\int_{B(a,2a_1r)} (f(x))^p d\mu(x) \right)^{q/p}.$$

Now observe that if $\rho(a, x) < r$ and $\rho(a, y) > 2a_1r$, then $\rho(a, y) > 2a_1\rho(a, x)$. Consequently, using the facts $\mu \in (RDC)$ (see (1)), $0 < \lambda_1 < 1 - \alpha p$ and condition (3) we have

$$\begin{aligned} S_2 &\leq c \int_{B(a,r)} \left(\int_{\rho(a,y)>r} \frac{f(y)}{\mu B(a, \rho(a, y))^{1-\alpha}} d\mu(y) \right)^q d\nu(x) \\ &= \nu(B) \left[\sum_{k=0}^{\infty} \int_{B(a,\eta_1^{k+1}r) \setminus B(a,\eta_1^k r)} \frac{f(y)}{\mu B(a, \rho(a, y))^{1-\alpha}} d\mu(y) \right]^q \\ &\leq c \nu(B) \left[\sum_{k=0}^{\infty} \left(\int_{B(a,\eta_1^{k+1}r)} (f(y))^p d\mu(y) \right)^{1/p} \right. \\ &\quad \left. \times \left(\int_{B(a,\eta_1^{k+1}r) \setminus B(a,\eta_1^k r)} \mu B(a, \rho(a, y))^{(\alpha-1)p'} d\mu(y) \right)^{1/p'} \right]^q \\ &\leq c \|f\|_{L^{p,\lambda_1}(X,\mu)}^q \nu(B) \left(\sum_{k=0}^{\infty} \mu B(a, \eta_1^{k+1}r)^{\lambda_1/p+\alpha-1+1/p'} \right)^q \\ &\leq c \|f\|_{L^{p,\lambda_1}(X,\mu)}^q \nu(B) \mu(B)^{(\lambda_1/p+\alpha-1/p)q} \left(\sum_{k=0}^{\infty} \eta_2^{k(\lambda_1/p+\alpha-1/p)} \right)^q \\ &\leq c \|f\|_{L^{p,\lambda_1}(X,\mu)}^q \mu(B)^{q\lambda_1/p} = c \|f\|_{L^{p,\lambda_1}(X,\mu)}^q \mu(B)^{\lambda_2}, \end{aligned}$$

where the positive constant c does not depend on B . Now the result follows immediately. \square

Proof of Theorem 3.2. *Sufficiency:* Assuming $\alpha = 1/p - 1/q$ and $\mu = \nu$ in Theorem 3.1 we have that K_α is bounded from $L^{p,\lambda_1}(X, \mu)$ to $L^{q,\lambda_2}(X, \mu)$.

Necessity: Suppose that K_α is bounded from $L^{p,\lambda_1}(X, \mu)$ to $L^{q,\lambda_2}(X, \mu)$. Then by Theorem 3.1 we have

$$\mu(B)^{1/q-1/p+\alpha} \leq c.$$

The conditions $\mu(X) = \infty$ and $\mu\{x\} = 0$, for all $x \in X$, implies that $\alpha = 1/p - 1/q$. \square

Proof of Theorem 3.3. Let $f \geq 0$. For $x, a \in X$, let us introduce the following notation:

$$E_1(x) := \left\{ y : \frac{\rho(a, y)}{\rho(a, x)} < \frac{1}{2a_1} \right\}, \quad E_2(x) := \left\{ y : \frac{1}{2a_1} \leq \frac{\rho(a, y)}{\rho(a, x)} \leq 2a_1 \right\},$$

$$E_3(x) := \left\{ y : 2a_1 < \frac{\rho(a, y)}{\rho(a, x)} \right\}.$$

For $i = 1, 2, 3, r > 0$ and $a \in X$, we denote

$$S_i := \int_{\rho(a,x) < r} \rho(a, x)^\gamma \left(\int_{E_i(x)} f(y) \rho(x, y)^{\alpha-1} d\mu(y) \right)^q d\mu(x).$$

If $y \in E_1(x)$, then $\rho(a, x) < 2a_1 a_0 \rho(x, y)$. Hence, it is easy to see that

$$S_1 \leq C \int_B \rho(a, x)^{\gamma+q(\alpha-1)} \left(\int_{\rho(a,y) < \rho(a,x)} f(y) d\mu(y) \right)^q d\mu(x).$$

Taking into account the condition $\gamma < (1 - \alpha)q - 1$ we have

$$\int_{\rho(a,x) > t} \rho(a, x)^{\gamma+q(\alpha-1)} d\mu(x) = \sum_{n=0}^{\infty} \int_{B(a, 2^{k+1}t) \setminus B(a, 2^k t)} (\rho(a, x))^{\gamma+q(\alpha-1)q} d\mu(x)$$

$$\leq c \sum_{n=0}^{\infty} (2^k t)^{\gamma+q(\alpha-1)+1} = c t^{\gamma+q(\alpha-1)+1},$$

while the condition $\beta < p - 1$ implies

$$\int_{\rho(a,x) < t} \rho(a, x)^{\beta(1-p')+1} d\mu(x) \leq c t^{\beta(1-p')+1}.$$

Hence

$$\sup_{a \in X, t > 0} \left(\int_{\rho(a,x) > t} \rho(a, x)^{\gamma+q(\alpha-1)} d\mu(x) \right)^{1/q} \left(\int_{B(a,t)} \rho(a, y)^{\beta(1-p')} d\mu(y) \right)^{1/p'}$$

$$< \infty.$$

Now using Theorem C we have

$$S_1 \leq c \left(\int_B \rho(a, x)^\beta (f(y)) d\mu(y) \right)^{q/p} \leq c \|f\|_{M_\beta^{p,\lambda_1}(X,\mu)}^q r^{\lambda_1 q/p} = c \|f\|_{M_\beta^{p,\lambda_1}(X,\mu)}^q r^{\lambda_2}.$$

Further, observe that if $\rho(a, y) > 2a_1\rho(a, x)$, then $\rho(a, y) \leq a_1\rho(a, x) + a_1\rho(a, y) \leq \rho(a, y)/2 + a_1\rho(a, x)$. Hence $\rho(a, y)/(2a_1) \leq \rho(a, x)$. Consequently, using the growth condition for μ , the fact $\lambda_1 < \beta - \alpha p + 1$ and Lemma E we find that

$$\begin{aligned}
 S_3 &\leq c \int_{B(a,r)} \rho(a, x)^\gamma \left(\int_{\rho(a,y) > \rho(a,x)} \frac{f(y)}{\rho(a, y)^{1-\alpha}} d\mu(y) \right)^q d\mu(x) \\
 &\leq c \int_{B(a,r)} \rho(a, x)^\gamma \left(\sum_{k=0}^{\infty} \int_{B(a, 2^{k+1}\rho(a,x)) \setminus B(a, 2^k\rho(a,x))} \frac{f(y)}{\rho(a, y)^{1-\alpha}} d\mu(y) \right)^q d\mu(x) \\
 &\leq c \int_{B(a,r)} \rho(a, x)^\gamma \left[\sum_{k=0}^{\infty} \left(\int_{B(a, 2^{k+1}\rho(a,x))} f^p(y) \rho(a, y)^\beta d\mu(y) \right)^{1/p} \right. \\
 &\quad \left. \times \left(\int_{B(a, 2^{k+1}\rho(a,x)) \setminus B(a, 2^k\rho(a,x))} \rho(a, y)^{\beta(1-p') + (\alpha-1)p'} d\mu(y) \right)^{1/p'} \right]^q d\mu(x) \\
 &\leq c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q \int_{B(a,r)} \rho(a, x)^\gamma \\
 &\quad \times \left(\sum_{k=0}^{\infty} (2^k \rho(a, x))^{\lambda_1/p + \alpha - 1 - \beta/p} (\mu B(a, 2^{k+1}\rho(a, x)))^{1/p'} \right)^q d\mu(x) \\
 &\leq c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q \int_{B(a,r)} \rho(a, x)^\gamma \left(\sum_{k=0}^{\infty} (2^k \rho(a, x))^{\lambda_1/p + \alpha - 1 - \beta/p} \right)^q d\mu(x) \\
 &\leq c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q \int_{B(a,r)} \rho(a, x)^{(\lambda_1/p + \alpha - 1 - \beta/p)q + \gamma} d\mu(x) \\
 &= c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q \int_{B(a,r)} \rho(a, x)^{\lambda_1 q/p - 1} d\mu(x) \leq c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q r^{\lambda_1 q/p} \\
 &= c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q r^{\lambda_2}.
 \end{aligned}$$

So, we conclude that

$$S_3 \leq c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q r^{\lambda_2}.$$

To estimate S_2 we consider two cases. First assume that $\alpha < 1/p$. Let

$$E_{k,r} := \{x : 2^k r \leq \rho(a, x) < 2^{k+1} r\},$$

$$F_{k,r} := \{x : 2^{k-1} r/a_1 \leq \rho(a, x) < a_1 2^{k+2} r\}.$$

Assume that $p^* = p/(1 - \alpha p)$. By Hölder's inequality, Corollary B and the assumption $\gamma = q(1/p + \beta/p - \alpha) - 1$ we have

$$\begin{aligned}
 S_2 &= \sum_{k=-\infty}^{-1} \int_{E_{k,r}} \rho(a, x)^\gamma \left(\int_{E_2(x)} f(y) \rho(x, y)^{\alpha-1} d\mu(y) \right)^q d\mu(x) \\
 &\leq \sum_{k=-\infty}^{-1} \left(\int_{E_{k,r}} \rho(a, x)^\gamma \left(\int_{E_2(x)} f(y) \rho(x, y)^{\alpha-1} d\mu(y) \right)^{p^*} d\mu(x) \right)^{q/p'} \\
 &\quad \times \left(\int_{E_{k,r}} \rho(a, x)^{\gamma p^*/(p^*-q)} d\mu(x) \right)^{(p^*-q)/p^*} \\
 &\leq c \sum_{k=-\infty}^{-1} 2^{k(\gamma+(p^*-q)/p^*)} \left(\int_{F_{k,r}} I_\alpha(f \chi_{F_{k,r}})(x)^{p^*} d\mu(x) \right)^{q/p'} \\
 &\leq c \sum_{k=-\infty}^{-1} 2^{k(\gamma+(p^*-q)/p^*)} \left(\int_{F_{k,r}} (f(x))^p d\mu(x) \right)^{q/p} \\
 &\leq c \left(\int_{B(a, 2a_1 r)} \rho(a, x)^\beta (f(x))^p d\mu(x) \right)^{q/p} \\
 &\leq c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q r^{\lambda_1 q/p} = c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q r^{\lambda_2}.
 \end{aligned}$$

Let us now consider the case $1/p < \alpha < 1$.

First notice that (see [13])

$$\int_{E_2(x)} (\rho(x, y))^{(\alpha-1)p'} d\mu(y) \leq c \rho(a, x)^{1+(\alpha-1)p'},$$

where the positive constant c does not depend on a and x .

This estimate and Hölder's inequality yield

$$\begin{aligned}
 S_2 &\leq c \sum_{k=-\infty}^{-1} \left(\int_{E_{k,r}} \rho(a, x)^{\gamma+[(\alpha-1)p'+1]q/p'} \left(\int_{E_2(x)} (f(y))^p d\mu(y) \right)^{q/p} d\mu(x) \right)^{q/p'} \\
 &\leq c \sum_{k=-\infty}^{-1} \left(\int_{E_{k,r}} \rho(a, x)^{\gamma+[(\alpha-1)p'+1]q/p'} d\mu(x) \right) \left(\int_{F_{k,r}} (f(y))^p d\mu(y) \right)^{q/p} \\
 &\leq c \sum_{k=-\infty}^{-1} (2^k r)^{\gamma+[(\alpha-1)p'+1]q/p'+1} \left(\int_{F_{k,r}} (f(y))^p d\mu(y) \right)^{q/p} \\
 &= c \sum_{k=-\infty}^{-1} 2^{k\beta q/p} \left(\int_{F_{k,r}} (f(y))^p d\mu(y) \right)^{q/p} \leq c \left(\int_{B(a, 2a_1 r)} (f(y))^p \rho(a, y)^\beta d\mu(y) \right)^{q/p} \\
 &\leq c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q r^{\lambda_1 q/p} = c \|f\|_{M_\beta^{p, \lambda_1}(X, \mu)}^q r^{\lambda_2}.
 \end{aligned}$$

Now the result follows immediately. \square

Proof of Theorem 3.4. Let $f \geq 0$. Suppose that $a \in X$ and $r > 0$. Suppose also that $f_1 = f \chi_{B(a, 2a_1r)}$ and $f_2 = f - f_1$. Then $I_\alpha f = I_\alpha f_1 + I_\alpha f_2$. Consequently,

$$\begin{aligned} \int_{B(a,r)} (I_\alpha f(x))^q d\mu(x) &\leq 2^{q-1} \left(\int_{B(a,r)} (I_\alpha f_1(x))^q d\mu(x) + \int_{B(a,r)} (I_\alpha f_2(x))^q d\mu(x) \right) \\ &:= 2^{q-1} (S_{a,r}^{(1)} + S_{a,r}^{(2)}). \end{aligned}$$

Due to Theorem A and the condition $s\lambda_1/p = \lambda_2/q$ we have

$$\begin{aligned} S_{a,r}^{(1)} &\leq c \left(\int_{B(a, 2a_1r)} (f(x))^p d\mu(x) \right)^{q/p} \\ &= c \left(\frac{1}{(2a_1r)^{\lambda_1 s}} \int_{B(a, 2a_1r)} (f(x))^p dx \right)^{q/p} r^{\lambda_1 s q/p} \leq c \|f\|_{M^{p, \lambda_1 s}(X, \mu)}^q r^{\lambda_2}. \end{aligned}$$

Now observe that if $x \in B(a, r)$ and $y \in X \setminus B(a, 2a_1r)$, then $\rho(a, y)/2a_1 \leq \rho(x, y)$. Hence Hölder's inequality, condition (2) and the condition $0 < \lambda_1 < p/q$ yield

$$\begin{aligned} I_\alpha f_2(x) &= \int_{X \setminus B(a, 2a_1r)} f(y)/\rho(x, y)^{1-\alpha} d\mu(y) \\ &= \sum_{k=0}^{\infty} \left(\int_{B(a, 2^{k+2}a_1r) \setminus B(a, 2^{k+1}a_1r)} (f(y))^p d\mu(y) \right)^{1/p} \\ &\quad \times \left(\int_{B(a, 2^{k+2}a_1r) \setminus B(a, 2^{k+1}a_1r)} \rho(a, y)^{(\alpha-1)p'} d\mu(y) \right)^{1/p'} \\ &\leq c \sum_{k=0}^{\infty} \left(\frac{1}{(2^{k+1}a_1r)^{\lambda_1 s}} \int_{B(a, 2^{k+1}a_1r)} (f(y))^p d\mu(y) \right)^{1/p} \\ &\quad \times (2^k a_1r)^{\lambda_1 s/p + \alpha - 1 + s/p'} \\ &\leq c \|f\|_{M^{p, \lambda_1 s}(X, \mu)}^q r^{\lambda_1 s/p + \alpha - 1 + s/p'}. \end{aligned}$$

Consequently, by the assumptions $s\lambda_1/p = \lambda_2/q$ and $s = pq(1 - \alpha)/(pq + p - q)$ we conclude that

$$S_{a,r}^{(2)} \leq c \|f\|_{M^{p, \lambda_1 s}(X, \mu)}^q r^{(\lambda_1 s/p + \alpha - 1 + s/p')q + s} = c \|f\|_{M^{p, \lambda_1 s}(X, \mu)}^q r^{\lambda_2}.$$

Summarizing the estimates derived above we finally have the desired result. \square

Acknowledgements

The second and third authors were partially supported by the INTAS Grant no. 05-1000008-8157 and the Georgian National Science Foundation Grant no. GNSF/ST07/3-169.

The authors express their gratitude to the referee and Professor Christian Berg for their remarks and suggestions.

References

- [1] D.R. Adams, A trace inequality for generalized potentials, *Studia Math.* 48 (1973) 99–105.
- [2] D.R. Adams, A note on Riesz potentials, *Duke Math. J.* 42 (4) (1975) 765–778.
- [3] V. Burenkov, H.V. Guliyev, Necessary and sufficient conditions for boundedness of the maximal operator in local Morrey-type spaces, *Studia Math.* 163 (2) (2004) 157–176.
- [4] I.V. Burenkov, H.V. Guliyev, V.S. Guliyev, Necessary and sufficient conditions for the boundedness of the fractional maximal operator in local Morrey-type spaces, *Dokl. Akad. Nauk* 409 (4) (2006) 443–447 (in Russian).
- [5] D. Edmunds, V. Kokilashvili, A. Meskhi, *Bounded and Compact Integral Operators, Mathematics and its Applications*, vol. 543, Kluwer Academic Publishers, Dordrecht, Boston, London, 2002.
- [6] Eridani, V. Kokilashvili, A. Meskhi, Morrey spaces and fractional integral operators, Preprint No. 65, School of Mathematical Sciences, GC University, Lahore, 2007.
- [7] J. García-Cuerva, A.E. Gatto, Boundedness properties of fractional integral operators associated to non-doubling measures, *Studia Math.* 162 (3) (2004) 245–261.
- [8] J. García-Cuerva, J.M. Martell, Two-weight norm inequalities for maximal operators and fractional integrals on non-homogeneous spaces, *Indiana Univ. Math. J.* 50 (3) (2001) 1241–1280.
- [9] I. Genebashvili, A. Gogatishvili, V. Kokilashvili, Solution of two-weight problems for integral transforms with positive kernels, *Georgian Math. J.* 3 (1) (1996) 319–342.
- [10] I. Genebashvili, A. Gogatishvili, V. Kokilashvili, M. Krbec, *Weight Theory for Integral Transforms on Spaces of Homogeneous Type*, Pitman Monographs and Surveys in Pure and Applied Mathematics, vol. 92, Longman, Harlow, 1998.
- [11] V. Kokilashvili, Weighted estimates for classical integral operators, in: *Nonlinear Analysis, Function Spaces and Applications IV*, Roudnice nad Labem, vol. 119, Teubner-Texte Math, Leipzig, 1990, pp. 86–103.
- [12] V. Kokilashvili, A. Meskhi, Fractional integrals on measure spaces, *Frac. Calc. Appl. Anal.* 4 (4) (2001) 1–24.
- [13] V. Kokilashvili, A. Meskhi, On some weighted inequalities for fractional integrals on non-homogeneous spaces, *Z. Anal. Anwend.* 24 (4) (2005) 871–885.
- [14] Y. Komori, T. Mizuhara, Notes on commutators and Morrey spaces, *Hokkaido Math. J.* 32 (2003) 345–353.
- [15] J. Peetre, On the theory of $\mathcal{L}^{p,\lambda}$ spaces, *J. Funct. Anal.* 4 (1969) 71–87.
- [16] S. Samko, A.A. Kilbas, O.I. Marichev, *Fractional Integrals and Derivatives. Theory and Applications*, Gordon and Breach, London, 1993.
- [17] Y. Sawano, H. Tanaka, Morrey spaces for nondoubling measures, *Acta Math. Sinica* 21 (6) (2005) 1535–1544.
- [18] E.M. Stein, G. Weiss, Fractional integrals on n -dimensional Euclidean spaces, *J. Math. Mech.* 7 (4) (1958) 503–514.
- [19] J.O. Strömberg, A. Torchinsky, *Weighted Hardy Spaces*, Lecture Notes in Mathematics, vol. 1381, Springer, Berlin, 1989.

Morrey spaces and fractional integral operators

ORIGINALITY REPORT

23%

SIMILARITY INDEX

17%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

www.emis.de

Internet Source

2%

2

w3.ualg.pt

Internet Source

1%

3

Shi, Yanlong, and Xiangxing Tao. "The boundedness of sublinear operators on Morrey-Herz spaces over the homogeneous type space", *Analysis Mathematica*, 2013.

Publication

1%

4

Usman Ashraf. "Boundedness and compactness of positive integral operators on cones of homogeneous groups", *Positivity*, 10/28/2008

Publication

1%

5

fac.ksu.edu.sa

Internet Source

1%

6

dualibra.com

Internet Source

1%

7

A. Meskhi. "CRITERIA FOR THE

BOUNDEDNESS AND COMPACTNESS OF INTEGRAL TRANSFORMS WITH POSITIVE KERNELS", Proceedings of the Edinburgh Mathematical Society, 06/2001

Publication

1%

8

Loukas Grafakos. "Maximal function characterizations of Hardy spaces on RD-spaces and their applications", Science in China Series A Mathematics, 12/2008

Publication

1%

9

tel.archives-ouvertes.fr

Internet Source

1%

10

Detlef Müller, Dachun Yang. "A difference characterization of Besov and Triebel-Lizorkin spaces on RD-spaces", Forum Mathematicum, 2009

Publication

1%

11

archive.org

Internet Source

1%

12

epubl.luth.se

Internet Source

1%

13

Cichocki, Andrzej, and Shun-ichi Amari. "Families of Alpha- Beta- and Gamma-Divergences: Flexible and Robust Measures of Similarities", Entropy, 2010.

Publication

1%

14	ddd.uab.cat Internet Source	1%
15	www.heldermann-verlag.de Internet Source	1%
16	V. S. Guliyev. "Boundedness of maximal, potential type, and singular integral operators in the generalized variable exponent Morrey type spaces", Journal of Mathematical Sciences, 10/2010 Publication	1%
17	www.mat.ucm.es Internet Source	1%
18	Mubariz G. Hajibayov. "Generalized potentials in variable exponent Lebesgue spaces on homogeneous spaces", Mathematische Nachrichten, 01/2011 Publication	<1%
19	Universitext, 2015. Publication	<1%
20	www.docme.ru Internet Source	<1%
21	www.fmi-plovdiv.org Internet Source	<1%
22	gking.harvard.edu Internet Source	<1%

23 arxiv.org Internet Source <1%

24 journal.taiwanmathsoc.org.tw Internet Source <1%

25 **GUOEN HU. "WEIGHTED ESTIMATES FOR SINGULAR INTEGRAL OPERATORS WITH NONSMOOTH KERNELS AND APPLICATIONS", Journal of the Australian Mathematical Society, 12/2008** Publication <1%

26 users.minet.uni-jena.de Internet Source <1%

27 **David R. Adams. "Morrey Potentials and Harmonic Maps", Communications in Mathematical Physics, 09/07/2011** Publication <1%

28 azjm.azvs.az Internet Source <1%

29 www.scm.org.co Internet Source <1%

30 mipa.unimes.fr Internet Source <1%

31 hal.archives-ouvertes.fr Internet Source <1%

docplayer.com.br

32

Internet Source

<1%

33

hpep3.eps.s.u-tokyo.ac.jp

Internet Source

<1%

34

Fofana, Ibrahim, Falé Roland Faléa, and Bérenger Akon Kpata. "A class of subspaces of Morrey spaces and norm inequalities on Riesz potential operators", Afrika Matematika, 2015.

Publication

<1%

35

nkiso.u-tokai.ac.jp

Internet Source

<1%

36

Mubariz G. Hajibayov. "Weighted Estimates of Generalized Potentials in Variable Exponent Lebesgue Spaces on Homogeneous Spaces", Recent Trends in Toeplitz and Pseudodifferential Operators, 2010

Publication

<1%

37

Temlyakov, Vladimir. "Appendix", Advanced Courses in Mathematics - CRM Barcelona, 2015.

Publication

<1%

38

www.uam.es

Internet Source

<1%

39

epdf.tips

Internet Source

<1%

40	cat4mba.com Internet Source	<1%
41	www.archive.org Internet Source	<1%
42	D. E. Edmunds. <i>Mathematische Nachrichten</i> , 06/2003 Publication	<1%
43	www.anstuocmath.ro Internet Source	<1%
44	www.ams.org Internet Source	<1%
45	users.uoa.gr Internet Source	<1%
46	orbilu.uni.lu Internet Source	<1%
47	vdocuments.mx Internet Source	<1%
48	mai.liu.se Internet Source	<1%
49	manualzz.com Internet Source	<1%
50	www.mth.kcl.ac.uk Internet Source	<1%

51

fsdona2011.uni-jena.de

Internet Source

<1%

52

Datuashvili, T.. "CATEGORICAL,
HOMOLOGICAL, AND HOMOTOPICAL
PROPERTIES OF ALGEBRAIC OBJECTS.",
Journal of Mathematical Sciences

Publication

<1%

53

Victor I. Burenkov. "Necessary and Sufficient
Conditions for the Boundedness of the Riesz
Potential in Local Morrey-type Spaces",
Potential Analysis, 04/2009

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Morrey spaces and fractional integral operators

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14
