

# CHARACTERIZATIONS FOR THE GENERALIZED FRACTIONAL INTEGRAL OPERATORS ON MORREY SPACES

*by* Eridani Eridani

---

**Submission date:** 13-Dec-2018 11:04AM (UTC+0800)

**Submission ID:** 1056162363

**File name:** CHARACTERIZATIONS\_FOR\_THE\_GENERALIZED.pdf (287.13K)

**Word count:** 6917

**Character count:** 28263

67

## CHARACTERIZATIONS FOR THE GENERALIZED FRACTIONAL INTEGRAL OPERATORS ON MORREY SPACES

ERIDANI, HENDRA GUNAWAN, EIICHI NAKAI AND YOSHIHIRO SAWANO

(Communicated by B. Opic)

50

*Abstract.* We present some characterizations for the boundedness of the generalized fractional integral operators on Morrey spaces. The characterizations follow from two key estimates, one for the norm of some functions in Morrey spaces, and another for the values of the corresponding fractional integrals. We prove three theorems about necessary and sufficient conditions. We show that these theorems are independent by giving some examples. We also obtain counterparts for the weak generalized Morrey spaces.

### 1. Introduction

95 In this paper, for a measurable function  $\rho : (0, \infty) \rightarrow (0, \infty)$ , we are interested in the generalized fractional integral operator  $I_\rho$  given by the formula

$$24 \quad I_\rho f(x) := \int_{\mathbb{R}^d} \frac{\rho(|x-y|)}{|x-y|^d} f(y) dy, \quad x \in \mathbb{R}^d,$$

for any suitable function  $f$  on  $\mathbb{R}^d$ . This generalized fractional integral operator was initially investigated in [27]. Nowadays many authors have been culminating important observations about  $I_\rho$  especially in connection with Morrey spaces. These spaces cover Lebesgue spaces as special cases and seem to describe the behavior of  $I_\rho$  well. In order to highlight what we shall prove in this paper, we take up the works [3, 6, 7, 18, 22, 25, 28, 35], where we formulated sufficient conditions on  $\rho$  for  $I_\rho$  to be bounded on Morrey spaces  $L_{p,\phi}$  with  $1 \leq p < \infty$  and  $\phi$  a function from  $(0, \infty)$  to itself. We aim to show that these conditions are necessary as well. We characterize the boundedness by estimating the norm of the characteristic functions of balls and the function  $\phi(|\cdot|)$ , as well as the value of the corresponding fractional integrals.

Hereafter, we assume that

$$\int_0^1 \frac{\rho(s)}{s} ds < \infty,$$

30

---

*Mathematics subject classification* (2010): Primary 42B35; Secondary 26A33, 46E30, 42B20, 43A15.  
*Keywords and phrases:* Fractional integrals, bounded operators, Morrey spaces.

so that the fractional integrals  $I_\rho f$  are well-defined, at least for characteristic functions of balls. In addition, we shall also assume that  $\rho$  satisfies the *growth condition*: there exist constants  $C_1 > 0$  and  $0 < 2k_1 < k_2 < \infty$  such that

$$\sup_{r/2 < s \leq r} \rho(s) \leq C_1 \int_{k_1 r}^{k_2 r} \frac{\rho(s)}{s} ds, \quad r > 0. \tag{1.1}$$

This condition is weaker than the usual *doubling condition*: there exists a constant  $C_2 > 0$  such that

$$\frac{1}{C_2} \leq \frac{\rho(r)}{\rho(s)} \leq C_2$$

whenever  $r$  and  $s$  satisfy

$$r, s > 0 \text{ and } \frac{1}{2} \leq \frac{r}{s} \leq 2.$$

See [40] for some examples and more explanation about these two conditions.

In the present paper we work on generalized Morrey spaces. For a certain function  $\phi : (0, \infty) \rightarrow (0, \infty)$ , we say that a function  $f$  belongs to the generalized Morrey space  $L_{p,\phi} = L_{p,\phi}(\mathbb{R}^d)$ , where  $1 \leq p < \infty$ , if

$$\|f\|_{L_{p,\phi}} := \sup_{a \in \mathbb{R}^d, r > 0} \frac{1}{\phi(r)} \left[ \frac{1}{|B(a,r)|} \int_{B(a,r)} |f(x)|^p dx \right]^{1/p} < \infty.$$

Note that if  $\phi(r) := r^{(\lambda-d)/p}$  for some  $1 \leq p < \infty$  and  $0 \leq \lambda < d$ , then  $L_{p,\phi}(\mathbb{R}^d) = L_{p,\lambda}(\mathbb{R}^d)$ , see (1.8) below. In [26, p. 446] we justified that  $\phi$  is a nonincreasing function such that  $t \mapsto \phi(t)^p t^d$  is a nondecreasing for  $L_{p,\phi}(\mathbb{R}^d) \neq \{0\}$ . We refer to [25, 29, 32] and [40, Section 1] for more information about these spaces.

Here we shall assume that  $\phi : (0, \infty) \rightarrow (0, \infty)$  is *almost decreasing* [that is, if  $r \leq s$ , then  $\phi(r) \geq C_3 \phi(s)$ ], and that  $r^d \phi^p(r)$  is *almost increasing*, [that is, if  $r \leq s$ , then  $r^d \phi^p(r) \leq C'_3 s^d \phi^p(s)$ ]. These two conditions implies that  $\phi$  also satisfies the *doubling condition*. Denote by  $\mathcal{G}_p$  the set of all functions  $\phi : (0, \infty) \rightarrow (0, \infty)$  such that  $\phi$  is almost decreasing and that  $r \mapsto r^{d/p} \phi(r)$  is almost increasing. Now we present three different criteria for the boundedness of  $I_\rho$ . For convenience, write  $\tilde{\rho}(r) := \int_0^r \frac{\rho(t)}{t} dt$ . We prove the following theorems about the boundedness of  $I_\rho$  on generalized Morrey spaces.

**THEOREM 1.1.** *Let  $1 < p < q < \infty$  and  $\phi \in \mathcal{G}_p$ . Assume*

$$\int_r^\infty \frac{\phi(s)\rho(s)}{s} ds \leq C \phi(r)\rho(r) \quad (r > 0) \tag{1.2}$$

*for some constant  $C > 0$ . Then  $I_\rho$  is bounded from  $L_{p,\phi}(\mathbb{R}^d)$  to  $L_{q,\phi^{p/q}}(\mathbb{R}^d)$  if and only if there exists a constant  $C' > 0$  such that*

$$\tilde{\rho}(r) \leq C' \phi(r)^{p/q-1} \quad (r > 0). \tag{1.3}$$

2 THEOREM 1.2. Let  $1 < p < q < \infty$  and let  $\phi \in \mathcal{G}_p$ . Assume

$$\int_0^r \frac{\phi(s)s^{d/p}}{s} ds \leq C\phi(r)r^{d/p} \quad (r > 0) \tag{1.4}$$

for some constant  $C > 0$ . Then  $I_p$  is bounded from  $L_{p,\phi}(\mathbb{R}^d)$  to  $L_{q,\phi^{p/q}}(\mathbb{R}^d)$  if and only if there exists a constant  $C' > 0$  such that

$$\phi(r) \int_0^r \frac{\rho(t)}{t} dt + \int_r^\infty \frac{\phi(t)\rho(t)}{t} dt \leq C'\phi(r)^{p/q} \quad (r > 0). \tag{1.5}$$

THEOREM 1.3. Let  $\phi, \psi \in \mathcal{G}_1$ . Assume that

$$\int_0^r \frac{\phi(s)s^d}{s} ds \leq C\phi(r)r^d \quad (r > 0) \tag{1.6}$$

for some constant  $C > 0$ . Then  $I_p$  is bounded from  $L_{1,\phi}(\mathbb{R}^d)$  to  $L_{1,\psi}(\mathbb{R}^d)$  if and only if there exists a constant  $C' > 0$  such that

$$\phi(r) \int_0^r \frac{\rho(t)}{t} dt + \int_r^\infty \frac{\phi(t)\rho(t)}{t} dt \leq C'\psi(r) \quad (r > 0). \tag{1.7}$$

81 Note that the integral operators such as  $(1 - \Delta)^{-\alpha}$  and  $L^{-\alpha}$ , where  $L$  is a suitable elliptic differential operator and  $\alpha > 0$ , fall under this scope. Also, if a measurable function  $V : \mathbb{R}^d \rightarrow (0, \infty)$  satisfies the reverse Hölder inequality, that is, there exist some constants  $C > 0, q \gg 1$  such that, for all balls  $B$ ,  $\left(\int_B V(x)^q \frac{dx}{|B|}\right)^{1/q} \leq C \int_B V(x) \frac{dx}{|B|}$ , then the operators  $V^\gamma(-\Delta + V)^{-\beta}$  with  $0 \leq \gamma \leq \beta \leq 1$  and  $V^\gamma \partial_j(-\Delta + V)^{-\beta}$  with  $0 \leq \gamma \leq \frac{1}{2} \leq \beta \leq 1, \beta - \gamma \geq \frac{1}{2}$  and  $j = 1, 2, \dots, d$  also fall under this scope [20]. We refer to [13, Sections 3 and 4] for a detailed description of these facts.

A few remarks concerning the conditions on the theorems may be in order.

REMARK 1.4.

- (i) Theorems 1.1-1.3 extend those obtained in [7] where the authors considered the classical Riesz potential.
- (ii) In Theorem 1.2, to prove the sufficiency, there is no need to assume (1.4).
- (iii) The condition (1.5) appeared in [18] originally and it later appeared in a bilinear estimate of the form  $g \cdot I_\alpha f$  (see [39, Theorem 1.6]).
- (iv) In Theorem 1.3, to prove the sufficiency, there is no need to assume (1.6).
- (v) It follows immediately that the right-hand side of (1.7) equals

$$\phi(r) \int_0^r \frac{\rho(t)}{t} dt + \int_r^\infty \frac{\phi(t)\rho(t)}{t} dt = \int_0^\infty \frac{\phi(\max(r,t))\rho(t)}{t} dt. \tag{1.7}$$

- (vi) The condition (1.7) is known to be sufficient in [28, Theorem 3.2].

41

Our results can be readily transplanted into those for Morrey spaces and Lebesgue spaces. For  $1 \leq p < \infty$  and  $0 \leq \lambda < d$ , recall that the Morrey space  $L^{p,\lambda} = L^{p,\lambda}(\mathbb{R}^d)$  consists of all locally integrable functions  $f$  on  $\mathbb{R}^d$  for which

$$\|f : L^{p,\lambda}\| := \sup_{a \in \mathbb{R}^d, r > 0} \left[ \frac{1}{r^\lambda} \int_{B(a,r)} |f(x)|^p dx \right]^{\frac{1}{p}} < \infty. \tag{1.8}$$

See [33] for more information about these spaces. Observe that, with norm coincidence,  $L^{p,0}(\mathbb{R}^d) = L^p(\mathbb{R}^d)$  for  $1 \leq p < \infty$ . We note that if  $\rho(r) = r^\alpha$ , with  $0 < \alpha < d$ , then  $I_\rho = I_\alpha$  is the classical fractional integral operator, also known as the Riesz potential, which is bounded from  $L^p(\mathbb{R}^d)$  to  $L^q(\mathbb{R}^d)$  if and only if  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{d}$ , where  $1 < p, q < \infty$  [43]. The necessary part is usually proved by using the scaling arguments. See [6, 18, 19, 46] for some recent results on the boundedness properties of  $I_\rho$ .

6

Theorems 1.1–1.2 both characterize the kernel function  $\rho$  for which  $I_\rho$  is bounded from  $L^p(\mathbb{R}^d)$  to  $L^q(\mathbb{R}^d)$  for  $1 < p < q < \infty$ .

46

**COROLLARY 1.5.** *Let  $1 < p < q < \infty$ . The operator  $I_\rho$  is bounded from  $L^p(\mathbb{R}^d)$  to  $L^q(\mathbb{R}^d)$  if and only if  $\rho(r) \leq Cr^{d(1/p-1/q)}$  for all  $r > 0$ .*

For  $\rho(r) = r^\alpha$ , Corollary 1.5 reads that the operator  $I_\rho$  is bounded from  $L^p(\mathbb{R}^d)$  to  $L^q(\mathbb{R}^d)$  if and only if  $\frac{\alpha}{d} = \frac{1}{p} - \frac{1}{q}$ , where  $1 < p < q < \infty$ .

With Theorems 1.1–1.3 we can characterize the function  $\rho$  for which  $I_\rho$  is bounded from one Morrey space to another.

The next corollary generalizes the previous characterization in Corollary 1.5.

80

**COROLLARY 1.6.** *Let  $1 < p < q < \infty$  and  $0 \leq \lambda < d$ . Assume that  $\rho$  satisfies (1.1). Then the operator  $I_\rho$  is bounded from  $L^{p,\lambda}(\mathbb{R}^d)$  to  $L^{q,\lambda}(\mathbb{R}^d)$  precisely when one of the following equivalent conditions holds.*

- (a)  $\rho(r) \leq Cr^{(d-\lambda)(1/p-1/q)}$  for all  $r > 0$ .
- (b)  $\tilde{\rho}(r) = \int_0^r \frac{\rho(s)}{s} ds \leq Cr^{(d-\lambda)(1/p-1/q)}$  for all  $r > 0$ .

In Sugano’s modified setting, these conditions corresponds to [45, the formula (18)], for example.

Generalized Morrey spaces occur naturally. We give a proposition, which improves [41, Theorem 5.1]. We write

$$\ell_{-1,0}(r) := \begin{cases} \log \frac{1}{r} & (0 < r < e^{-1}), \\ 1 & (e^{-1} \leq r). \end{cases}$$

**PROPOSITION 1.7.** *Let  $s \in (0, d)$ . Define*

$$\psi(r) := \frac{r^d \ell_{-1,0}(r)}{(1+r)^s} \quad (r > 0).$$

19 Then there exists a constant  $C_s > 0$  such that

$$\|f : L_{1,\psi}\| \leq C_s \|(1 - \Delta)^{s/2} f : L^{1,d-s}\|$$

holds for all  $f \in L^{1,d-s}(\mathbb{R}^d)$  with  $(1 - \Delta)^{s/2} f \in L^{1,d-s}(\mathbb{R}^d)$ .

The proof will be given by Example 5 below. Note that we cannot delete  $\ell_{-1,0}$  because it is a necessary condition for this estimate as Example 5 shows.

A passage of Theorems 1.1, 1.2 and 1.3 from generalized Morrey spaces to weak generalized Morrey spaces can be readily done. For  $p \in [1, \infty)$  and  $\phi : (0, \infty) \rightarrow (0, \infty)$ , recall that  $L_{p,\phi,\text{weak}} = L_{p,\phi,\text{weak}}(\mathbb{R}^d)$  is the set of all functions such that

$$\|f : L_{p,\phi,\text{weak}}\| := \sup_{a \in \mathbb{R}^n, r > 0} \frac{1}{\phi(r)} \left( \frac{\sup_{t > 0} t^p m(B(a,r), f, t)}{|B(a,r)|} \right)^{1/p} < \infty, \tag{1.9}$$

where

$$m(B(a,r), f, t) := |\{x \in B(a,r) : |f(x)| > t\}|.$$

We remark that some prefer to use  $M_{p,\phi}(\mathbb{R}^n)$  instead of  $L_{p,\phi}(\mathbb{R}^n)$  and that, likewise, some prefer to use  $WM_{p,\phi}(\mathbb{R}^n)$  instead of  $L_{p,\phi,\text{weak}}(\mathbb{R}^n)$ .

We have counterparts of three theorems above.

79 THEOREM 1.8. Let  $1 \leq p < q < \infty$  and let  $\phi \in \mathcal{G}_p$ . Assume (1.2). Then  $I_\rho$  is bounded from  $L_{p,\phi}(\mathbb{R}^d)$  to  $L_{q,\phi^{p/q},\text{weak}}(\mathbb{R}^d)$  if and only if (1.3) holds. 6

1 THEOREM 1.9. Let  $1 \leq p < q < \infty$  and let  $\phi \in \mathcal{G}_p$ . Assume (1.4). Then  $I_\rho$  is bounded from  $L_{p,\phi}(\mathbb{R}^d)$  to  $L_{q,\phi^{p/q},\text{weak}}(\mathbb{R}^d)$  if and only if (1.5) holds. 6

THEOREM 1.10. Let  $\phi, \psi \in \mathcal{G}_1$ . Assume that (1.6). Then  $I_\rho$  is bounded from  $L_{1,\phi}(\mathbb{R}^d)$  to  $L_{1,\psi,\text{weak}}(\mathbb{R}^d)$  if and only if (1.7) holds. 78

Here we will recall some works related to generalized fractional integral operators and generalized Morrey spaces.

In [17, 25, 31, 35], the authors generalized Morrey spaces to various directions. In [25], generalized Morrey spaces with variable growth condition are defined. The work [35] is a passage from [25] to the metric measure space whose underlying measure does not satisfy the doubling condition. The paper [34] is a counterpart to the weak type spaces. A further generalization is done in [17] and some related examples can be found in [21, 36]. As another generalization, in [31], Morrey spaces are generalized to martingale Morrey spaces. In [23], the authors applied generalized Morrey spaces to grasp the limiting case. In [33] Peetre proved there the boundedness of singular integral operator on generalized Morrey-Campanato spaces. The weak variants of the boundedness of the maximal operators, fractional integral operators and singular integral operators in generalized Morrey spaces were also investigated in the papers [11, 12].

In [2, 10, 14, 15, 24, 37, 42, 45] generalized fractional integral operators are investigated. The works [14, 15] are oriented to the boundedness of  $I_\rho$  and Guliyev and

Mustafayev used (1.2), (1.5) and (1.6) as a sufficient condition. In [42], the author placed themselves in the setting of non-doubling measure spaces and in [42, Theorem 2.1] the condition (1.6) showed up. In [24], the author extended the Gagliardo-Nirenberg inequality for  $I_\alpha$ , the case when  $\rho(t) = t^\alpha$ , to the one for  $I_\rho$ . An intersection of two classical Morrey spaces with the same parameter  $p$  can be regarded as a generalized Morrey space. Sugano investigated the boundedness of fractional integral operators on intersections of Morrey spaces [45], where she postulated conditions stronger than (1.2) and (1.5).

Other operators such as commutators are taken up in [12, 37]. In [12, (5.1)] Guliyev, Karaman, Seymur and Shukurov essentially considered (1.5) in order to show the boundedness of the fractional maximal operator of order  $\alpha$ .

In order to show the boundedness of commutators generated by BMO functions and  $I_\alpha$ , in [37], the authors postulated  $\phi$  on (1.2) and (1.5).

Hereafter, the letter  $C$  denotes a positive constant whose value may differ from line to line, which may depend on  $d, \rho, p$  and  $q$ , but not on the functions  $f$  and the variables  $x$ .

This paper is organized as follows: In Section 2, we shall give the norm estimates of the characteristic functions of balls and the function  $\phi(|\cdot|)$ , and calculate the image by  $I_\rho$  of these functions. Based upon these preliminary results, we shall prove Theorems 1.1–1.3 and 1.8–1.10, the main results in Section 3. Some examples are presented in Section 4 and they will show that Theorems 1.1–1.3 and 1.8–1.10 are independent.

### 2. Some norm and integral estimates

Let us first consider the characteristic functions of balls. For every  $R > 0$ , let  $B_R := B(0, R)$  be the ball centered at 0 with radius  $R$ , and  $\chi_{B_R}$  be the characteristic function of  $B_R$ . Recall  $\tilde{\rho}(r) = \int_0^r \frac{\rho(s)}{s} ds$ . Also we write  $B(x, R) = \{y \in \mathbb{R}^d : |x - y| < R\}$ .

The following lemmas will be used several times in this paper.

**LEMMA 2.1.** *There exists a constant  $C > 0$  such that the inequality  $\tilde{\rho}(R/2) \leq CI_\rho \chi_{B_R}(x)$  holds whenever  $x \in B_{R/2}$  and  $R > 0$ .*

*Proof.* Take  $x \in B_{R/2}$ . We write the integral in full:

$$I_\rho \chi_{B_R}(x) = \int_{\mathbb{R}^d} \frac{\rho(|x - y|)}{|x - y|^d} \chi_{B_R}(y) dy = \int_{B_R} \frac{\rho(|x - y|)}{|x - y|^d} dy.$$

A geometric observation shows that  $B(x, R/2) \subseteq B(0, R)$ . Hence, we have

$$I_\rho \chi_{B_R}(x) \geq \int_{B(x, R/2)} \frac{\rho(|x - y|)}{|x - y|^d} dy = C \int_0^{R/2} \frac{\rho(s)}{s} ds.$$

Note that we only use the spherical coordinates to obtain the last integral.  $\square$

LEMMA 2.2. For every  $R > 0$  and a measurable function  $\phi : (0, \infty) \rightarrow (0, \infty)$  satisfying the doubling condition

$$\frac{1}{C} \leq \frac{\phi(s)}{\phi(r)} \leq C \quad (0 < r/2 \leq s \leq 2r), \tag{2.1}$$

the inequality

$$C^{-1} \int_{2R}^{\infty} \frac{\phi(t)\rho(t)}{t} dt \leq I_{\rho}g_R(x) \leq C \int_{2R/3}^{\infty} \frac{\phi(t)\rho(t)}{t} dt$$

holds whenever  $x \in B_{R/3}$ , where  $g_R(x) := \phi(|x|)\chi_{B_R^c}(x)$ .

*Proof.* We prove the right-hand inequality, the left-hand inequality being similar. A geometric observation shows that  $|x - y| \sim |y|$  for all  $x \in B_{R/3}$  and  $y \in \mathbb{R}^d \setminus B_{2R/3}$ . Since  $\phi$  satisfies (2.1), then

$$\begin{aligned} I_{\rho}g_R(x) &= \int_{\mathbb{R}^d \setminus B_R} \frac{\phi(|y|)\rho(|x-y|)}{|x-y|^d} dy \\ &\leq \int_{\mathbb{R}^d \setminus B(x, 2R/3)} \frac{\phi(|y|)\rho(|x-y|)}{|x-y|^d} dy \\ &= \int_{\mathbb{R}^d \setminus B_{2R/3}} \phi(|x-y|) \frac{\rho(|y|)}{|y|^d} dy \\ &\leq C \int_{2R/3}^{\infty} \frac{\phi(t)\rho(t)}{t} dt \text{ for } x \in B_{R/3}. \end{aligned}$$

It remains to write the most right-hand side in terms of the spherical coordinates.  $\square$

The lemma below gives an estimate for the norm of  $\chi_{B_R}$  in  $L_{p,\phi}(\mathbb{R}^d)$ .

LEMMA 2.3. Let  $1 \leq p < \infty$  and  $\phi \in \mathcal{G}_p$ . There exists a constant  $C > 0$  such that  $C^{-1}\phi(R)^{-1} \leq \|\chi_{B_R} : L_{p,\phi}\| \leq C\phi(R)^{-1}$  for all  $R > 0$ .

Lemma 2.3 is proven in [8, Proposition A] and [26, Lemma 3.3]. See [7] as well.

LEMMA 2.4. Let  $1 \leq p < \infty$  and  $\phi \in \mathcal{G}_p$ . Assume that there exists a constant  $C > 0$  such that

$$\int_0^R \phi(t)t^{d/p-1} dt \leq C\phi(R)R^{d/p} \quad (R > 0). \tag{2.2}$$

Then the function  $x \mapsto \phi(|x|)$  belongs to  $L_{p,\phi}(\mathbb{R}^d)$ .

*Proof.* First note that (2.2) is equivalent to

$$\frac{1}{r^d} \int_0^r \phi(t)t^{d-1} dt \leq C\phi(r)^p \quad (r > 0) \tag{2.3}$$



in view of [30, Lemma 7.1]. Let  $\phi_1(r) = \inf_{t \leq r} \phi(t)$ . Then,  $\phi_1(r)$  is a non-increasing function such that  $L_{p,\phi}(\mathbb{R}^d)$  and  $L_{p,\phi_1}(\mathbb{R}^d)$  are isomorphic [26, p. 446]. Hence, we can assume that  $\phi$  itself is decreasing. In this case  $x \mapsto \phi(|x|)$  is radial decreasing, so that

$$\left[ \frac{1}{|B(a,r)|} \int_{B(a,r)} \phi(|x|)^p dx \right]^{1/p} \leq \left[ \frac{1}{|B_r|} \int_{B_r} \phi(|x|)^p dx \right]^{1/p} \quad (a \in \mathbb{R}^d). \tag{2.4}$$

Combining (2.3) and (2.4) and using the spherical coordinate, we obtain the desired result.  $\square$

### 3. Proofs of main results

In this section we prove our main results, six theorems in Section 1. We prove their sufficiency in Subsection 3.1 and then prove their necessity in Subsection 3.2.

#### 3.1. Proof of sufficiency

We remark that (1.5) includes (1.3). We prove the estimate (3.1). Once we prove (3.1), the estimate (3.1) gives us the boundedness of  $I_\rho$  from  $L_{p,\phi}(\mathbb{R}^d)$  to  $L_{q,\phi^{p/q}}(\mathbb{R}^d)$  as we shall see below. Here we use the fact that the Hardy-Littlewood maximal operator  $M$  is bounded on  $L_{p,\phi}(\mathbb{R}^d)$ , if  $p > 1$  and  $\phi$  is almost decreasing [29, 35]. See [25, 43, 44] for more information about  $M$  on the space  $L_{p,\phi}(\mathbb{R}^d)$ .

LEMMA 3.1. Let  $1 \leq p < q \leq \infty$  and let  $\phi \in \mathcal{G}_p$ . We assume (1.2) and (1.3), or we assume (1.5). If we normalize the norm of  $f$  by  $\|f\|_{L_{p,\phi}} = 1$ , then we have

$$|I_\rho f(x)| \leq C \left( [Mf(x)]^{p/q} + \inf_{r>0} \phi(r)^{p/q} \right), \quad x \in \mathbb{R}^d. \tag{3.1}$$

*Proof.* First we may assume that  $\phi$  is continuous and strictly decreasing (see [29, Proposition 3.4]). Recall that  $k_1$  and  $k_2$  appeared in the condition (1.1) of  $\rho$ . Let  $\rho^*(r) = \int_{k_1 r}^{k_2 r} \frac{\rho(s)}{s} ds$ . We have

$$|I_\rho f(x)| \leq C \left[ \sum_{j=-\infty}^{-1} + \sum_{j=0}^{\infty} \frac{\rho^*(2^j r)}{(2^j r)^d} \int_{|x-y| < 2^j r} |f(y)| dy \right]$$

for given  $x \in \mathbb{R}^d$  and  $r > 0$ . Let  $\Sigma_I$  and  $\Sigma_{II}$  be the first and second summations above. Now we invoke the overlapping property in [40]:

$$\begin{aligned} \sum_{j=-\infty}^{-1} \rho^*(2^j r) &\leq \sum_{j=-\infty}^{-1} \int_{2^j k_1 r}^{2^j k_2 r} \frac{\rho(s)}{s} ds \\ &= \int_0^{k_2 r} \left( \sum_{j=-\infty}^{-1} \chi_{[2^j k_1 r, 2^j k_2 r]}(s) \right) \frac{\rho(s)}{s} ds \\ &\leq C \int_0^{k_2 r} \frac{\rho(s)}{s} ds \\ &\leq C \tilde{\rho}(k_2 r) \end{aligned}$$

and

$$\begin{aligned} \sum_{j=0}^{\infty} \rho^*(2^j r) \phi(2^j r) &= \int_{k_1 r}^{\infty} \left( \sum_{j=0}^{\infty} \chi_{[2^j k_1 r, 2^{j+1} k_1 r]}(s) \frac{\rho(s)}{s} \phi(2^j r) \right) ds \\ &\leq C \int_{k_1 r}^{\infty} \left( \sum_{j=0}^{\infty} \chi_{[2^j k_1 r, 2^{j+1} k_1 r]}(s) \right) \frac{\rho(s)}{s} \phi(s) ds \\ &\leq C \int_{k_1 r}^{\infty} \frac{\rho(s)}{s} \phi(s) ds. \end{aligned}$$

Then, we have

$$\begin{aligned} \Sigma_I &\leq C \sum_{j=-\infty}^{-1} \rho^*(2^j r) Mf(x) \leq C \tilde{\rho}(k_2 r) Mf(x) \leq C \phi(r)^{p/q-1} Mf(x), \\ \Sigma_{II} &\leq C \sum_{j=0}^{\infty} \rho^*(2^j r) \phi(2^j r) \|f : L_{p,\phi}\| \leq C \int_{k_1 r}^{\infty} \frac{\rho(s) \phi(s)}{s} ds. \end{aligned}$$

We use (1.2) or (1.5) now. By the doubling property of  $\phi$ , we obtain  $\Sigma_{II} \leq C \phi(r)^{p/q}$ . Hence,

$$|I_{\rho} f(x)| \leq C \phi(r)^{p/q-1} [Mf(x) + \phi(r)] \quad (\text{for all } r > 0). \tag{3.2}$$

First assume  $Mf(x) \leq \inf_{r>0} \phi(r)$ . Then, (3.1) is immediate from (3.2).

Next, we assume  $Mf(x) > \inf_{r>0} \phi(r)$ . Since  $\|f : L_{p,\phi}\| = 1$ , we have

$$1 \geq \frac{1}{\phi(r)} \left( \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)|^p dy \right)^{1/p} \geq \frac{1}{\phi(r)} \cdot \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy.$$

Hence

$$\frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy \leq \phi(r)$$

for all  $r > 0$ . This implies

$$\frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy \leq \sup_{R>0} \phi(R)$$

for all  $r > 0$ . Since  $r > 0$  and  $x \in \mathbb{R}^d$  are arbitrary, it follows that  $Mf(x) \leq \sup_{r>0} \phi(r)$ . We can thus find  $R > 0$  such that  $Mf(x) \sim 2\phi(R)$  and, with this  $R$ , we can obtain (3.1).  $\square$

*Proof of Theorems 1.1 and 1.2 (Sufficiency).* Let  $\|f : L_{p,\phi}\| = 1$ . Let  $B = B(z,s)$  be an arbitrary ball. If we integrate (3.1), then we have

$$\frac{1}{|B|} \int_B |I_{\rho} f(x)|^q dx \leq C \left( \frac{1}{|B|} \int_B [Mf(x)]^p dx + \inf_{r>0} \phi(r)^p \right), \quad x \in \mathbb{R}^d.$$

If we divide both sides by  $\phi(s)^p$ , then we have

$$\frac{1}{\phi(s)^p |B|} \int_B |I_{\rho} f(x)|^q dx \leq C \left( \frac{1}{\phi(s)^p |B|} \int_B [Mf(x)]^p dx + 1 \right) \leq C$$

by virtue of the boundedness of the maximal operator  $M$  on  $L_{p,\phi}(\mathbb{R}^n)$ . The ball  $B$  being arbitrary, we obtain the desired result.  $\square$

*Proof of Theorem 1.3 (Sufficiency).* Let  $\|f : L_{p,\phi}\| = 1$ . For a ball  $B(z, r)$ , let  $f_1 = f\chi_{B(z, 2r)}$  and  $f_2 = f - f_1$ . Then a geometric observation shows  $B(z, r) \subset B(y, 3r)$  for all  $y \in B(z, 2r)$ . Hence by the Fubini theorem and the normalization,

$$\begin{aligned} \int_{B(z,r)} |I_\rho f_1(x)| dx &\leq \int_{B(z,r)} \left( \int_{B(z,2r)} |f(y)| \frac{\rho(|x-y|)}{|x-y|^d} dy \right) dx \\ &\leq \int_{B(z,2r)} \left( \int_{B(y,3r)} |f(y)| \frac{\rho(|x-y|)}{|x-y|^d} dx \right) dy \\ &\leq C\bar{\rho}(3r)\phi(3r)r^d \\ &\leq C\psi(r)r^d. \end{aligned}$$

Here for the last inequality we used (1.7) and the doubling condition of  $\psi$ . Thus, the estimate for  $f_1$  is valid. As for  $f_2$ , we let  $x \in B(z, r)$ . Then we have

$$|I_\rho f_2(x)| \leq \int_{B(z,2r)^c} |f(y)| \frac{\rho(|x-y|)}{|x-y|^d} dy \leq \int_{B(x,r)^c} |f(y)| \frac{\rho(|x-y|)}{|x-y|^d} dy$$

and decomposing the right-hand side dyadically as we did in the proof of Theorem 1.1 for  $\Sigma_{II}$ , we obtain

$$|I_\rho f_2(x)| \leq \sum_{j=1}^{\infty} \int_{B(x,2^j r) \setminus B(x,2^{j-1} r)} |f(y)| \frac{\rho(|x-y|)}{|x-y|^d} dy \leq C \int_{2k_1 r}^{\infty} \frac{\phi(t)\rho(t)}{t} dt.$$

If we use (1.7) and the doubling condition of  $\psi$ , then we obtain  $|I_\rho f_2(x)| \leq C\psi(r)$ . Thus, the estimate for  $f_2$  is valid as well.  $\square$

*Proof of Theorems 1.8 and 1.9 (Sufficiency).* We normalize  $f$  so that we have  $\|f : L_{p,\phi}\| = 1$ . By virtue of (3.1), we have

$$\begin{aligned} &\frac{1}{\phi(r)^{p/q}} \left( \frac{t^q m(B(a, r), |I_\rho f|, t)}{|B(a, r)|} \right)^{1/q} \\ &\leq C \frac{1}{\phi(r)^{p/q}} \left( \frac{t^q m(B(a, r), [Mf]^{p/q}, t/2)}{|B(a, r)|} \right)^{1/q} \\ &\quad + C \frac{1}{\phi(r)^{p/q}} \left( \frac{t^q m(B(a, r), \inf_{r' > 0} \phi(r')^{p/q}, t/2)}{|B(a, r)|} \right)^{1/q} \\ &\leq C \frac{1}{\phi(r)^{p/q}} \left( \frac{t^q m(B(a, r), [Mf]^{p/q}, t/2)}{|B(a, r)|} \right)^{1/q} + C \frac{1}{\phi(r)^{p/q}} \inf_{r' > 0} \phi(r')^{p/q} \\ &\leq C (\|Mf\|_{p,\phi, \text{weak}})^{p/q} + C \\ &\leq C. \end{aligned}$$

Here for the second to the last inequality, we have used [29, Theorem 6.1]. Theorem 1.9 can be proved in the same way.  $\square$

*Proof of Theorem 1.10 (Sufficiency).* Under the condition (1.7), in the proof of the sufficiency of Theorem 1.3 we already established that  $I_\rho$  is bounded from  $L_{1,\phi}(\mathbb{R}^d)$  to  $L_{1,\psi}(\mathbb{R}^d)$ , which is stronger than the boundedness from  $L_{1,\phi}(\mathbb{R}^d)$  to  $L_{1,\psi,\text{weak}}(\mathbb{R}^d)$ .  $\square$

REMARK 3.2. The proof of the sufficient part is similar to, but not the same as, that in [18, 28]. In this paper, we do not assume that  $\rho$  satisfies the doubling condition nor that  $\phi$  is surjective, as we did in [18].

### 3.2. Proof of necessity

*Proof of Theorem 1.1 (Necessity).* Note that

$$\tilde{\rho}(R/2) \leq C \left[ \frac{1}{|B_{R/2}|} \int_{B_{R/2}} |I_\rho \chi_{B_R}(x)|^q dx \right]^{1/q} \tag{3.3}$$

by virtue of a pointwise estimate in Lemma 2.1. Notice also that

$$\|I_\rho \chi_{B_R} : L_{q,\phi^{p/q}}\| \leq C \|\chi_{B_R} : L_{p,\phi}\| \tag{3.4}$$

since  $I_\rho$  is assumed bounded. If we combine (3.3), (3.4), Lemma 2.3, and the doubling property of  $\phi$ , we have

$$\begin{aligned} \tilde{\rho}(R/2) &\leq C \phi(R/2)^{p/q} \phi(R/2)^{-p/q} \left[ \frac{1}{|B_{R/2}|} \int_{B_{R/2}} |I_\rho \chi_{B_R}(x)|^q dx \right]^{1/q} \\ &\leq C \phi(R/2)^{p/q} \|I_\rho \chi_{B_R} : L_{q,\phi^{p/q}}\| \\ &\leq C \phi(R/2)^{p/q} \phi(R)^{-1} \\ &\leq C \phi(R/2)^{p/q-1}, \end{aligned}$$

for all  $R > 0$ . This completes the proof.  $\square$

*Proof of Theorem 1.2 (Necessity).* The exactly same argument as we did for Theorem 1.1 works and we conclude

$$\int_0^r \frac{\rho(t)}{t} dt \leq C \phi(r)^{p/q-1}$$

for  $r > 0$ .

Meanwhile, by virtue of Lemma 2.2, we obtain

$$\int_{2R}^\infty \frac{\phi(t)\rho(t)}{t} dt \sim \left( \frac{1}{R^d} \int_{B_{R/3}} I_\rho g(x)^q dx \right)^{1/q} \leq C \phi(R)^{p/q} \|I_\rho g_R : L_{q,\phi^{p/q}}\|.$$

Since  $I_\rho$  is bounded, we obtain

$$\int_{2R}^\infty \frac{\phi(t)\rho(t)}{t} dt \leq C\phi(R)^{p/q} \|g_R : L_{p,\phi}\|.$$

Now we invoke Lemma 2.4 to conclude

$$\int_{2R}^\infty \frac{\phi(t)\rho(t)}{t} dt \leq C\phi(R)^{p/q} \leq C\phi(2R)^{p/q}.$$

Thus, Theorem 1.2 is proven.  $\square$

*Proof of Theorem 1.3 (Necessity).* Assume that  $I_\rho$  is bounded from  $L_{1,\phi}(\mathbb{R}^d)$  to  $L_{1,\psi}(\mathbb{R}^d)$ . By Lemma 2.1, we obtain

$$\tilde{\rho}(r) \sim r^{-d} \int_{B_{r/2}} I_\rho \chi_{B_r}(x) dx \leq \psi\left(\frac{r}{2}\right) \|I_\rho \chi_{B_r} : L_{1,\psi}\|.$$

Since  $\psi \in \mathcal{G}_1$  and  $I_\rho$  is assumed bounded from  $L_{1,\phi}(\mathbb{R}^d)$  to  $L_{1,\psi}(\mathbb{R}^d)$ , it follows that  $\tilde{\rho}(r) \leq C\psi(r) \|\chi_{B_r} : L_{1,\phi}\|$ . Since  $\|\chi_{B_r} : L_{1,\phi}\| \sim \phi(r)^{-1}$ , we conclude  $\tilde{\rho}(r) \leq C\frac{\psi(r)}{\phi(r)}$ . Meanwhile, by Lemma 2.2, we have

$$\int_r^\infty \frac{\rho(t)\phi(t)}{t} dt \leq C\psi\left(\frac{r}{6}\right) \|I_\rho g_r : L_{1,\psi}\| \leq C\psi(r) \|g_r : L_{1,\phi}\| \leq C\psi(r).$$

Thus, Theorem 1.3 is proved.  $\square$

*Proof of Theorem 1.8 (Necessity).* Choose  $C > 0$  from Lemma 2.1 and write it for  $C_0$ . By Lemma 2.1 we have

$$\begin{aligned} \sup_{t>0} t^q m(B_{R/2}, I_\rho \chi_{B_R}(x), t) &\geq \left(\frac{\tilde{\rho}(R/2)}{C_0}\right)^q m\left(B_{R/2}, I_\rho \chi_{B_R}, \frac{\tilde{\rho}(R/2)}{C_0}\right) \\ &\geq \left(\frac{\tilde{\rho}(R/2)}{C_0}\right)^q |B_{R/2}|. \end{aligned}$$

If we arrange this inequality, then we have

$$\tilde{\rho}(R/2) \leq C \left[ \frac{\sup_{t>0} t^q m(B_{R/2}, I_\rho \chi_{B_R}, t)}{|B_{R/2}|} \right]^{1/q}. \tag{3.5}$$

Therefore

$$\begin{aligned} \tilde{\rho}(R/2) &\leq C \phi(R/2)^{p/q} \phi(R/2)^{-p/q} \left[ \frac{\sup_{t>0} t^q m(B_{R/2}, I_\rho \chi_{B_R}, t)}{|B_{R/2}|} \right]^{1/q} \\ &\leq C \phi(R/2)^{p/q} \|I_\rho \chi_{B_R} : L_{q,\phi^{p/q},\text{weak}}\| \\ &\leq C \phi(R/2)^{p/q} \phi(R)^{-1} \\ &\leq C \phi(R/2)^{p/q-1}, \end{aligned}$$

for all  $R > 0$ . This completes the proof.  $\square$

*Proof of Theorem 1.9 (Necessity).* The exactly same argument as we did for Theorem 1.8 works and we conclude

$$\int_0^r \frac{\rho(t)}{t} dt \leq C\phi(r)^{p/q-1}$$

for  $r > 0$ . Meanwhile, by virtue of Lemma 2.2, we obtain

$$\begin{aligned} & \sup_{t>0} t^q m(B_{R/3}, I_\rho g_{B_R}, t) \\ & \geq \left( \frac{1}{C} \int_{2R}^\infty \frac{\phi(t)\rho(t)}{t} dt \right)^q m\left(B_{R/3}, I_\rho g_{B_R}, \frac{1}{C} \int_{2R}^\infty \frac{\phi(t)\rho(t)}{t} dt\right) \\ & \geq \left( \frac{1}{C} \int_{2R}^\infty \frac{\phi(t)\rho(t)}{t} dt \right)^q |B_{R/3}|. \end{aligned}$$

Hence it follows that

$$\begin{aligned} \int_{2R}^\infty \frac{\phi(t)\rho(t)}{t} dt & \leq C \left[ \frac{\sup_{t>0} t^q m(B_{R/3}, I_\rho g_{B_R}, t)}{|B_{R/3}|} \right]^{1/q} \\ & \leq C\phi(R/3)^{p/q} \|I_\rho g_R : L_{q,\phi^{p/q},\text{weak}}\| \\ & \leq C\phi(R/3)^{p/q} \|g_R : L_{p,\phi}\| \leq C\phi(R/3)^{p/q} \leq C\phi(2R)^{p/q}. \end{aligned}$$

Thus, Theorem 1.9 is proven.  $\square$

*Proof of Theorem 1.10 (Necessity).* Assume that  $I_\rho$  is bounded from  $L_{1,\phi}(\mathbb{R}^d)$  to  $L_{1,\psi,\text{weak}}(\mathbb{R}^d)$ . By Lemma 2.1, we obtain

$$\tilde{\rho}(R/2) \leq C \frac{\sup_{t>0} t m(B_{R/2}, I_\rho \chi_{B_R}(x), t)}{|B_{R/2}|} \leq \psi\left(\frac{R}{2}\right) \|I_\rho \chi_{B_R} : L_{1,\psi,\text{weak}}\|.$$

Since  $\psi \in \mathcal{G}_1$  and  $I_\rho$  is assumed bounded from  $L_{1,\phi}(\mathbb{R}^d)$  to  $L_{1,\psi,\text{weak}}(\mathbb{R}^d)$ , it follows that

$$\tilde{\rho}(r) \leq C\psi(r) \|\chi_{B_{2r}} : L_{1,\phi}\| \sim \frac{\psi(r)}{\phi(r)}.$$

Meanwhile, by Lemma 2.2, we have

$$\begin{aligned} \int_r^\infty \frac{\rho(t)\phi(t)}{t} dt & \leq C\psi\left(\frac{r}{6}\right) \|I_\rho g_r : L_{1,\psi,\text{weak}}\| \\ & \leq C\psi(r) \|g_r : L_{1,\phi}\| \\ & \leq C\psi(r). \end{aligned}$$

Thus, Theorem 1.10 is proved.  $\square$

**4. Examples showing Theorems 1.1–1.3 are independent**

In this section, we show by examples that Theorems 1.1–1.3 are of independent interest. Here and below we write

$$\ell_{\beta_1, \beta_2}(r) := \begin{cases} (\log \frac{1}{r})^{-\beta_1} & (0 < r < e^{-1}), \\ 1 & (e^{-1} \leq r \leq e), \\ (\log r)^{\beta_2} & (e < r). \end{cases}$$

This function is used to describe the “log”-growth and “log”-decay properties. Also, we fix  $p$  and  $q$  so that  $1 < p < q < \infty$ .

**EXAMPLE 1.** Let  $\mu_1, \mu_2$  satisfy  $\mu_1, \mu_2 \geq 0$ . Set  $\alpha := \frac{d}{p} - \frac{d}{q}$  and  $\beta_i := (\frac{p}{q} - 1)\mu_i$  for  $i = 1, 2$ . Define  $\rho(r) := r^\alpha \ell_{\beta_1, \beta_2}(r)$ ,  $\phi(r) := r^{-\frac{d}{p}} \ell_{\mu_1, \mu_2}(r)$  for  $r > 0$ . Then this pair  $(\rho, \phi)$  fulfills the assumptions (1.2) and (1.3) in Theorem 1.1 but fails (1.4) in Theorem 1.2. More precisely, since  $\alpha > 0$ , we have  $\tilde{\rho}(r) \sim \rho(r)$  and  $\int_r^\infty \frac{\phi(t)\rho(t)}{t} dt \sim \phi(r)\rho(r)$ .

Example 1 is an endpoint case of the next example.

**EXAMPLE 2.** Let  $\lambda$  satisfy  $0 < (\frac{p}{q} - 1)\lambda < d$  and  $-\frac{d}{p} < \lambda < 0$ . Take  $\mu_1, \mu_2$  arbitrarily. Set  $\alpha := (\frac{p}{q} - 1)\lambda$  and  $\beta_i := (\frac{p}{q} - 1)\mu_i$  for  $i = 1, 2$ . Define  $\rho(r) := r^\alpha \ell_{\beta_1, \beta_2}(r)$  and  $\phi(r) := r^\lambda \ell_{\mu_1, \mu_2}(r)$  for  $r > 0$ . Then this pair  $(\rho, \phi)$  fulfills the assumptions (1.2)–(1.4) in Theorems 1.1 and 1.2. Indeed,  $\tilde{\rho}(r) \sim \rho(r)$ ,  $\int_r^\infty \frac{\phi(t)\rho(t)}{t} dt \sim \phi(r)\rho(r)$ .

The next example concerns the case when the spaces are close to  $L^\infty(\mathbb{R}^d)$  and the smoothing order of  $I_\rho$  is “almost 0”.

**EXAMPLE 3.** Let  $\mu_1, \mu_2 < 0$ . Set  $\beta_1 := (\frac{p}{q} - 1)\mu_1 + 1 \in (1, \infty)$  and  $\beta_2 := (\frac{p}{q} - 1)\mu_2 - 1 \in (-1, \infty)$ . Define  $\rho(r) := \ell_{\beta_1, \beta_2}(r)$  and  $\phi(r) := \ell_{\mu_1, \mu_2}(r)$  for  $r > 0$ . Then this pair  $(\rho, \phi)$  fulfills the assumptions (1.4) and (1.5) in Theorem 1.2 but fails (1.2) in Theorem 1.1. More precisely, for all  $r > 0$ , we have  $\tilde{\rho}(r) \sim \ell_{\beta_1-1, \beta_2+1}(r)$ , and  $\int_r^\infty \frac{\phi(t)\rho(t)}{t} dt \sim \ell_{\mu_1+\beta_1-1, \mu_2+\beta_2+1}(r)$ .

We consider a case when the target space is close to  $L^\infty(\mathbb{R}^d)$ .

**EXAMPLE 4.** Let  $\alpha, \beta_1, \mu_1, \mu_2$  satisfy  $0 < \alpha < \frac{d}{p}$ ,  $\mu_1 + \beta_1 < 1$ ,  $\mu_2 < 0$ . Set  $\beta_2 := (\frac{p}{q} - 1)\mu_2 - 1 \in (-1, \infty)$ . Define  $\rho(r) := \min(1, r^\alpha)\ell_{\beta_1, \beta_2}(r)$  and  $\phi(r) := \max(1, r^{-\alpha})\ell_{\mu_1, \mu_2}(r)$  for  $r > 0$ . Then this pair  $(\rho, \phi)$  fulfills the assumptions (1.4) and (1.5) in Theorem 1.2 but fails (1.2) in Theorem 1.1. More precisely,  $\phi(r)\tilde{\rho}(r) \sim \ell_{\mu_1+\beta_1, \mu_2+\beta_2+1}(r)$  and  $\int_r^\infty \frac{\phi(t)\rho(t)}{t} dt \sim \ell_{\mu_1+\beta_1-1, \mu_2+\beta_2+1}(r)$  for  $r > 0$ .

We conclude this paper by proving Proposition 1.7.

EXAMPLE 5. Let  $0 < s < d$ . Define  $\phi(r) := r^{-s}$  and  $\psi(r) := (1+r)^{-s} \ell_{-1,0}(r)$  for  $r > 0$ . Let  $\rho(r) := r^d G_s(r)$ , where  $G_s$  denotes the Bessel kernel, the kernel of  $(1 - \Delta)^{s/2}$ . Observe that  $\tilde{\rho}(r) \sim \min(r^s, 1)$  and hence  $\phi(r)\tilde{\rho}(r) \sim \min(1, r^{-s})$ . Note also that  $\int_r^\infty \frac{\phi(t)\rho(t)}{t} dt \sim \begin{cases} \log(e/r) & (r < 1), \\ r^{d-s} G_s(r) & (r \geq 1). \end{cases}$  Then we have  $\phi(r)\tilde{\rho}(r) + \int_r^\infty \frac{\phi(t)\rho(t)}{t} dt \sim \psi(r)$ . Hence it follows from Theorem 1.3 that  $\|I_\rho f : L_{1,\psi}\| \leq C\|f : L_{1,\phi}\|$ , extending Proposition 1.7. This triple  $(\rho, \phi, \psi)$  fulfills the assumptions (1.7) and (1.6) in Theorem 1.3 but it fails (1.2) in Theorem 1.1 and (1.5) in Theorem 1.2.

*Acknowledgements.* The research was initiated when the first and the second authors visited Kyoto University under the GCOE 2011 Project. Eiichi Nakai is supported by Grant-in-Aid for Scientific Research (C), No. 24540159, Japan Society for the Promotion of Science. Yoshihiro Sawano is supported by Grant-in-Aid for Young Scientists (B) No. 24740085 Japan Society for the Promotion of Science.

The authors are grateful to Professor Vagif Guliyev for his generosity of sending us the works [14, 15]. The authors are grateful to the anonymous referee for his/her pointing out the paper [2].

#### REFERENCES

- [1] D. R. ADAMS, *A note on Riesz potentials*, Duke Math. J. **42** (1975), 765–778.
- [2] W. S. BUDHI AND J. LINDIARTI, *Boundedness of multilinear generalized fractional integral operators in generalized Morrey space*, Far East J. Math. Sci. **57**, 1 (2011), 91–104.
- [3] V. BURENKOV AND V. GULIYEV, *Necessary and sufficient conditions for the boundedness of the Riesz potential in local Morrey-type spaces*, Potential Anal. **30**, 3 (2009), 211–249.
- [4] V. I. BURENKOV, A. GOGATISHVILI, V. S. GULIYEV AND R. CH. MUSTAFAYEV, *Boundedness of the Riesz potential in local Morrey-type spaces*, Potential Analysis **35**, 1 (2011), 67–87.
- [5] F. CHIARENZA AND M. FRASCA, *Morrey spaces and Hardy-Littlewood maximal function*, Rend. Mat. **7** (1987), 273–279.
- [6] ERIDANI, H. GUNAWAN, AND E. NAKAI, *On generalized fractional integral operators*, Sci. Math. Jpn. **60** (2004), 539–550.
- [7] ERIDANI, M. I. UTOYO, AND H. GUNAWAN, *A characterization for fractional integral operators on generalized Morrey spaces*, Anal. Theory Appl. **28**, 3 (2012), 263–267.
- [8] ERIDANI AND Y. SAWANO, *Fractional Integral Operators in Generalized Morrey Spaces Defined on Metric Measure Spaces*, Proc. A. Razmadze Math. Inst. **158** (2012), 13–24.
- [9] ERIDANI, AND M. I. UTOYO, *A note on Nakai's results*, Uzbek. Mat. Zh. **4** (2009), 172–179.
- [10] A. D. GADJIEV, *On generalized potential-type integral operators, Dedicated to Roman Taberski on the occasion of his 70th birthday*, Funct. Approx. Comment. Math. **25** (1997), 37–44.
- [11] V. S. GULIYEV, *Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces*, Journal of inequalities and applications, Volume **2009**, Article ID 503948, 20 pages.
- [12] V. S. GULIYEV, S. S. ALIYEV, T. KARAMAN, AND P. S. SHUKUROV, *Boundedness of sublinear operators and commutators on generalized Morrey spaces*, Integral Equations and Operator Theory **71**, Issue 3 (2011), 327–355.
- [13] V. S. GULIYEV, J. HASANOV, AND Y. ZEREN, *Necessary and sufficient conditions for the boundedness of the Riesz potential in modified Morrey spaces*, J. Math. Inequal. **5**, 4 (2011), 491–506.
- [14] V. S. GULIYEV, S. S. ALIYEV, AND R. CH. MUSTAFAYEV, *On generalized fractional integrals*, Trans. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **21**, 4 (2001), Math. Mech., 63–71, 237.



- [15] V. S. GULIYEV, AND R. CH. MUSTAFAYEV, *On generalized fractional integrals on spaces of homogeneous type*, Khazar Journal of Mathematics **2**, 2 (2006), 49–60.
- [16] V. S. GULIYEV, AND R. MUSTAFAYEV, *Boundedness of the anisotropic maximal and anisotropic singular integral operators in generalized Morrey spaces*, Acta Math. Sin. (Engl. Ser.) **27**, 12 (2011), 2361–2370.
- [17] V. S. GULIYEV, AND Y. SAWANO, *Linear operators on Generalized Morrey spaces with non-doubling measures*, to appear in Publ. Math. Debrecen.
- [18] H. GUNAWAN, *A note on the generalized fractional integral operators*, J. Indones. Math. Soc. **9** (2003), 39–43.
- [19] H. GUNAWAN, AND ERIDANI, *Fractional integrals and generalized Olsen inequalities*, Kyungpook Math. J. **49** (2009), 31–39.
- [20] K. KURATA, AND S. SUGANO, *A remark on estimates for uniformly elliptic operators on weighted  $L^p$  spaces and Morrey spaces*, Math. Nachr. **209** (2000), 137–150.
- [21] L. LIGUANG, Y. SAWANO, AND D. YANG, *Morrey-type Spaces on Gauss Measure Spaces and Boundedness of Singular Integrals*, J. Geometrical Analysis, online.
- [22] Y. MIZUTA, E. NAKAI, T. OHNO, AND T. SHIMOMURA, *Boundedness of fractional integral operators on Morrey spaces and Sobolev embeddings for generalized Riesz potentials*, J. Math. Soc. Japan **62**, 3 (2010), 707–744.
- [23] Y. MIZUTA, E. NAKAI, T. OHNO, AND T. SHIMOMURA, *An elementary proof of Sobolev embeddings for Riesz potentials of functions in Morrey spaces  $L^{1,v,\beta}(G)$* , Hiroshima Math. J. **38**, 3 (2008), 425–436.
- [24] Y. MIZUTA, E. NAKAI, Y. SAWANO, AND T. SHIMOMURA, *Gagliardo-Nirenberg inequality for generalized Riesz potentials of functions in Musielak-Orlicz spaces*, Arch. Math. **98**, 3 (2012), 253–263.
- [25] E. NAKAI, *Hardy-Littlewood maximal operator, singular integral operators and the Riesz potentials on generalized Morrey spaces*, Math. Nachr. **166** (1994), 95–103.
- [26] E. NAKAI, *A characterization of pointwise multipliers on the Morrey spaces*, Sci. Math. **3** (2000), 445–454.
- [27] E. NAKAI, *On generalized fractional integrals*, Taiwanese J. Math. **5** (2001), 587–602.
- [28] E. NAKAI, *On generalized fractional integrals on the weak Orlicz spaces,  $BMO_\phi$ , the Morrey spaces and the Campanato spaces*, Function spaces, interpolation theory and related topics (Lund, 2000), 389–401, de Gruyter, Berlin, 2002.
- [29] E. NAKAI, *Orlicz-Morrey spaces and the Hardy-Littlewood maximal function*, Studia Math. **188**, 3 (2008), 193–221.
- [30] E. NAKAI, *A generalization of Hardy spaces  $H^p$  by using atoms*, Acta Math. Sin. (Engl. Ser.) **24**, 8 (2008), 1243–1268.
- [31] E. NAKAI AND G. SADASUE, *Martingale Morrey-Campanato spaces and fractional integrals*, J. Funct. Spaces Appl. **2012**, Art. ID 673929, 29 pp.
- [32] J. PEETRE, *On convolution operators leaving  $L^{p,\lambda}$  spaces invariant*, Ann. Mat. Pura Appl. **72** (1966), 295–304.
- [33] J. PEETRE, *On the theory of  $\mathcal{L}_{p,\lambda}$  spaces*, J. Funct. Anal. **4** (1969), 71–87.
- [34] Y. SAWANO,  *$l^q$ -valued extension of the fractional maximal operators for non-doubling measures via potential operators*, International Journal of Pure and Applied Mathematics **26**, 4 (2006), 505–523.
- [35] Y. SAWANO, *Generalized Morrey Spaces for non-doubling measures*, Nonlinear differ. equ. appl. **15** (2008), 413–425.
- [36] Y. SAWANO AND T. SHIMOMURA, *Sobolev embeddings for Riesz potentials of functions in non-doubling Morrey spaces of variable exponents*, Collectanea Mathematica, in press.
- [37] Y. SAWANO AND S. SHIRAI, *Compact commutators on Morrey spaces with non-doubling measures*, Georgian Math. J. **15**, 2 (2008), 353–376.
- [38] Y. SAWANO, S. SUGANO, AND H. TANAKA, *A note on generalized fractional integral operators on generalized Morrey spaces*, Bound. Value Probl. **2009**, Art. ID 835865, 18 pp.
- [39] Y. SAWANO, S. SUGANO, AND H. TANAKA, *Generalized fractional integral operators and fractional maximal operators in the framework of Morrey spaces*, Trans. Amer. Math. Soc. **363**, 12 (2011), 6481–6503.
- [40] Y. SAWANO, S. SUGANO AND H. TANAKA, *Orlicz-Morrey spaces and fractional operators*, Potential Anal. **36**, 4 (2012), 517–556.

- [41] Y. SAWANO AND H. WADADE, *On the Gagliardo-Nirenberg type inequality in the critical Sobolev-Morrey space*, J. Fourier Anal. Appl., online.
- [42] I. SIHWANINGRUM, S. MARYANI, AND H. GUNAWAN, *Weak type inequalities for fractional integral operators on generalized non-homogeneous Morrey spaces*, Anal. Theory and Appl. **28**, 1 (2012), 65–72.
- [43] E. M. STEIN, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, New Jersey, 1970.
- [44] E. M. STEIN, *Harmonic Analysis: Real Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, Princeton, New Jersey, 1993.
- [45] S. SUGANO, *Some inequalities for generalized fractional integral operators on generalized Morrey spaces*, Math. Inequal. Appl. **14**, 4 (2011), 849–865.
- [46] S. SUGANO AND H. TANAKA, *Boundedness of fractional integral operators on generalized Morrey spaces*, Sci. Math. Jpn. **58**, 3 (2003), 531–540 (Sci. Math. Jpn. Online **8** (2003), 233–242).

(Received October 6, 2012)

*Eridani*  
Department of Mathematics  
Airlangga University  
Surabaya 60115, Indonesia  
e-mail: eridani.dinadewi@gmail.com

*Hendra Gunawan*  
Department of Mathematics  
Bandung Institute of Technology  
Bandung 40132, Indonesia  
e-mail: hgunawan@math.itb.ac.id

*Eiichi Nakai*  
Department of Mathematics  
Ibaraki University  
Mito, Ibaraki 310-8512  
Japan  
e-mail: enakai@mx.ibaraki.ac.jp

*Yoshihiro Sawano*  
Department of Mathematics and Information Science  
Tokyo Metropolitan University  
1-1 Minami-Ohsawa, Hachioji, 192-0397  
Japan  
e-mail: yoshihiro-sawano@celery.ocn.ne.jp

# CHARACTERIZATIONS FOR THE GENERALIZED FRACTIONAL INTEGRAL OPERATORS ON MORREY SPACES

## ORIGINALITY REPORT

25%

SIMILARITY INDEX

14%

INTERNET SOURCES

23%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1

Yoshihiro Sawano. "Theory of Besov Spaces", Springer Nature America, Inc, 2018

Publication

1%

2

Tanaka, Hitoshi, and Hendra Gunawan. "The Local Trace Inequality for Potential Type Integral Operators", Potential Analysis, 2013.

Publication

1%

3

Nakai, E.. "Hardy spaces with variable exponents and generalized Campanato spaces", Journal of Functional Analysis, 20120501

Publication

1%

4

[isbfs.mns.kyutech.ac.jp](http://isbfs.mns.kyutech.ac.jp)

Internet Source

1%

5

[itu.diva-portal.org](http://itu.diva-portal.org)

Internet Source

1%

6

Seeger, A.. "Bounds for singular fractional integrals and related Fourier integral operators", Journal of Functional Analysis, 20030401

Publication

1%

7

[www.springerprofessional.de](http://www.springerprofessional.de)

Internet Source

<1%

8

Kokilashvili, Vakhtang, Mieczysław Mastyło, and Alexander Meskhi. "Fractional integral operators between Banach function lattices", Nonlinear Analysis Theory Methods & Applications, 2015.

Publication

<1%

9	LIN TANG. "ENDPOINT ESTIMATES FOR MULTILINEAR FRACTIONAL INTEGRALS", Journal of the Australian Mathematical Society, 06/2008 Publication	<1%
10	<a href="http://www.acadsci.fi">www.acadsci.fi</a> Internet Source	<1%
11	<a href="http://www.emis.de">www.emis.de</a> Internet Source	<1%
12	<a href="http://mathsoc.jp">mathsoc.jp</a> Internet Source	<1%
13	Operator Theory Advances and Applications, 2014. Publication	<1%
14	SHI-HUA CHEN. "THE EFFECTIVE MASS OF STRONG-COUPLING MAGNETOPOLARONS IN A PARABOLIC QUANTUM DOT", Modern Physics Letters B, 2011 Publication	<1%
15	You-qiang Wang. "A new class of deprivation-based generalized Gini indices", Economic Theory, 09/2000 Publication	<1%
16	Yasuo Komori. "Weighted Morrey spaces and a singular integral operator", Mathematische Nachrichten, 02/2009 Publication	<1%
17	<a href="http://dosyayukleme.ahievran.edu.tr">dosyayukleme.ahievran.edu.tr</a> Internet Source	<1%
18	<a href="http://hrcak.srce.hr">hrcak.srce.hr</a> Internet Source	<1%
19	<a href="http://www.heldermann-verlag.de">www.heldermann-verlag.de</a> Internet Source	<1%

- |    |  |     |
|----|--|-----|
| 20 | Sautbekova, Merey, and Winfried Sickel. "Strong summability of Fourier series and Morrey spaces", <i>Analysis Mathematica</i> , 2014.<br>Publication                           | <1% |
| 21 | Vagif S. Guliyev. "Boundedness of Sublinear Operators and Commutators on Generalized Morrey Spaces", <i>Integral Equations and Operator Theory</i> , 09/16/2011<br>Publication | <1% |
| 22 | Yasuo Komori. "Factorization of functions in $H^1(\mathbb{R}^n)$ and generalized Morrey spaces", <i>Mathematische Nachrichten</i> , 04/2006<br>Publication                     | <1% |
| 23 | <a href="http://boundaryvalueproblems.springeropen.com">boundaryvalueproblems.springeropen.com</a><br>Internet Source  | <1% |
| 24 | <a href="http://alpha.math.uga.edu">alpha.math.uga.edu</a><br>Internet Source  | <1% |
| 25 | Hua Wang. "Boundedness of intrinsic square functions on generalized Morrey spaces", <i>Georgian Mathematical Journal</i> , 2014<br>Publication                                 | <1% |
| 26 | <a href="http://www.mathsoc.jp">www.mathsoc.jp</a><br>Internet Source  | <1% |
| 27 | <a href="http://journalofinequalitiesandapplications.springeropen.com">journalofinequalitiesandapplications.springeropen.com</a><br>Internet Source                            | <1% |
| 28 | <a href="http://www.mia-journal.com">www.mia-journal.com</a><br>Internet Source  | <1% |
| 29 | preview-<br><a href="http://journalofinequalitiesandapplications.springeropen.com">journalofinequalitiesandapplications.springeropen.com</a><br>Internet Source                | <1% |
| 30 | "Concrete Operators, Spectral Theory, Operators in Harmonic Analysis and Approximation", Springer Nature America, Inc,   | <1% |

2014

Publication

---

31 Dachun Yang, Dongyong Yang, Guoen Hu. "The Hardy Space  $H_1$  with Non-doubling Measures and Their Applications", Springer Nature America, Inc, 2013

Publication

---

32 [nardus.mpn.gov.rs](http://nardus.mpn.gov.rs)

Internet Source

---

33 Nakai, Eiichi, and Gaku Sadasue. "Maximal function on generalized martingale Lebesgue spaces with variable exponent", Statistics & Probability Letters, 2013.

Publication

---

34 Evgeniya Burtseva, Natasha Samko. "On weighted generalized fractional and Hardy-type operators acting between Morrey-type spaces", Fractional Calculus and Applied Analysis, 2017

Publication

---

35 Brandon Carter. "Gradient Formula for Linearly Self-Interacting Branes", Communications in Mathematical Physics, 04/01/2003

Publication

---

36 [www.mat.ucm.es](http://www.mat.ucm.es)

Internet Source

---

37 [www.mis.hiroshima-u.ac.jp](http://www.mis.hiroshima-u.ac.jp)

Internet Source

---

38 [journals.cms.math.ca](http://journals.cms.math.ca)

Internet Source

---

39 Daniel Aalto. "The discrete maximal operator in metric spaces", Journal d Analyse Mathématique, 05/2010

Publication

---

[amarun.net](http://amarun.net)

40	Internet Source	<1%
41	"Function Spaces and Inequalities", Springer Nature, 2017 Publication	<1%
42	<a href="http://repository.dl.itc.u-tokyo.ac.jp">repository.dl.itc.u-tokyo.ac.jp</a> Internet Source	<1%
43	<a href="http://www.kkuniyuk.com">www.kkuniyuk.com</a> Internet Source	<1%
44	<a href="http://lnx.mathmods.eu">lnx.mathmods.eu</a> Internet Source	<1%
45	<a href="http://imagine.enpc.fr">imagine.enpc.fr</a> Internet Source	<1%
46	<a href="http://jeomj.rmi.acnet.ge">jeomj.rmi.acnet.ge</a> Internet Source	<1%
47	Eroglu, Ahmet M. "Boundedness of fractional oscillatory integral operators and their commutators on generalized Morrey spaces", Boundary Value Problems, 2013. Publication	<1%
48	<a href="http://www.jstage.jst.go.jp">www.jstage.jst.go.jp</a> Internet Source	<1%
49	Iida, Takeshi. "Weighted estimates of higher order commutators generated by BMO-functions and the fractional integral operator on Morrey spaces", Journal of Inequalities and Applications, 2016. Publication	<1%
50	<a href="http://repository.kulib.kyoto-u.ac.jp">repository.kulib.kyoto-u.ac.jp</a> Internet Source	<1%
51	<a href="http://lightning.itcs.uiuc.edu">lightning.itcs.uiuc.edu</a> Internet Source	<1%

- 52 Yoshihiro Mizuta. "Sobolev's inequality for Riesz potentials of functions in Morrey spaces of integral form", *Mathematische Nachrichten*, 06/29/2010  
Publication <1%
- 
- 53 [siba-ese.unile.it](http://siba-ese.unile.it)  
Internet Source <1%
- 
- 54 [matwbn.icm.edu.pl](http://matwbn.icm.edu.pl)  
Internet Source <1%
- 
- 55 [www.math.helsinki.fi](http://www.math.helsinki.fi)  
Internet Source <1%
- 
- 56 A. S. Balakishiyev, E. A. Gadjieva, F. Gürbüz, A. Serbetci. "Boundedness of some sublinear operators and their commutators on generalized local Morrey spaces", *Complex Variables and Elliptic Equations*, 2017  
Publication <1%
- 
- 57 [www.kurims.kyoto-u.ac.jp](http://www.kurims.kyoto-u.ac.jp)  
Internet Source <1%
- 
- 58 Mizuta, Yoshihiro, and Tetsu Shimomura. "Weighted Morrey spaces of variable exponent and Riesz potentials : Weighted Morrey spaces of variable exponent and Riesz potentials", *Mathematische Nachrichten*, 2014.  
Publication <1%
- 
- 59 [www.scribd.com](http://www.scribd.com)  
Internet Source <1%
- 
- 60 Yoshihiro Sawano, Hiroko Yoshida. "A predual of a predual of  $B\sigma$  and its applications to commutators", *Science China Mathematics*, 2018  
Publication <1%
- 
- 61 [link.springer.com](http://link.springer.com)  
Internet Source <1%



<1%

62

[actamath.savbb.sk](http://actamath.savbb.sk)

Internet Source

<1%

63

[www.stt.msu.edu](http://www.stt.msu.edu)

Internet Source

<1%

64

[ajmaa.org](http://ajmaa.org)

Internet Source

<1%

65

Eridani, A.. "Morrey spaces and fractional integral operators", *Expositiones Mathematicae*, 2009

Publication

<1%

66

[adatbank.transindex.ro](http://adatbank.transindex.ro)

Internet Source

<1%

67

[kenkyu-web.bur.osaka-kyoiku.ac.jp](http://kenkyu-web.bur.osaka-kyoiku.ac.jp)

Internet Source

<1%

68

[www.jams.or.jp](http://www.jams.or.jp)

Internet Source

<1%

69

Elman J. Ibrahimov, Ali Akbulut. "The Hardy–Littlewood–Sobolev theorem for Riesz potential generated by Gegenbauer operator", *Transactions of A. Razmadze Mathematical Institute*, 2016

Publication

<1%

70

Haibo LIN, Dachun YANG. "Pointwise multipliers for localized morrey-campanato spaces on RD-spaces", *Acta Mathematica Scientia*, 2014

Publication

<1%

71

*Analysis Partial Differential Equations and Applications*, 2009.

Publication

<1%

72

Guliyev, Vagif S., and Lubomira G. Softova. "Global Regularity in Generalized Morrey Spaces of Solutions to Nondivergence Elliptic Equations with VMO Coefficients", Potential Analysis, 2013.

Publication

<1%

73

Yoshihiro Sawano. "Generalized Morrey Spaces for Non-doubling Measures", Nonlinear Differential Equations and Applications NoDEA, 12/2008

Publication

<1%

74

Applied and Numerical Harmonic Analysis, 2015.

Publication

<1%

75

Yoshihiro Sawano. "Atomic Decompositions of Hardy Spaces with Variable Exponents and its Application to Bounded Linear Operators", Integral Equations and Operator Theory, 2013

Publication

<1%

76

[math.usm.my](http://math.usm.my)

Internet Source

<1%

77

[facta.junis.ni.ac.rs](http://facta.junis.ni.ac.rs)

Internet Source

<1%

78

[media.neliti.com](http://media.neliti.com)

Internet Source

<1%

79

Dag Lukkassen, Lars-Erik Persson, Natasha Samko. "Hardy Type Operators In Local Vanishing Morrey Spaces On Fractal Sets", Fractional Calculus and Applied Analysis, 2015

Publication

<1%

80

Komori-Furuya, Yasuo, Katsuo Matsuoka, Eiichi Nakai, and Yoshihiro Sawano. "Applications of Littlewood-Paley Theory for  $\lambda$ -Morrey Spaces to the Boundedness of Integral Operators", Journal

<1%

## of Function Spaces and Applications, 2013.

Publication

81

Yoshihiro Sawano. "Besov-Morrey spaces and Triebel-Lizorkin-Morrey spaces on domains", *Mathematische Nachrichten*, 08/18/2010

Publication

<1%

82

[www.math.sns.it](http://www.math.sns.it)

Internet Source

<1%

83

Hakim, Denny Ivanal, and Yoshihiro Sawano. "Interpolation of generalized Morrey spaces", *Revista Matemática Complutense*, 2016.

Publication

<1%

84

Yoshihiro Sawano, Tetsu Shimomura. "Sobolev embeddings for Riesz potentials of functions in non-doubling Morrey spaces of variable exponents", *Collectanea Mathematica*, 2013

Publication

<1%

85

Fumi-Yuki Maeda. "Trudinger's Inequality and Continuity of Potentials on Musielak–Orlicz–Morrey Spaces", *Potential Analysis*, 05/01/2012

Publication

<1%

86

Gala, Sadek, Yoshihiro Sawano, and Hitoshi Tanaka. "A new Beale-Kato-Majda criteria for the 3D magneto-micropolar fluid equations in the Orlicz-Morrey space", *Mathematical Methods in the Applied Sciences*, 2012.

Publication

<1%

87

Hendra Gunawan, Eder Kikianty, Christopher Schwanke. "Discrete Morrey spaces and their inclusion properties", *Mathematische Nachrichten*, 2017

Publication

<1%

88

Sadek Gala, Yoshihiro Sawano, Hitoshi Tanaka. "A remark on two generalized Orlicz–Morrey spaces", *Journal of Approximation Theory*, 2015

<1%

89 GuliyeV, Vagif, Shamsiyya Muradova, Mehriban Omarova, and Lubomira Softova. "Gradient estimates for parabolic equations in generalized weighted Morrey spaces", Acta Mathematica Sinica English Series, 2016.

Publication

90 Lucas B. Oliveira, Guilherme S. Oliveira, Jackson Piazza, Murilo Cervi, Ricardo N. Prado, Alysson R. Seidel. "Fixed Frequency Self-Oscillating Electronic Ballast Design Procedure", 2008 IEEE Industry Applications Society Annual Meeting, 2008

Publication

91 "Mathematical Analysis and Applications—Plenary Lectures", Springer Nature America, Inc, 2018

Publication

92 Noriko Mizoguchi. "Non-existence of radial backward self-similar blow-up solutions with sign change", Proceedings of the Royal Society of Edinburgh Section A Mathematics, 08/2011

Publication

93 Lin, Haibo, Suqing Wu, and Dachun Yang. "Boundedness of certain commutators over non-homogeneous metric measure spaces", Analysis and Mathematical Physics, 2016.

Publication

94 GuliyeV, Vagif S., Fatih Deringoz, and Javanshir J. Hasanov. " $(\Phi, \Psi)$ -admissible potential operators and their commutators on vanishing Orlicz-Morrey spaces", Collectanea mathematica, 2015.

Publication

95 Humberto Rafeiro, Stefan Samko. "Fractional

---

integrals and derivatives: mapping properties",  
Fractional Calculus and Applied Analysis, 2016

Publication

---

<1%

---

Exclude quotes      Off

Exclude matches      Off

Exclude bibliography      On

# CHARACTERIZATIONS FOR THE GENERALIZED FRACTIONAL INTEGRAL OPERATORS ON MORREY SPACES

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---

PAGE 11

---

PAGE 12

---

PAGE 13

---

PAGE 14

---

PAGE 15

---

PAGE 16

---

PAGE 17

---