

141
accesses
0
downloads



Institute of Mathematics

of the Czech Academy of Sciences

Mathematica Bohemica, Vol. 143, No. 4, pp. 409-417, 2018

Norm estimates for Bessel-Riesz operators on generalized Morrey spaces

Mochammad Idris, Hendra Gunawan, Eridani

Received May 11, 2017. First published March 7, 2018.

Abstract: We revisit the properties of Bessel-Riesz operators and present a different proof of the boundedness of these operators on generalized Morrey spaces. We also obtain an estimate for the norm of these operators on generalized Morrey spaces in terms of the norm of their kernels on an associated Morrey space. As a consequence of our results, we reprove the boundedness of fractional integral operators on generalized Morrey spaces, especially of exponent 1 , and obtain a new estimate for their norm.

Keywords: Bessel-Riesz operator; fractional integral operator; generalized Morrey space

Classification MSC: 42B20, 26A33, 42B25, 26D10

DOI: 10.21136/MB.2018.0045-17

PDF available at: [Institute of Mathematics CAS](#) [Digital Mathematics Library](#)

References:

- [1] *D. R. Adams*: A note on Riesz potentials. Duke Math. J. 42 (1975), 765-778. [DOI 10.1215/S0012-7094-75-04265-9](https://doi.org/10.1215/S0012-7094-75-04265-9) | [MR 0458158](https://mr.ams.org/) | [Zbl 0336.46038](https://zbmath.org/)
- [2] *F. Chiarenza, M. Frasca*: Morrey spaces and Hardy-Littlewood maximal function. Rend. Mat. Appl., VII. Ser. 7 (1987), 679-693. [MR 0985999](https://mr.ams.org/) | [Zbl 0717.42023](https://zbmath.org/)
- [3] *L. Grafakos*: Classical Fourier Analysis. Graduate Texts in Mathematics 249. Springer, New York (2008). [DOI 10.1007/978-0-387-09432-8](https://doi.org/10.1007/978-0-387-09432-8) | [MR 2445437](https://mr.ams.org/) | [Zbl 1220.42001](https://zbmath.org/)
- [4] *H. Gunawan, Eridani*: Fractional integrals and generalized Olsen inequalities. Kyungpook Math. J. 49 (2009), 31-39. [DOI 10.5666/KMJ.2009.49.1.031](https://doi.org/10.5666/KMJ.2009.49.1.031) | [MR 2527370](https://mr.ams.org/) | [Zbl 1181.26016](https://zbmath.org/)
- [5] *H. Gunawan, D. I. Hakim, K. M. Limanta, A. A. Masta*: Inclusion properties of generalized Morrey spaces. Math. Nachr. 290 (2017), 332-340. [DOI 10.1002/mana.201500425](https://doi.org/10.1002/mana.201500425) | [MR 3607107](https://mr.ams.org/) | [Zbl 1361.42023](https://zbmath.org/)
- [6] *G. H. Hardy, J. E. Littlewood*: Some properties of fractional integrals I. Math. Z. 27 (1928), 565-606. [DOI 10.1007/BF01171116](https://doi.org/10.1007/BF01171116) | [MR 1544927](https://mr.ams.org/) | [JFM 54.0275.05](https://jfm.cambridge.org/)
- [7] *G. H. Hardy, J. E. Littlewood*: Some properties of fractional integrals II. Math. Z. 34 (1932), 403-439. [DOI 10.1007/BF01180596](https://doi.org/10.1007/BF01180596) | [MR 1545260](https://mr.ams.org/) | [Zbl 0003.15601](https://zbmath.org/)
- [8] *M. Idris, H. Gunawan, Eridani*: The boundedness of Bessel-Riesz operators on generalized Morrey spaces. Aust. J. Math. Anal. Appl. 13 (2016), Article ID 9, 10 pages. [MR 3518943](https://mr.ams.org/) | [Zbl 1342.42015](https://zbmath.org/)
- [9] *M. Idris, H. Gunawan, J. Lindiarni, Eridani*: The boundedness of Bessel-Riesz operators on Morrey spaces. Int. Symp. On Current Progress in Mathematics and Sciences 2015. AIP Conf. Proc 1729 (2016), Article No. 020006. [DOI 10.1063/1.4946909](https://doi.org/10.1063/1.4946909)
- [10] *E. H. Lieb, M. Loss*: Analysis. Graduate Studies in Mathematics 14. AMS, Providence (2001). [DOI 10.1090/gsm/014](https://doi.org/10.1090/gsm/014) | [MR 1817225](https://mr.ams.org/) | [Zbl 0966.26002](https://zbmath.org/)
- [11] *E. Nakai*: Hardy-Littlewood maximal operator, singular integral operators and the Riesz potentials on generalized Morrey spaces. Math. Nachr. 166 (1994), 95-103. [DOI 10.1002/mana.19941660108](https://doi.org/10.1002/mana.19941660108) | [MR 1273325](https://mr.ams.org/) | [Zbl 0837.42008](https://zbmath.org/)
- [12] *J. Peetre*: On the theory of $\mathcal{L}_{p,\lambda}$ spaces. J. Functional Analysis 4 (1969), 71-87. [DOI 10.1016/0022-1236\(69\)90022-6](https://doi.org/10.1016/0022-1236(69)90022-6) | [MR 0241965](https://mr.ams.org/) | [Zbl 0175.42602](https://zbmath.org/)
- [13] *S. L. Sobolev*: On a theorem of functional analysis. Am. Math. Soc., Transl., II. Ser. 34 (1963), 39-68 Translated from Mat. Sb., N. Ser. 4 (1938), 471-497. [DOI 10.1090/trans2/034/02](https://doi.org/10.1090/trans2/034/02) | [Zbl 0131.11501](https://zbmath.org/)

Affiliations: *Mochammad Idris*, Department of Mathematics, Institute of Technology Bandung, Bandung 40132, Indonesia, e-mail: idemath@gmail.com (*permanent address*: Department of Mathematics, Lambung Mangkurat University, Banjarbaru Campus, Banjarbaru 70714, Indonesia); *Hendra Gunawan*, Department of Mathematics, Institute of Technology Bandung, Bandung 40132, Indonesia, e-mail: hgunawan@math.itb.ac.id; *Eridani*, Department of Mathematics, Airlangga University, Campus C Mulyorejo, Surabaya 60115, Indonesia, e-mail: eridani.dinadewi@gmail.com

[\[List of online first articles\]](#) [\[Contents of Mathematica Bohemica\]](#)
[\[Full text of the older issues of Mathematica Bohemica at DML-CZ\]](#)

© [Institute of Mathematics CAS](#), 2020

PDF available at:



[Institute of Mathematics CAS](#)

free access

...



[Digital Mathematics Library](#)

fulltext not available (moving wall 0 months)

NORM ESTIMATES FOR BESSEL-RIESZ OPERATORS ON
GENERALIZED MORREY SPACES

MOCHAMMAD IDRIS, HENDRA GUNAWAN, Bandung,
ERIDANI, Surabaya

Received May 11, 2017. First published March 7, 2018.
Communicated by Dagmar Medková

Abstract. We revisit the properties of Bessel-Riesz operators and present a different proof of the boundedness of these operators on generalized Morrey spaces. We also obtain an estimate for the norm of these operators on generalized Morrey spaces in terms of the norm of their kernels on an associated Morrey space. As a consequence of our results, we reprove the boundedness of fractional integral operators on generalized Morrey spaces, especially of exponent 1, and obtain a new estimate for their norm.

Keywords: Bessel-Riesz operator; fractional integral operator; generalized Morrey space

MSC 2010: 42B20, 26A33, 42B25, 26D10

1. INTRODUCTION

Integral operators such as maximal operators and fractional integral operators have been studied extensively in the last four decades. Here we are interested in Bessel-Riesz operators, which are related to fractional integral operators. Let $0 < \alpha < n$ and $\gamma \geq 0$. The operator $I_{\alpha,\gamma}$ which maps every $f \in L_{\text{loc}}^p(\mathbb{R}^n)$, $1 \leq p < \infty$, to

$$I_{\alpha,\gamma}f(x) := \int_{\mathbb{R}^n} K_{\alpha,\gamma}(x-y)f(y) \, dy = K_{\alpha,\gamma} * f(x), \quad x \in \mathbb{R}^n,$$

where $K_{\alpha,\gamma}(x) := |x|^{\alpha-n}(1+|x|)^{-\gamma}$, is called *Bessel-Riesz operator*, and the kernel $K_{\alpha,\gamma}$ is called *Bessel-Riesz kernel*. The boundedness of these operators on Morrey spaces and on generalized Morrey spaces was studied in [8] and [9].

The first and second authors are supported by ITB Research & Innovation Program 2016.

Let $1 \leq p < \infty$ and $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be of class \mathcal{G}_p , that is, φ is almost decreasing (there exists $C > 0$ such that $\varphi(r) \geq C\varphi(s)$ for $r \leq s$) and $\varphi^p(r)r^n$ is almost increasing (there exists $C > 0$ such that $\varphi^p(r)r^n \leq C\varphi^p(s)s^n$ for $r \leq s$). Clearly if φ is of class \mathcal{G}_p , then φ satisfies the *doubling condition*, that is, there exists $C > 0$ such that $C^{-1} \leq \varphi(r)/\varphi(s) \leq C$ whenever $1 \leq rs^{-1} \leq 2$. We define the *generalized Morrey space* $L^{p,\varphi}(\mathbb{R}^n)$ to be the set of all functions $f \in L^p_{\text{loc}}(\mathbb{R}^n)$ for which

$$\|f\|_{L^{p,\varphi}} := \sup_{B=B(a,r)} \frac{1}{\varphi(r)} \left(\frac{1}{|B|} \int_B |f(x)|^p dx \right)^{1/p} < \infty,$$

where $|B|$ denotes the Lebesgue measure of B . (Recall that the Lebesgue measure of $B = B(a, r)$ is $|B(a, r)| = C_n r^n$ for every $a \in \mathbb{R}^n$ and $r > 0$, where $C_n > 0$ depends only on n .)

If $1 \leq p \leq q < \infty$ and $\varphi(r) := C_n r^{-n/q}$, $r > 0$, then $L^{p,\varphi}(\mathbb{R}^n)$ is the classical Morrey space $L^{p,q}(\mathbb{R}^n)$, which is equipped with

$$\|f\|_{L^{p,q}} := \sup_{B=B(a,r)} |B|^{1/q-1/p} \left(\int_B |f(x)|^p dx \right)^{1/p}.$$

Particularly, for $p = q$, $L^{p,p}(\mathbb{R}^n)$ is the Lebesgue space $L^p(\mathbb{R}^n)$.

In [9], we know that for $\gamma > 0$, $K_{\alpha,\gamma}$ is a member of $L^t(\mathbb{R}^n)$ spaces for some values of t depending on α and γ . It follows from Young's inequality (see [3]) that

$$\|I_{\alpha,\gamma} f\|_{L^q} \leq \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^p}, \quad f \in L^p(\mathbb{R}^n)$$

whenever $1 \leq p < t'$, $1/q = 1/p - 1/t'$ (where t' denotes the dual exponent of t) and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$. This tells us that $I_{\alpha,\gamma}$ is bounded from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$ with $\|I_{\alpha,\gamma}\|_{L^p \rightarrow L^q} \leq \|K_{\alpha,\gamma}\|_{L^t}$. In [8], it is also shown that $I_{\alpha,\gamma}$ is bounded on generalized Morrey spaces but without a good estimate for its norm as on Morrey spaces. We shall now refine the results by estimating the norms of the operators more carefully through the membership of K_α in Morrey spaces.

Note that for $\gamma = 0$, $I_{\alpha,0} = I_\alpha$ is the *fractional integral operator* with kernel $K_\alpha(x) := |x|^{\alpha-n}$. Hardy and Littlewood [6], [7] and Sobolev [13] proved the boundedness of I_α on Lebesgue spaces. The boundedness of I_α on Morrey spaces is proved by Peetre [12], and improved by Adams [1] and Chiarenza and Frasca [2]. Later, Nakai [11] obtained the boundedness of I_α on generalized Morrey spaces, which can be viewed as an extension of Spanne's result. In 2009, Gunawan and Eridani [4] proved the boundedness of I_α on generalized Morrey spaces which extends Adams' and Chiarenza-Frasca's results.

In this paper, we give a new proof of the boundedness of $I_{\alpha,\gamma}$ on generalized Morrey spaces. At the same time, an upper bound for the norm of the operators is obtained. As a consequence of our result, we have an estimate for the norm of I_α (from a generalized Morrey space to another) in terms of the norm of K_α on the associated Morrey space. A lower bound for the norm of the operators is discussed in Section 3.

2. THE BOUNDEDNESS OF $I_{\alpha,\gamma}$ ON GENERALIZED MORREY SPACES

We begin with a lemma about the membership of K_α in some Morrey spaces. Note that throughout this paper, the letters C and C_k denote constants which may change from line to line.

Lemma 2.1. *If $0 < \alpha < n$, then $K_\alpha \in L^{s,t}(\mathbb{R}^n)$, where $1 \leq s < t = n/(n - \alpha)$.*

Proof. Let $0 < \alpha < n$. Take an arbitrary $B = B(a, R)$, where $a \in \mathbb{R}^n$ and $R > 0$. For $1 \leq s < t = n/(n - \alpha)$ we observe that

$$\begin{aligned} |B|^{s/t-1} \int_B K_\alpha^s(x) \, dx &\leq |B(0, R)|^{s/t-1} \int_{B(0, R)} |x|^{(\alpha-n)s} \, dx \\ &\leq CR^{n(s/t-1)} R^{n(1-s/t)} = C. \end{aligned}$$

By taking the supremum over $B = B(a, R)$ we obtain $\|K_\alpha\|_{L^{s,t}}^s \leq C$. Hence $K_\alpha \in L^{s,t}(\mathbb{R}^n)$. \square

Remark 2.2. For $0 < \alpha < n$ and $\gamma > 0$ we know that $K_{\alpha,\gamma} \in L^t(\mathbb{R}^n)$ for $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$, see [9]. By the inclusion property of Morrey spaces (see [5]) we have $K_{\alpha,\gamma} \in L^t(\mathbb{R}^n) = L^{t,t}(\mathbb{R}^n) \subseteq L^{s,t}(\mathbb{R}^n)$ for $1 \leq s \leq t$ and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$. Moreover, because $K_{\alpha,\gamma}(x) \leq K_\alpha(x)$ for every $x \in \mathbb{R}^n$, $K_{\alpha,\gamma}$ is also contained in $L^{s,t}(\mathbb{R}^n)$ for $1 \leq s < t = n/(n - \alpha)$.

As a counterpart of the results in [8] and [9], we have the following theorem on the boundedness of $I_{\alpha,\gamma}$ on Morrey spaces. Note particularly that the estimate holds for $p_1 = 1$.

Theorem 2.3. *If $0 < \alpha < n$ and $\gamma \geq 0$, then $I_{\alpha,\gamma}$ is bounded from $L^{p_1,q_1}(\mathbb{R}^n)$ to $L^{p_2,q_2}(\mathbb{R}^n)$ with*

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,q_2}} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}}, \quad f \in L^{p_1,q_1}(\mathbb{R}^n)$$

whenever $1 \leq p_1 \leq q_1 < n/\alpha$, $1/p_2 = 1/p_1 - 1/s'$, and $1/q_2 = 1/q_1 - 1/t'$, with $1 \leq s < t = n/(n - \alpha)$ for $\gamma \geq 0$ or $1 \leq s \leq t$ and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$ for $\gamma > 0$.

Theorem 2.3 is in fact a special case of the boundedness of $I_{\alpha,\gamma}$ on generalized Morrey spaces, which is stated in the following theorem.

Theorem 2.4. *Let $0 < \alpha < n$ and $\gamma \geq 0$. If $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is of class \mathcal{G}_{p_1} such that $\int_R^\infty \varphi(r)r^{n/t'-1} dr \leq C\varphi(R)R^{n/t'}$ for every $R > 0$, then $I_{\alpha,\gamma}$ is bounded from $L^{p_1,\varphi}(\mathbb{R}^n)$ to $L^{p_2,\psi}(\mathbb{R}^n)$, where $\psi(r) := \varphi(r)r^{n/t'}$, with*

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C\|K_{\alpha,\gamma}\|_{L^{s,t}}\|f\|_{L^{p_1,\varphi}}, \quad f \in L^{p_1,\varphi}(\mathbb{R}^n)$$

whenever $1 \leq p_1 < n/\alpha$ and $1/p_2 = 1/p_1 - 1/s'$, with $1 \leq s < t = n/(n - \alpha)$ for $\gamma \geq 0$ or $1 \leq s \leq t$ and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$ for $\gamma > 0$.

Proof. Suppose that $\gamma > 0$ and all the hypotheses hold. For $f \in L^{p_1,\varphi}(\mathbb{R}^n)$ and $B = B(a, R)$, where $a \in \mathbb{R}^n$ and $R > 0$, write

$$f := f_1 + f_2 := f_{\chi_{\tilde{B}}} + f_{\chi_{\tilde{B}^c}},$$

where $\tilde{B} = B(a, 2R)$ and \tilde{B}^c denotes its complement. To estimate $I_{\alpha,\gamma}f_1$, we observe that for every $x \in B$, Hölder's inequality gives

$$\begin{aligned} |I_{\alpha,\gamma}f_1(x)| &\leq \int_{\tilde{B}} K_{\alpha,\gamma}(x-y)|f(y)| dy \\ &= \int_{\tilde{B}} K_{\alpha,\gamma}^{s/p_2}(x-y)|f(y)|^{p_1/p_2} K_{\alpha,\gamma}^{(p_2-s)/p_2}(x-y)|f(y)|^{(p_2-p_1)/p_2} dy \\ &\leq \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y)|f(y)|^{p_1} dy \right)^{1/p_2} \\ &\quad \times \left(\int_{\tilde{B}} K_{\alpha,\gamma}^{(p_2-s)/(p_2-1)}(x-y)|f(y)|^{(p_2-p_1)/(p_2-1)} dy \right)^{1/p_2'}. \end{aligned}$$

Meanwhile, we have

$$\begin{aligned} \int_{\tilde{B}} K_{\alpha,\gamma}^{(p_2-s)/(p_2-1)}(x-y)|f(y)|^{(p_2-p_1)/(p_2-1)} dy \\ \leq \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y) dy \right)^{p_2'(1/s-1/p_2)} \left(\int_{\tilde{B}} |f(y)|^{p_1} dy \right)^{p_2'/s'}. \end{aligned}$$

Therefore we obtain

$$\begin{aligned} |I_{\alpha,\gamma}f_1(x)| &\leq \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y)|f(y)|^{p_1} dy \right)^{1/p_2} \\ &\quad \times \left(\int_{\tilde{B}} K_{\alpha,\gamma}^s(x-y) dy \right)^{1/s-1/p_2} \left(\int_{\tilde{B}} |f(y)|^{p_1} dy \right)^{1/s'}. \end{aligned}$$

$$\begin{aligned} &\leq \left(\int_{\bar{B}} K_{\alpha,\gamma}^s(x-y) |f(y)|^{p_1} dy \right)^{1/p_2} \\ &\quad \times CR^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'}. \end{aligned}$$

We then take the p_2 th power and integrate both sides over B to get

$$\begin{aligned} &\int_B |I_{\alpha,\gamma} f_1(x)|^{p_2} dx \\ &\leq \int_B \int_{\bar{B}} K_{\alpha,\gamma}^s(x-y) |f(y)|^{p_1} dy dx \\ &\quad \times (CR^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'})^{p_2}. \end{aligned}$$

By Fubini's theorem we have

$$\begin{aligned} &\int_B |I_{\alpha,\gamma} f_1(x)|^{p_2} dx \\ &\leq \int_{\bar{B}} |f(y)|^{p_1} \left(\int_B K_{\alpha,\gamma}^s(x-y) dx \right) dy \\ &\quad \times (CR^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'})^{p_2} \\ &\leq CR^{n(1-s/t)} \|K_{\alpha,\gamma}\|_{L^{s,t}}^s \int_{\bar{B}} |f(y)|^{p_1} dy \\ &\quad \times (R^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'})^{p_2} \\ &\leq CR^{n(1-s/t)+n} \varphi^{p_1}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^s \|f\|_{L^{p_1,\varphi}}^{p_1} \\ &\quad \times (R^{n(1-s/t)(1/s-1/p_2)+n/s'} \varphi^{p_1/s'}(2R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{1-s/p_2} \|f\|_{L^{p_1,\varphi}}^{p_1/s'})^{p_2} \\ &\leq C|B|\psi^{p_2}(R) \|K_{\alpha,\gamma}\|_{L^{s,t}}^{p_2} \|f\|_{L^{p_1,\varphi}}^{p_2}, \end{aligned}$$

whence

$$\frac{1}{\psi(R)} \left(\frac{1}{|B|} \int_B |I_{\alpha,\gamma} f_1(x)|^{p_2} dx \right)^{1/p_2} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}}.$$

Next, we estimate $I_{\alpha,\gamma} f_2$. For every $x \in B = B(a, R)$ we observe that

$$\begin{aligned} |I_{\alpha,\gamma} f_2(x)| &\leq \int_{\bar{B}^c} K_{\alpha,\gamma}(x-y) |f(y)| dy \\ &\leq \int_{|x-y| \geq R} K_{\alpha,\gamma}(x-y) |f(y)| dy \\ &= \sum_{k=0}^{\infty} \int_{2^k R \leq |x-y| < 2^{k+1} R} K_{\alpha,\gamma}(x-y) |f(y)| dy \\ &\leq \sum_{k=0}^{\infty} K_{\alpha,\gamma}(2^k R) \int_{2^k R \leq |x-y| < 2^{k+1} R} |f(y)| dy \end{aligned}$$

$$\begin{aligned}
&\leq C \sum_{k=0}^{\infty} K_{\alpha,\gamma}(2^k R)(2^k R)^{n/p_1'} \left(\int_{2^k R \leq |x-y| < 2^{k+1} R} |f(y)|^{p_1} dy \right)^{1/p_1} \\
&\leq C \|f\|_{L^{p_1,\varphi}} \sum_{k=0}^{\infty} K_{\alpha,\gamma}(2^k R)(2^k R)^n \varphi(2^k R).
\end{aligned}$$

For every $k \in \mathbb{Z}$ we have

$$\begin{aligned}
K_{\alpha,\gamma}(2^k R) &\leq C(2^k R)^{-n/s} \left(\int_{2^k R \leq |x-y| < 2^{k+1} R} K_{\alpha,\gamma}^s(x-y) dy \right)^{1/s} \\
&\leq C(2^k R)^{-n/t} \|K_{\alpha,\gamma}\|_{L^{s,t}}.
\end{aligned}$$

Since $\int_R^\infty \varphi(r)r^{n/t'-1} dr \leq C\varphi(R)R^{n/t'}$, we get

$$\begin{aligned}
|I_{\alpha,\gamma}f_2(x)| &\leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}} \sum_{k=0}^{\infty} (2^k R)^{n/t'} \varphi(2^k R) \\
&\leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}} \int_R^\infty \varphi(r)r^{n/t'-1} dr \\
&\leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}} \varphi(R)R^{n/t'} \\
&= C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}} \psi(R).
\end{aligned}$$

Raising to the p_2 th power and integrating over B we obtain

$$\int_B |I_{\alpha,\gamma}f_2(x)|^{p_2} dx \leq C (\|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}})^{p_2} \psi^{p_2}(R) |B|,$$

whence

$$\frac{1}{\psi(R)} \left(\frac{1}{|B|} \int_B |I_{\alpha,\gamma}f_2(x)|^{p_2} dx \right)^{1/p_2} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}}.$$

Combining the two estimates for $I_{\alpha,\gamma}f_1$ and $I_{\alpha,\gamma}f_2$ we obtain

$$\frac{1}{\psi(R)} \left(\frac{1}{|B|} \int_B |I_{\alpha,\gamma}f(x)|^{p_2} dx \right)^{1/p_2} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}}.$$

Since this inequality holds for every $a \in \mathbb{R}^n$ and $R > 0$, it follows that

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\varphi}},$$

as desired.

We may repeat the same argument and use Lemma 2.1 to obtain the same inequality for the case where $\gamma = 0$ and $1 \leq s < t = n/(n - \alpha)$. \square

Remark 2.5. Theorems 2.3 and 2.4 give us upper estimates for the norm of the Bessel-Riesz operators (from one Morrey space to another). In particular, for $\gamma = 0$ we have an estimate for the norm of the fractional integral operator I_α in terms of the norm of its kernel (on the associated Morrey space), which follows from the inequality

$$\|I_\alpha f\|_{L^{p_2, \psi}} \leq C \|K_\alpha\|_{L^{s, t}} \|f\|_{L^{p_1, \varphi}}$$

for $1 \leq p_1 < n/\alpha$ and $1/p_2 = 1/p_1 - 1/s'$, with $1 \leq s < t = n/(n - \alpha)$.

In the following section, we discuss lower estimates for the norm of the operators in terms of the norm of the Bessel-Riesz kernel (on some Morrey spaces).

3. AN ESTIMATE FOR THE NORM OF THE OPERATORS

Recall that if $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ are normed spaces and if $T: (X, \|\cdot\|_X) \rightarrow (Y, \|\cdot\|_Y)$ is a linear operator, then the norm of T (from X to Y) is defined as

$$\|T\|_{X \rightarrow Y} := \sup_{f \neq 0} \frac{\|Tf\|_Y}{\|f\|_X}.$$

Knowing that the Bessel-Riesz operator $I_{\alpha, \gamma}$ is a linear operator on Morrey spaces, we would like to estimate the norm of $I_{\alpha, \gamma}$ from a (generalized) Morrey space to another. We obtain the following result.

Theorem 3.1. *Let $0 < \alpha < n$, $\gamma \geq 0$, and φ be of class \mathcal{G}_{p_1} where $1 \leq p_1 < n/\alpha$. If $\varphi(r)r^n$ is almost increasing and for every $R > 0$ we have*

- (i) $\int_R^\infty \varphi(r)r^{n/t'-1} dr \leq C_1 \varphi(R)R^{n/t'}$,
- (ii) $\int_0^R \varphi^{p_1}(r)r^{n-1} dr \leq C_2 \varphi^{p_1}(R)R^n$, and
- (iii) $\int_0^R r^{n-1}/\varphi^{s'}(r)r^{ns'} dr \leq C_3 R^n/\varphi^{s'}(R)R^{ns'}$, where $1 \leq p_1 < t$ and $1 < s < t = n/(n - \alpha)$ for $\gamma \geq 0$ or $1 \leq p_1 \leq t$, $1 < s \leq t$, and $n/(n + \gamma - \alpha) < t < n/(n - \alpha)$ for $\gamma > 0$,

then we have

$$C_4 \|K_{\alpha, \gamma}\|_{L^{p_1, t}} \leq \|I_{\alpha, \gamma}\|_{L^{p_1, \varphi} \rightarrow L^{p_2, \psi}} \leq C_5 \|K_{\alpha, \gamma}\|_{L^{s, t}}$$

whenever $1/p_2 = 1/p_1 - 1/s'$ and $\psi(r) := \varphi(r)r^{n/t'}$. In particular, for $\gamma = 0$, $1 \leq p_1 < t$ and $1 < s < t = n/(n - \alpha)$ we have

$$C_4 \|K_\alpha\|_{L^{p_1, t}} \leq \|I_\alpha\|_{L^{p_1, \varphi} \rightarrow L^{p_2, \psi}} \leq C_5 \|K_\alpha\|_{L^{s, t}}$$

whenever $1/p_2 = 1/p_1 - 1/s'$ and $\psi(r) := \varphi(r)r^{n/t'}$.

Proof. Suppose that $\gamma > 0$ and all the hypotheses hold. By Theorem 2.4 we already have

$$\|I_{\alpha,\gamma}\|_{L^{p_1,\varphi}\rightarrow L^{p_2,\psi}} \leq C\|K_{\alpha,\gamma}\|_{L^{s,t}}.$$

To prove the lower estimate, put $\varrho(r) := \varphi(r)r^n$. Let $B = B(a, R)$, where $a \in \mathbb{R}^n$ and $R > 0$. By our assumptions on φ we have

$$|B|^{1/t}\psi(R)\left(\frac{1}{|B|}\int_B \varrho^{-s'}(|x|) dx\right)^{1/s'} \leq C\varphi(R)R^{n/s}\left(\int_0^R \frac{r^{n-1}}{\varphi^{s'}(r)r^{ns'}} dr\right)^{1/s'} \leq C.$$

Now take $f_0(x) := \varphi(|x|)$. Here $\|f_0\|_{L^{p_1,\varphi}} \approx 1$. Moreover, one may compute that

$$I_{\alpha,\gamma}f_0(x) \geq \int_{B(x,2|x|)} K_{\alpha,\gamma}(x-y)f_0(y) dy \geq CK_{\alpha,\gamma}(2x)\varphi(|x|)|x|^n = C\varrho(|x|)K_{\alpha,\gamma}(x)$$

for every $x \in \mathbb{R}^n$. It follows that

$$\|\varrho(\cdot)K_{\alpha,\gamma}\|_{L^{p_2,\psi}} \leq C\|I_{\alpha,\gamma}f_0\|_{L^{p_2,\psi}} \leq C\|I_{\alpha,\gamma}\|_{L^{p_1,\varphi}\rightarrow L^{p_2,\psi}}.$$

Next, by Hölder's inequality we have

$$\left(\int_B K_{\alpha,\gamma}^{p_1}(x) dx\right)^{1/p_1} \leq \left(\int_B \varrho^{-s'}(|x|) dx\right)^{1/s'} \left(\int_B (\varrho(|x|)K_{\alpha,\gamma}(x))^{p_2} dx\right)^{1/p_2},$$

whence

$$\begin{aligned} |B|^{1/t-1/p_1}\left(\int_B K_{\alpha,\gamma}^{p_1}(x) dx\right)^{1/p_1} &\leq |B|^{1/t}\psi(R)\left(\frac{1}{|B|}\int_B \varrho^{-s'}(|x|) dx\right)^{1/s'} \\ &\quad \times \frac{1}{\psi(R)}\left(\frac{1}{|B|}\int_B (\varrho(|x|)K_{\alpha,\gamma}(x))^{p_2} dx\right)^{1/p_2} \\ &\leq C\|I_{\alpha,\gamma}\|_{L^{p_1,\varphi}\rightarrow L^{p_2,\psi}}. \end{aligned}$$

By taking the supremum over $B = B(a, R)$ we conclude that

$$C\|K_{\alpha,\gamma}\|_{L^{p_1,t}} \leq \|I_{\alpha,\gamma}\|_{L^{p_1,\varphi}\rightarrow L^{p_2,\psi}},$$

as desired. The same argument applies for the case where $\gamma = 0$ with $1 \leq p_1 < t$ and $1 < s < t = n/(n - \alpha)$. \square

Remark 3.2. One may observe that the constants C_4 and C_5 in Theorem 3.1 depend on φ , n , p_1 , s , and t , but not on α and γ . Although the lower and the upper bound are not comparable, we may still get useful information from these estimates,

especially for the norm of the operator I_α from $L^{p_1, \varphi}(\mathbb{R}^n)$ to $L^{p_2, \psi}(\mathbb{R}^n)$. Observe that for $1 \leq p_1 < t = n/(n - \alpha)$ we have $\|K_\alpha\|_{L^{p_1, t}}^{p_1} = C/((\alpha - n)p_1 + n) \geq C/\alpha$. Hence, if all the hypotheses in Theorem 3.1 hold for the case where $\gamma = 0$, then we obtain $\|I_\alpha\|_{L^{p_1, \varphi} \rightarrow L^{p_2, \psi}} \geq C/\alpha$, which blows up when $\alpha \rightarrow 0^+$. For $\varphi(r) := r^{-n/q_1}$ with $1 \leq p_1 < q_1 < \min\{s, n/\alpha\}$ and $1 < s < n/(n - \alpha)$, our result reduces to the estimate $\|I_\alpha\|_{L^{p_1, q_1} \rightarrow L^{p_2, q_2}} \geq C/\alpha$, where $1/p_2 = 1/p_1 - 1/s'$ and $1/q_2 = 1/q_1 - \alpha/n$. A similar behavior of the norm of I_α from $L^{p_1}(\mathbb{R}^n)$ to $L^{p_2}(\mathbb{R}^n)$ for $1/p_2 = 1/p_1 - \alpha/n$ when $\alpha \rightarrow 0^+$ is observed in [10], Chapter 4.

Acknowledgement. All authors would like to thank the anonymous referee for his/her careful reading and useful comments on the earlier version of this paper.

References

- [1] *D. R. Adams*: A note on Riesz potentials. *Duke Math. J.* *42* (1975), 765–778. [zbl](#) [MR](#) [doi](#)
- [2] *F. Chiarenza, M. Frasca*: Morrey spaces and Hardy-Littlewood maximal function. *Rend. Mat. Appl.*, VII. Ser. *7* (1987), 679–693. [zbl](#) [MR](#)
- [3] *L. Grafakos*: *Classical Fourier Analysis*. Graduate Texts in Mathematics 249. Springer, New York, 2008. [zbl](#) [MR](#) [doi](#)
- [4] *H. Gunawan, Eridani*: Fractional integrals and generalized Olsen inequalities. *Kyungpook Math. J.* *49* (2009), 31–39. [zbl](#) [MR](#) [doi](#)
- [5] *H. Gunawan, D. I. Hakim, K. M. Limanta, A. A. Masta*: Inclusion properties of generalized Morrey spaces. *Math. Nachr.* *290* (2017), 332–340. [zbl](#) [MR](#) [doi](#)
- [6] *G. H. Hardy, J. E. Littlewood*: Some properties of fractional integrals I. *Math. Z.* *27* (1928), 565–606. [zbl](#) [MR](#) [doi](#)
- [7] *G. H. Hardy, J. E. Littlewood*: Some properties of fractional integrals II. *Math. Z.* *34* (1932), 403–439. [zbl](#) [MR](#) [doi](#)
- [8] *M. Idris, H. Gunawan, Eridani*: The boundedness of Bessel-Riesz operators on generalized Morrey spaces. *Aust. J. Math. Anal. Appl.* *13* (2016), Article ID 9, 10 pages. [zbl](#) [MR](#)
- [9] *M. Idris, H. Gunawan, J. Lindiarni, Eridani*: The boundedness of Bessel-Riesz operators on Morrey spaces. *Int. Symp. On Current Progress in Mathematics and Sciences 2015*. AIP Conf. Proc. vol. 1729, 2016, Article No. 020006. [doi](#)
- [10] *E. H. Lieb, M. Loss*: *Analysis*. Graduate Studies in Mathematics 14. AMS, Providence, 2001. [zbl](#) [MR](#) [doi](#)
- [11] *E. Nakai*: Hardy-Littlewood maximal operator, singular integral operators and the Riesz potentials on generalized Morrey spaces. *Math. Nachr.* *166* (1994), 95–103. [zbl](#) [MR](#) [doi](#)
- [12] *J. Peetre*: On the theory of $\mathcal{L}_{p, \lambda}$ spaces. *J. Functional Analysis* *4* (1969), 71–87. [zbl](#) [MR](#) [doi](#)
- [13] *S. L. Sobolev*: On a theorem of functional analysis. *Am. Math. Soc.*, Transl., II. Ser. *34* (1963), 39–68; Translated from *Mat. Sb.*, N. Ser. *4* (1938), 471–497. [zbl](#) [doi](#)

Authors' addresses: Mochammad Idris, Department of Mathematics, Institute of Technology Bandung, Bandung 40132, Indonesia, e-mail: idemath@gmail.com (*permanent address*: Department of Mathematics, Lambung Mangkurat University, Banjarbaru Campus, Banjarbaru 70714, Indonesia); Hendra Gunawan, Department of Mathematics, Institute of Technology Bandung, Bandung 40132, Indonesia, e-mail: hgunawan@math.itb.ac.id; Eridani, Department of Mathematics, Airlangga University, Campus C Mulyorejo, Surabaya 60115, Indonesia, e-mail: eridani.dinadewi@gmail.com.

also developed by scimago:

 SCIMAGO INSTITUTIONS RANKINGS**SJR**

Scimago Journal & Country Rank

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Mathematica Bohemica

Country[Czech Republic](#) -  [SJR Ranking of Czech Republic](#)**Subject Area and Category**[Mathematics](#)
[Mathematics \(miscellaneous\)](#)**Publisher**[Akademie Ved Ceske Republiky](#)**Publication type**[Journals](#)**ISSN**

08627959

Coverage

2011-ongoing

 [Join the conversation about this journal](#)


6

H Index


Quartiles

+


Mathematics (miscellaneous)




● SJR



● Total Cites ● Self-Cites




Citations per document




- Cites / Doc. (4 years)
- Cites / Doc. (3 years)
- Cites / Doc. (2 years)


● External Cites per Doc ● Cites per Doc +




● % International Collaboration +



● Citable documents ● Non-citable documents +



● Cited documents ● Uncited documents +



Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimag
```

Mathematica Bohemica ←

Q3 Mathematics (miscellaneous)

best quartile

Leave a comment

Name

Email

(will not be published)

I'm not a robot



reCAPTCHA
Privacy - Terms

[Submit](#)

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:

ScopusFollow us on [@ScimagoJR](#)

Scimago Lab, Copyright 2007-2018. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)