

The Boundedness of Bessel- Riesz Operators On Morrey Spaces

by Eridani Eridani

Submission date: 14-Dec-2018 05:11PM (UTC+0800)

Submission ID: 1056967397

File name: 2016-AIPConfProc1729_IGLE_aip_proc.pdf (269.69K)

Word count: 2405

Character count: 10009



The boundedness of Bessel-Riesz operators on Morrey spaces

Mochammad Idris, Hendra Gunawan, Janny Lindiarni, and Eridani

Citation: *AIP Conference Proceedings* **1729**, 020006 (2016); doi: 10.1063/1.4946909

View online: <http://dx.doi.org/10.1063/1.4946909>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/1729?ver=pdfcov>

Published by the AIP Publishing

Articles you may be interested in

[Boundedness and invertibility of multidimensional integral operators with anisotropically homogeneous kernels in weighted \$L^p\$ -spaces](#)

AIP Conf. Proc. **1637**, 663 (2014); 10.1063/1.4904637

10

[Commutators of Hardy operators in vanishing Morrey spaces](#)

AIP Conf. Proc. **1493**, 859 (2012); 10.1063/1.4765588

2

[Boundedness of Weighted Singular Integral Operators on a Carleson Curve in Grand Lebesgue Spaces](#)

AIP Conf. Proc. **1281**, 490 (2010); 10.1063/1.3498517

12

[Boundedness properties of fermionic operators](#)

J. Math. Phys. **51**, 083503 (2010); 10.1063/1.3464264

2

[Boundedness of linear integral operators in weighted \$L^p\$ spaces](#)

J. Math. Phys. **16**, 1522 (1975); 10.1063/1.522702

The Boundedness of Bessel-Riesz Operators On Morrey Spaces

Mochammad Idris^{1,a)}, Hendra Gunawan¹, Janny Lindiarni¹ and Eridani²

¹Department of Mathematics, Institut Teknologi Bandung, Bandung 40132, Indonesia

²Department of Mathematics, Airlangga University, Campus C, Mulyorejo, Surabaya 60115, Indonesia

^{a)}Corresponding author: mochedris@students.itb.ac.id

Abstract. In this paper, we shall discuss about Bessel-Riesz operators. Kurata *et al.* have investigated their boundedness on generalized Morrey spaces with weight. The boundedness of these operators on Lebesgue spaces and Morrey spaces will be reproved using a different approach. Moreover, we also find the norm of the operators are bounded by the norm of the kernels.

Keywords: Bessel-Riesz operators, Hardy-Littlewood maximal operator, Morrey spaces.

INTRODUCTION

Let $0 < \gamma, 0 < \alpha < n$ and define

$$I_{\alpha,\gamma}f(x) := \int_{\mathbb{R}^n} K_{\alpha,\gamma}(x-y)f(y)dy$$

for every $f \in L^p_{loc}(\mathbb{R}^n)$, where $p \geq 1$, $K_{\alpha,\gamma}(x) := \frac{|x|^{n-\alpha}}{(1+|x|)^\gamma}$, $x \in \mathbb{R}^n$. Here, $K_{\alpha,\gamma}$ can be viewed as multiple of two kernels, $K_{\alpha,\gamma}(x) = J_\gamma(x)K_\alpha(x)$ for every $x \in \mathbb{R}^n$. In [1], J_γ and K_α are known as *Bessel kernel* and *Riesz kernel*. So, $K_{\alpha,\gamma}$ is called *Bessel-Riesz kernel* and $I_{\alpha,\gamma}$ is called *Bessel-Riesz operator*.

For $\gamma = 0$, we have $I_{\alpha,0} = I_\alpha$ (is called *fractional integral operator* or *Riesz potential* [1]). Studies about I_α were started since 1920's. Hardy-Littlewood [2, 3] and Sobolev [4] proved the boundedness of I_α on *Lebesgue spaces* through the inequality $\|I_\alpha f\|_{L^q} \leq C_p \|f\|_{L^p}$, for every $f \in L^p(\mathbb{R}^n)$, $1 < p < \frac{n}{\alpha}$, and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$.

For $1 \leq p \leq q$, the (classical) Morrey space $L^{p,q}(\mathbb{R}^n)$ is defined by

$$L^{p,q}(\mathbb{R}^n) := \{f \in L^p_{loc}(\mathbb{R}^n) : \|f\|_{L^{p,q}} < \infty\},$$

where $\|f\|_{L^{p,q}} := \sup_{r>0, a \in \mathbb{R}^n} r^{n(1/q-1/p)} \left(\int_{|x-a|<r} |f(x)|^p dx \right)^{1/p}$. We have an inclusion property for Morrey spaces $L^q(\mathbb{R}^n) = L^{q,q}(\mathbb{R}^n) \subseteq L^{p,q}(\mathbb{R}^n) \subseteq L^{1,q}(\mathbb{R}^n)$.

On Morrey spaces, Spanne [5] has shown that I_α is bounded form $L^{p_1,q_1}(\mathbb{R}^n)$ to $L^{p_2,q_2}(\mathbb{R}^n)$ for $1 < p_1 < q_1 < \frac{n}{\alpha}$, $\frac{1}{p_2} = \frac{1}{p_1} - \frac{\alpha}{n}$, and $\frac{1}{q_2} = \frac{1}{q_1} - \frac{\alpha}{n}$. Furthermore, Adams [6] and Chiarenza-Frasca [7] obtained a stronger result.

Theorem 1 [Adams, Chiarenza-Frasca] If $0 < \alpha < n$ then we have

$$\|I_\alpha f\|_{L^{p_2,q_2}} \leq C_{p_1,q_1} \|f\|_{L^{p_1,q_1}},$$

for every $f \in L^{p_1,q_1}(\mathbb{R}^n)$ where $1 < p_1 < q_1 < \frac{n}{\alpha}$, $\frac{1}{p_2} = \frac{1}{p_1} - \frac{\alpha}{n}$, and $\frac{1}{q_2} = \frac{1}{q_1} - \frac{\alpha}{n}$.

Meanwhile, we have $|I_{\alpha,\gamma}f(x)| \leq |I_\alpha f(x)|$, for every $f \in L^p_{loc}(\mathbb{R}^n)$. Using this inequality, $I_{\alpha,\gamma}$ is bounded on these spaces. In 1999, Kurata *et. al* [8] have proved that $W \cdot I_{\alpha,\gamma}$ is bounded on generalized Morrey spaces where W is a multiplication operator. Here, we shall discuss about the boundedness of $I_{\alpha,\gamma}$ on Lebesgue spaces and Morrey spaces. We shall see the influence of $K_{\alpha,\gamma}$ for the boundedness of $I_{\alpha,\gamma}$.

PRELIMINARY RESULTS

We can see that the Bessel-Riesz kernel vanishes faster at infinity than that the Riesz kernel. From this fact, we can show that the kernel of Bessel-Riesz is a member of some Lebesgue spaces. We begin with the following lemma.

Lemma 2 If $b > a > 0$ then $\sum_{k \in \mathbb{Z}} \frac{(u^k R)^a}{(1+u^k R)^b} < \infty$, for every $u > 1$ and $R > 0$.

Proof. Let $b > a > 0$, so that $b - a > 0$. We write $\sum_{k \in \mathbb{Z}} \frac{(u^k R)^a}{(1+u^k R)^b} = \sum_{k=-1}^{-\infty} \frac{(u^k R)^a}{(1+u^k R)^b} + \sum_{k=0}^{\infty} \frac{(u^k R)^a}{(1+u^k R)^b}$. Next, we estimate $\sum_{k=-1}^{-\infty} \frac{(u^k R)^a}{(1+u^k R)^b} \leq \sum_{k=-1}^{-\infty} (u^k R)^a < \infty$ and $\sum_{k=0}^{\infty} \frac{(u^k R)^a}{(1+u^k R)^b} \leq \sum_{k=0}^{\infty} (u^k R)^{a-b} < \infty$. Hence, we obtain $\sum_{k \in \mathbb{Z}} \frac{(u^k R)^a}{(1+u^k R)^b} < \infty$. ■

Lemma 2 is useful to prove the membership of $K_{\alpha,\gamma}$ in some Lebesgue spaces.

Theorem 3 If $0 < \alpha < n$ and $0 < \gamma$ then $K_{\alpha,\gamma} \in L^t(\mathbb{R}^n)$ and $\|K_{\alpha,\gamma}\|_{L^t} \sim \left(\sum_{k \in \mathbb{Z}} \frac{(2^k R)^{(\alpha-n)t+n}}{(1+2^k R)^{\gamma t}} \right)^{\frac{1}{t}}$, for $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$.

Proof. Suppose $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$ where $0 < \gamma$, $0 < \alpha < n$, so that $(\alpha - n)t + n > 0$. For arbitrary $R > 0$, write

$$\int_{\mathbb{R}^n} |K_{\alpha,\gamma}(y)|^t dy = \int_{\mathbb{R}^n} \frac{|y|^{(\alpha-n)t}}{(1+|y|)^{\gamma t}} dy = C_1 \int_{\mathbb{R}^+} \frac{r^{(\alpha-n)t+n-1}}{(1+r)^{\gamma t}} dr = C_1 \sum_{k \in \mathbb{Z}} \int_{2^k R \leq r < 2^{k+1} R} \frac{r^{(\alpha-n)t+n-1}}{(1+r)^{\gamma t}} dr, \quad C_1 > 0.$$

We obtain $\int_{\mathbb{R}^n} |K_{\alpha,\gamma}(y)|^t dy \leq C_1 \sum_{k \in \mathbb{Z}} \frac{1}{(1+2^k R)^{\gamma t}} \int_{2^k R \leq r < 2^{k+1} R} r^{(\alpha-n)t+n-1} dr = C_2 \sum_{k \in \mathbb{Z}} \frac{(2^k R)^{(\alpha-n)t+n}}{(1+2^k R)^{\gamma t}}$, $C_2 = \frac{C_1(2^{(\alpha-n)t+n}-1)}{(\alpha-n)t+n}$ and $\int_{\mathbb{R}^n} |K_{\alpha,\gamma}(y)|^t dy \geq \frac{C_1}{2^{\gamma t}} \sum_{k \in \mathbb{Z}} \frac{1}{(1+2^k R)^{\gamma t}} \int_{2^k R \leq r < 2^{k+1} R} r^{(\alpha-n)t+n-1} dr = C_3 \sum_{k \in \mathbb{Z}} \frac{(2^k R)^{(\alpha-n)t+n}}{(1+2^k R)^{\gamma t}}$, $C_3 = \frac{C_1(2^{(\alpha-n)t+n}-1)}{2^{\gamma t}[(\alpha-n)t+n]}$. Therefore $\int_{\mathbb{R}^n} |K_{\alpha,\gamma}(y)|^t dy \sim \sum_{k \in \mathbb{Z}} \frac{(2^k R)^{(\alpha-n)t+n}}{(1+2^k R)^{\gamma t}}$ for every $R > 0$. Using Lemma 2, take $t \in \left(\frac{n}{n+\gamma-\alpha}, \frac{n}{n-\alpha} \right)$, choose $u = 2$, and define $a := (\alpha - n)t + n$, $b := \gamma t$. We get $\sum_{k \in \mathbb{Z}} \frac{(2^k R)^{(\alpha-n)t+n}}{(1+2^k R)^{\gamma t}} < \infty$. Hence $K_{\alpha,\gamma} \in L^t(\mathbb{R}^n)$. ■

In this study, the membership of $K_{\alpha,\gamma}$ in Lebesgue spaces is an important result. With the result, we can use Young inequality [9] to prove the boundedness of $I_{\alpha,\gamma}$ on Lebesgue spaces.

Theorem 4 (Young's inequality) Let $1 \leq p, q, t \leq \infty$ satisfy $\frac{1}{q} + 1 = \frac{1}{p} + \frac{1}{t}$, then we have

$$\|g * f\|_{L^q} \leq \|g\|_{L^p} \|f\|_{L^t}$$

for every $g \in L^t(\mathbb{R}^n)$, $f \in L^p(\mathbb{R}^n)$.

Corollary 5 For $0 < \alpha < n$, $\gamma > 0$, we have

$$\|I_{\alpha,\gamma} f\|_{L^q} \leq \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^p}$$

for every $f \in L^p(\mathbb{R}^n)$ where $1 \leq p < t'$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, $\frac{1}{q} + 1 = \frac{1}{p} + \frac{1}{t}$.

By the above corollary, we can say that $I_{\alpha,\gamma}$ is bounded from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$. Moreover, norm of kernel Bessel-Riesz dominates norm of $I_{\alpha,\gamma} f$. Consequently in Lebesgue spaces, we obtain $\|I_{\alpha,\gamma}\| \leq \|K_{\alpha,\gamma}\|_{L^t}$.

We shall extend the boundedness of $I_{\alpha,\gamma}$ on Morrey spaces, but Young's inequality is not available on Morrey spaces. Using the Hardy-Littlewood maximal operator M , the boundedness of $I_{\alpha,\gamma}$ can be reproved on Lebesgue spaces and Morrey spaces. The operator M is defined by

$$Mf(x) := \sup_{x \in B} \frac{1}{|B|} \int_B |f(y)| dy, \quad x \in \mathbb{R}^n,$$

for every $f \in L^p_{loc}(\mathbb{R}^n)$ where $|B|$ denotes Lebesgue measure of ball $B = B(a, r)$ (centered at $a \in \mathbb{R}^n$ with radius $r > 0$). The supremum is taken over all open balls in \mathbb{R}^n . It is well known that the operator M is bounded on Lebesgue spaces $(L^p(\mathbb{R}^n), p > 1)$ [1, 10] and Morrey spaces [7].

MAIN RESULTS

In this section, we are going to discuss about the boundedness of the Bessel-Riesz operators on Morrey spaces. In the previous section, we have $K_{\alpha,\gamma} \in L'(\mathbb{R}^n)$ where $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$ and the inclusion property of Morrey spaces, so $K_{\alpha,\gamma} \in L^{s,t}(\mathbb{R}^n)$ where $1 \leq s \leq t$. Accordingly, we have the following theorem.

Theorem 6 Let $0 < \alpha < n$, $0 < \gamma$, then we have

$$\|I_{\alpha,\gamma} f\|_{L^{p_2,q_2}} \leq C_{p_1,q_1} \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}}$$

for every $f \in L^{p_1,q_1}(\mathbb{R}^n)$ where $1 < p_1 < q_1 < t'$, $1 \leq s \leq t$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, $\frac{1}{p_2} = \frac{1}{p_1} - \frac{q_1}{p_1 t'}$, and $\frac{1}{q_2} = \frac{1}{q_1} - \frac{1}{t'}$.

Proof. Suppose $0 < \alpha < n$, $0 < \gamma$ and take $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, $1 \leq s \leq t$. Let $f \in L^{p_1,q_1}(\mathbb{R}^n)$, $1 < p_1 < q_1 < t'$. For every $x \in \mathbb{R}^n$, write $I_{\alpha,\gamma} f(x) := I_1(x) + I_2(x)$ where $I_1(x) := \int_{|x-y|<R} \frac{|x-y|^{n-\alpha} f(y)}{(1+|x-y|)^\gamma} dy$ and $I_2(x) := \int_{|x-y|\geq R} \frac{|x-y|^{n-\alpha} f(y)}{(1+|x-y|)^\gamma} dy$, $R > 0$. To estimate I_1 and I_2 , we use dyadic decomposition. Now, estimate I_1

$$|I_1(x)| \leq C_1 \sum_{k=-1}^{-\infty} \frac{(2^k R)^{\alpha-n}}{(1+2^k R)^\gamma} \int_{2^k R \leq |x-y| < 2^{k+1} R} |f(y)| dy \leq C_2 Mf(x) \sum_{k=-1}^{-\infty} \frac{(2^k R)^{\alpha-n+n/s} (2^k R)^{n/s'}}{(1+2^k R)^\gamma}.$$

By using Hölder's inequality, we get

$$\begin{aligned} |I_1(x)| &\leq C_3 Mf(x) \left(\sum_{k=-1}^{-\infty} \frac{(2^k R)^{(\alpha-n)s+n}}{(1+2^k R)^{\gamma s}} \right)^{1/s} \left(\sum_{k=-1}^{-\infty} (2^k R)^n \right)^{1/s'} \\ &\leq C_4 Mf(x) \frac{\left(\int_{|x-y|<R} K_{\alpha,\gamma}^s(x-y) dy \right)^{1/s}}{R^{n(1/s-1/t)}} R^{n(1/s-1/t)'} \leq C_4 \|K_{\alpha,\gamma}\|_{L^{s,t}} Mf(x) R^{n/t'}. \end{aligned}$$

Hölder's inequality is used again to estimate I_2 :

$$|I_2(x)| \leq C_5 \sum_{k=0}^{\infty} \frac{(2^k R)^{\alpha-n}}{(1+2^k R)^\gamma} \int_{2^k R \leq |x| < 2^{k+1} R} |f(y)| dy \leq C_5 \sum_{k=0}^{\infty} \frac{(2^k R)^{\alpha-n}}{(1+2^k R)^\gamma} \left(\int_{2^k R \leq |x| < 2^{k+1} R} |f(y)|^{p_1} dy \right)^{1/p_1} (2^k R)^{n/p_1'}.$$

Next, we write

$$|I_2(x)| \leq C_6 \|f\|_{L^{p_1,q_1}} \sum_{k=0}^{\infty} \frac{(2^k R)^{\alpha-n+n/q_1}}{(1+2^k R)^\gamma} \frac{\left(\int_{2^k R \leq |x| < 2^{k+1} R} dy \right)^{1/s}}{(2^k R)^{n/s}} \leq C_6 \|f\|_{L^{p_1,q_1}} \sum_{k=0}^{\infty} \frac{\left(\int_{2^k R \leq |x| < 2^{k+1} R} \frac{|x-y|^{(\alpha-n)s}}{(1+|x-y|)^\gamma} dy \right)^{1/s}}{(2^k R)^{n/q_1-n} (2^k R)^{n/s}},$$

and we obtain $|I_2(x)| \leq C_6 \|f\|_{L^{p_1,q_1}} \|K_{\alpha,\gamma}\|_{L^{s,t}} \sum_{k=0}^{\infty} (2^k R)^{n/t'-n/q_1} \leq C_7 \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}} R^{n(1/t'-1/q_1)}$. Summing the two estimates, we get $|I_{\alpha,\gamma} f(x)| \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} (Mf(x) R^{n/t'} + \|f\|_{L^{p_1,q_1}} R^{n/t'-n/q_1})$, for each $x \in \mathbb{R}^n$.

Assume that f is not identically 0 and Mf is finite everywhere. Choose $R > 0$ such that $R^{n/q_1} = \frac{\|f\|_{L^{p_1,q_1}}}{Mf(x)}$. We get $|I_{\alpha,\gamma} f(x)| \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}}^{q_1/t'} Mf(x)^{1-q_1/t'}$. Define $\frac{1}{p_2} := \frac{(t'-q_1)}{p_1 t'}$ and $\frac{1}{q_2} := \frac{1}{q_1} - \frac{1}{t'}$. For arbitrary $r > 0$, we have

$$\left(\int_{|x|<r} |I_{\alpha,\gamma} f(x)|^{p_2} dx \right)^{1/p_2} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}}^{1-p_1/p_2} \left(\int_{|x|<r} |Mf(x)|^{p_1} dx \right)^{(1/p_2)}.$$

Divide by $r^{n/p_2-n/q_2}$ and take supremum to get

$$\begin{aligned} \|I_{\alpha,\gamma} f\|_{L^{p_2,q_2}} &= \sup_{r>0} \frac{\left(\int_{|x|<r} |I_{\alpha,\gamma} f(x)|^{p_2} dx \right)^{1/p_2}}{r^{n/p_2-n/q_2}} \\ &\leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}}^{1-p_1/p_2} \sup_{r>0} \frac{\left(\int_{|x|<r} |Mf(x)|^{p_1} dx \right)^{(1/p_2)}}{r^{n/p_2-n/q_2}} = C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}}^{1-p_1/p_2} \|Mf\|_{L^{p_1,q_1}}^{p_1/p_2}. \end{aligned}$$

Using the boundedness of M on Morrey spaces (Chiarenza-Frasca's Theorem [7]), we obtain an inequality $\|I_{\alpha,\gamma}f\|_{L^{p_2,q_2}} \leq C_{p_1,q_1} \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}}$. ■

By Theorem 6 and the inclusion property of Morrey spaces, for $1 \leq s \leq t$, we have

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,q_2}} \leq C_{p_1,q_1} \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,q_1}} \leq C_{p_1,q_1} \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,q_1}}$$

where $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$. We also obtain $\frac{p_2}{q_2} = \frac{p_1}{q_1}$. It is similar with Chiarenza-Frasca's result for the boundedness of fractional integral operators on Morrey spaces.

CONCLUDING REMARK

From the results of this study, we have seen that the norm of the Bessel-Riesz kernel dominates the norm of $I_{\alpha,\gamma}f$ for every f in Morrey space $L^{p,q}(\mathbb{R}^n)$ (p and q are suitable numbers). Moreover, using $K_{\alpha,\gamma} \in L^{s,t}(\mathbb{R}^n)$, $1 \leq s < t$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, the norm of the Bessel-Riesz kernel is closer to the norm of $I_{\alpha,\gamma}f$ than using $K_{\alpha,\gamma} \in L^t(\mathbb{R}^n)$. In the future, we shall continue this study to prove the boundedness of generalized Bessel-Riesz operators on Morrey spaces and generalized Morrey spaces.

ACKNOWLEDGMENTS

The first, second, and third authors are supported by ITB Research and Innovation Program 2015.

REFERENCES

1. E. M. Stein, *Singular Integral and Differentiability Properties of Functions* (Princeton University Press, Princeton, New Jersey, 1970).
2. G. H. Hardy and J. E. Littlewood, *Math. Zeit.* **27**, 565-606 (1927).
3. G. H. Hardy and J. E. Littlewood, *Math. Zeit.* **34**, 403-439 (1932).
4. S. L. Sobolev, *Amer. Math. Soc. Transl. Ser. 2* **34**, 39-68 (1963).
5. J. Peetre, *J. Funct. Anal.* **4**, 71-87 (1969).
6. D. R. Adams, *Duke Math. J.* **42**, 765-778 (1975).
7. F. Chiarenza and M. Frasca, *Rend. Mat.* **7**, 273-279 (1987).
8. K. Kurata, S. Nishigaki and S. Sugano, *Proc. Amer. Math. Soc.* **128**, 587-602 (1999).
9. L. Grafakos, *Classical Fourier Analysis* (Springer, New York, 2008).
10. E. M. Stein, *Harmonic Analysis: Real Variable Methods, Orthogonality, and Oscillatory Integrals* (Princeton University Press, Princeton, New Jersey, 1993).

The Boundedness of Bessel-Riesz Operators On Morrey Spaces

ORIGINALITY REPORT

16%

SIMILARITY INDEX

7%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Yuen Yiu. "New topological insulator thin film exhibits novel magnetic properties", Scilight, 2017
Publication 1%
- 2 "Function Spaces and Inequalities", Springer Nature, 2017
Publication 1%
- 3 A. H. Cahyana, A. Nadila, B. Ardiansah. "Unexpected synthesis of cinnamaldehyde-dimedone adduct using samarium (III) chloride", AIP Publishing, 2017
Publication 1%
- 4 Gunawan, Hendra, Denny I. Hakim, Kevin M. Limanta, and Al A. Masta. "Inclusion properties of generalized Morrey spaces : Inclusion properties of generalized Morrey spaces", Mathematische Nachrichten, 2016.
Publication 1%
- 5 Nguyen Minh Chuong. "Pseudodifferential 1%

Operators and Wavelets over Real and p-adic Fields", Springer Nature, 2018

Publication

6

Tanaka, Hitoshi, and Hendra Gunawan. "The Local Trace Inequality for Potential Type Integral Operators", Potential Analysis, 2013.

Publication

1%

7

www.m-hikari.com

Internet Source

1%

8

Yan Lin. "Strongly Singular Calderón–Zygmund Operator and Commutator on Morrey Type Spaces", Acta Mathematica Sinica English Series, 11/2007

Publication

1%

9

journals.smc.math.ca

Internet Source

1%

10

Guliyev, Vagif S., Fatih Deringoz, and Javanshir J. Hasanov. " (Φ, Ψ) -admissible potential operators and their commutators on vanishing Orlicz-Morrey spaces", Collectanea mathematica, 2015.

Publication

1%

11

Yoshihiro Mizuta. "Maximal functions, Riesz potentials and Sobolev embeddings on Musielak-Orlicz-Morrey spaces of variable exponent in \mathbb{R}^n ", Revista Matemática Complutense, 05/31/2011

1%

12

homepage.ruhr-uni-bochum.de

Internet Source

<1%

13

vdocuments.site

Internet Source

<1%

14

Der-Chen Chang. "Generalized Calderon-Zygmund operators on homogeneous groups and applications", *Applicable Analysis*, 05/2008

Publication

<1%

15

personal.fmipa.itb.ac.id

Internet Source

<1%

16

stat.epfl.ch

Internet Source

<1%

17

Yoshihiro Sawano. "Orlicz–Morrey Spaces and Fractional Operators", *Potential Analysis*, 06/15/2011

Publication

<1%

18

www.emis.de

Internet Source

<1%

19

pareto.uab.es

Internet Source

<1%

20

www.math.liu.se

Internet Source

<1%

21

home.hiroshima-u.ac.jp

Internet Source

<1%

22 Lu, Hsiao-feng, Amaro Barreal, David Karpuk, and Camilla Hollanti. "Efficiently sphere-decodable physical layer transmission schemes for wireless storage networks", EURASIP Journal on Advances in Signal Processing, 2016.
Publication <1%

23 pangu2002.org
Internet Source <1%

24 Vagif S. Guliyev, Fatih Deringoz. "Some Characterizations of Lipschitz Spaces via Commutators on Generalized Orlicz–Morrey Spaces", Mediterranean Journal of Mathematics, 2018
Publication <1%

25 Thomas Aronsson, Olof Johansson-Stenman, Tomas Sjögren. "Social Comparisons and Optimal Taxation in a Small Open Economy", The Scandinavian Journal of Economics, 2018
Publication <1%

26 S. V. Kislyakov. "On BMO-Regular Lattices of Measurable Functions. II", Journal of Mathematical Sciences, 09/2005
Publication <1%

27 Brian Kern. "Imaging through turbulence with a quadrature-phase optical interferometer", <1%

Applied Optics, 2005

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The Boundedness of Bessel-Riesz Operators On Morrey Spaces

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
