

Humoral and cellular immune response induced by antigenic protein of *Sarcoptes scabiei* var. *caprae*

by Poedji Hastutiek

Submission date: 05-Aug-2020 10:36AM (UTC+0800)

Submission ID: 1366058769

File name: 02_Humoral_and_celluler_immune....pdf (716.48K)

Word count: 4087

Character count: 21730

Humoral and cellular immune response induced by antigenic protein of *Sarcoptes scabiei* var. *caprae*

Nunuk Dyah Retno Lastuti¹, Wiwik Misaco Yuniarti², Poedji Hastutiek¹, Lucia Tri Suwanti¹ and Dony Chrismanto³

5
1. Department of Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia; 2. Department of Veterinary Clinic, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia; 3. Study Programme of Animal Health, Faculty of Vocation, Universitas Airlangga, Surabaya, Indonesia.

Corresponding author: Wiwik Misaco Yuniarti, e-mail: wiwikmisaco@yahoo.com

Co-authors: NDR: nunuk_dyah@fkh.unair.ac.id, PH: poedjihastutiek@gmail.com, LTS: tswant@gmail.com, DC: donychrismanto@gmail.com

Received: 17-01-2018, Accepted: 14-05-2018, Published online: 19-06-2018

doi: 10.14202/vetworld.2018.819-823 How to cite this article: Lastuti NDR, Yuniarti WM, Hastutiek P, Suwanti LT, Chrismanto D (2018) Humoral and cellular immune response induced by antigenic protein of *Sarcoptes scabiei* var. *caprae*, *Veterinary World*, 11(6): 819-823.

Abstract

Aim: Scabies is one of the most important diseases in goats and caused by a complex hypersensitivity process that involves both humoral and cell-mediated immune responses. This phenomenon shows that the variety of *Sarcoptes scabiei* has different characteristics of specific antigenic protein or different immune-dominant. This research aims to detect the humoral and cellular immune response of rabbits which were immunized with the protein of *S. scabiei* var. *caprae*.

Materials and Methods: This research was done as follows, identification and collection of *Sarcoptes scabiei* var. *caprae* from goat infected with scabies, separation of protein antigen from *S. scabiei* mites with ultrasonic sonicator, measurement of protein content with spectrophotometry, rabbit injection with 500 µg dose of antigen protein which was repeated 5 times (5x booster) every 2 weeks. Measurement of IgG titer using indirect ELISA, whereas to detect the expression of cellular immune response (TLR-9, CD4, and CD8) using Direct Immunofluorescence assay.

Results: Based on the statistical analysis, it showed that there was a significant enhancement (p<0.05) for optical density value or antibody titer and cellular immune response was shown by TLR-9, CD4, and CD8 expression in rabbit T lymphocytes which appear yellow to green fluorescent color using fluorescence microscope. The amount of fluorescence T lymphocytes showed a significant difference (p<0.05) between the control and various boosters.

Conclusion: Antigenic protein of *S. scabiei* var. *caprae* contains ligands, which are involved in the pathogen-associated molecular pattern that has an ability to induce humoral and cellular immune response in rabbit. Specifically, that TLR-9 is not only involved in innate immunity but also in adaptive immunity and can be used as alternative adjuvant development research.

Keywords: antigenic protein, CD4, CD8, immunoglobulin G, *Sarcoptes scabiei* var. *caprae*, toll-like receptor-9.

Introduction

17
Sarcoptic mange is one of the most economically important diseases in goats in Indonesia, but the control of this ectoparasite is limited [1]. An increasing number of treatment failures is being reported because of the drug resistance. Vaccination is an attractive ecological alternative to the use of acaricides for the parasite control. However, effective antiparasite vaccines against sarcoptic mange have not yet been developed [2]. 12

At present, it is considered as an emerging/re-emerging parasitic disease that threatens human and animal healthy globally [3]. Since the prevalence of the disease in human and animals is very high, the

1
economic losses caused by the disease are enormous. The development of a specific vaccine would be a sustainable option for the control of this disease [4].

Vaccine development requires preliminary research through immunogenic protein exploration of *Sarcoptes scabiei* which is isolated from goats. Previous studies have shown that *S. scabiei* var. *caprae* contains antigenic proteins with molecule weight 205.8 kDa, 57.3 kDa, 43.0 kDa; these are proteins have antigenic characteristics which are probably involved in scabies pathogenesis in goats [5].

Immunogenic protein can work maximally as a vaccine when it can stimulate T cells activation together with co-receptor CD4 and CD8, which plays a role in the immune system activation. CD4+ T cells will help B cells to produce antibodies and also it helps phagocytosis to destroy ingested microbes, while CD8+T cells will kill intracellular microbes [6-9]. It has been known that toll-like receptor (TLR) in lymphocytes act as an essential signal to regulate lymphocytes activation and proliferation, produce antibodies, and regulate antigen presentation. TLR stimulation through microbial product initiates signaling pathways

1
Copyright: Lastuti, et al. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

which activate not only the innate immunity but also the adaptive immunity [6,10-12]. To investigate that TLR plays a role in adaptive immunity, it requires research toward *S. scabiei* protein immunized in rabbits to detect cellular immune response and TLR-9 as markers. TLR-9 has been developed by some researchers as vaccine adjuvant containing synthetic oligodeoxynucleotide to fight infectious diseases, allergies, and cancers [13-16].

The aim of the study was to detect humoral and cellular immune response of rabbits which were immunized with *S. scabiei* var. *caprae* protein. Increasing optical density (OD) value or antibody titer (immunoglobulin G [IgG]) as a parameter of humoral immune response and TLR-9, CD4, and CD8 as a parameter of cellular immune response that could be used preliminary study for the development of subunit vaccine as an alternative for scabies prevention on goats in Indonesia.

2 Materials and Methods

Ethical approval

The entire research was conducted appropriately following the ethics in using experimental animals and has been approved by the Ethics Commission of the Faculty of Veterinary Medicine, Universitas Airlangga, No: 630-KE.

Isolation of *S. scabiei* mites and rabbits immunization

S. scabiei mites (adults, nymphs, and larvae) were isolated from domestic goats that showed clinical signs of scabies such as thickening of the skin, crust formation, and alopecia on the area around eyes, ears, mouth, and legs [5,6,17]. After the identification based on key identification from Soulsby [17], mites isolation was required for rabbits immunization by these following steps: Mites (around one thousand mites) were put into a Petri dish and mixed with phosphate-buffered saline (PBS) solution and strained to get the result free of crust. The result was washed with PBS solution and centrifuged at 3000 rpm for 10 min. The washing process was performed at least 3 times, to get mites free of dirty materials that were carried from the scraping process.

The deposited mites would be formed as pellets and kept in a freezer at -80°C , to be processed into homogenized protein [5]. The pellets (mites) with homogenizing medium: 100 mM Tris-HCL buffer pH 7.4, 100 mM NaCl, 5 mM EDTA, 5 mM MgCl_2 [18] were sonicated at 30 K Hz and this step was repeated for 16 times. Every sonication step lasts for 4 min with a break time of 2 min [5]. The sonication result solution was centrifuged at 16,000 rpm for 5 min; then, the protein concentration of the resulting supernatant was measured during visible spectrophotometer with 595 nm wavelength.

This research used six rabbits, 4 months old and prepared for the immunization trial at the laboratory of experimental animals, Faculty of Veterinary Medicine, Universitas Airlangga. Each of experimental rabbits was injected with 500 μg *S. scabiei* protein (0.3 ml) and added Freund adjuvant complete (Sigma, USA) with

ratio 1:1. Every 2 weeks, 5 times the injection was performed with the same protein as a booster. Before the first injection, about 10 ml of rabbit blood was taken for TLR-9, CD4, and CD8 examination as preliminary data (control) and whole blood examination was conducted at the end of first booster until the fifth booster.

Examination of humoral immune response (IgG) with direct ELISA

To assess the humoral immune response (IgG), blood samples were collected from the marginal ear vein before immunization (control), on the 14th day (1st booster), the 28th day (2nd booster), the 42th day (3rd booster), the 56th day (4th booster), and the 70th day (5th booster). Serum samples were obtained from blood and stored at -20°C until use.

High-binding microtiter plates (BIO-RAD, USA) were coated overnight at 4°C with 100 μl of antigen solution of *S. scabiei* (10 $\mu\text{l}/\text{ml}$ antigen in PBS) in each, then washed with 200 μl PBS-T20 for 3 times. After that, blocking with 4% skim milk (Maxcream, Indonesia) PBS-Tween as much as 200 $\mu\text{l}/\text{well}$ and incubated at 37°C for an hour, then washed with PBS-T20 for 3 times. Rabbits' serum from immunization result was added and tested at 1/200 dilution as much as 100 μl for each well. This stage was performed in twice.

For antibody measurement standard, antibody dilution was done that the negative control antibody titer had been known at dilution of 1/200 up to 1/204.800 and added 100 μl for each well and then incubated at 37°C for an hour and washed. The next step was the addition of conjugate IgG anti-rabbit (Promega Corporation, USA) with 1/4000 dilution as much as 100 μl per well and incubated at 37°C for one hour. Then, the plate was washed and added substrate pNPP into substrate buffer (diethanolamine 1 mg/ml) as much as 100 μl for each well. Incubation was done in a dark room at room temperature for 15-45 min. The reaction was stopped by adding 3N NaOH solution for 50 μl in each well, and the plate was read using ELISA reader (BIO-RAD, USA) with wavelength of 405 nm.

The ELISA reading was done as these following steps: OD value standard (positive or negative) was sorted by the maximum to the minimum OD value. As the negative control limit, it was taken OD value at dilution 1/200. The highest dilution of positive control antibody that is still showing positive value based on cutoff value (2 times the average negative control) is an antibody titer. Then, it was made a statistic range from antibody standard, for example, the maximum OD range >3.171 has antibody titer 204.800, between 2.999 and 3.171 has antibody titer 102.400, and so on up to the zero titer. For measuring sample OD value, put sample OD value on the titer range below. Antibody titer showed a negative result if the ELISA reading result between control rabbits and immunized rabbits obtained a similar OD value. Antibody titer showed a positive result if OD value from immunized rabbits was higher or at least 2 times larger than OD value from control rabbits [19].

Immunofluorescence technique for TLR-9, CD4, and CD8 examination

For the examination of TLR-9, CD4, and CD8, the isolation of PBMC in T cells was performed based on Boyum method (1968) with some modifications [19], as these following procedures: 5 ml whole blood and washed with 10% PBS, centrifuged at 1600 rpm, and temperature 10°C for 10 min, next the undercoat to be put on Ficoll isopaque. The mixture containing blood and Ficoll isopaque was centrifuged at 1600 rpm and temperature 10°C for 10 min. The resultant buffy coat was separated and washed with Tween PBS. TLR-9 examination was performed as these following stages, buffy coat, and 300 µl MEM (Sigma, USA) incubated at 37°C for an hour, the solution was fixed by absolute methanol, and blocked by PBS and 1% fetal calf serum for 15 min, next the solution was washed using Tween PBS and there was added the first antibody of monoclonal antibody TLR9-fluorescein isothiocyanate (FITC) labeled (IMGENEX Corp). The solution added 1% fetal calf serum and FITC conjugate. The result was examined by fluorescence microscope using 400x magnification, to find out whether any yellow-to-green fluorescence color from T cells expressing TLR-9. If there is fluorescence light from T cells showing an activated immune response, then the calculation is conducted toward the number of fluorescent T cells on 20 µl solution.

Examination of CD4 and CD8 was performed as these following: buffy coat was centrifuged at 1700 rpm for 5 min, the supernatant was removed, 30 µl 70% methanol was added to the pellet and make it into suspension. Next, the suspension was incubated for 5 min at room temperature and centrifuged at 10,000 rpm for 5 min. The supernatant was removed, washed with Tween PBS once, and centrifuged at 10,000 rpm for 5 min, then added 200 µl PBS and make it into suspension. 10 µl of suspension was taken, and put into a tube, added 10 µl Monoclonal Mouse Anti-Human CD4, and Clone MT310 (DakoCytomation, Denmark), Monoclonal Mouse Anti-Human CD8, Clone DK25 (DakoCytomation, Denmark). Make the suspension into suspension and incubated in dark room at 4°C for 30 min or 37°C for 30 min. A solution about 20 µl dropped on glass object, examined under a fluorescence microscope with 400x magnification, to detect the existence of fluorescence color of greenish yellow from T cell which expressed both CD4 and CD8. If there was fluorescence color of greenish yellow which showed activated immune response, then the counting toward the number of fluorescent T cells would be done.

Results

The humoral immune response was shown by OD value enhancement on 3rd booster that reached 16.3 times (mean 3.0424 ± 0.0313) compared to OD control value (mean 0.1838 ± 0.0552). Based on the statistical analysis (multivariate tests), it showed that there was a significant enhancement ($p < 0.05$) for both

values from OD value or antibody titer. From the data obtained, the increasing OD value started from day 14 and reached the peak on day 28 (2nd booster), the enhancement stays up to day 42 (3rd booster), and constantly increases up to day 70 or 5th booster (3.1226 ± 0.0430). As well as antibody titer measurement was performed by series of serum dilution then based on cutoff value, it can be known at which dilution antibodies can still form a specific reaction. According to the statistical analysis, it showed that between the control and treatment (booster) group, there is a significant increase ($p < 0.05$) (Figure-1).

The cellular immune response was shown by the expression of TLR-9, CD4, and CD8 in rabbit T cells which is marked by the yellow-to-green fluorescence color after rabbit immunization up to 5 times booster (Figure-2). In addition, the amount of fluorescence T cells was counted for each 20 µl buffy coat; it was shown the amount of fluorescence in T cells is increased in accordance with the treatment from various boosters (booster one up to booster five).




Figure-1: Level of optical density value and antibody titer (immunoglobulin G) of rabbit resulted from immunization with soluble mite protein.




Figure-2: Toll like receptor-9 (a), CD4 (b), CD8 (c) expression in Rabbit T cells, visualized by fluorescein isothiocyanate (400× magnification).

T_{H1} amount of T cells which expresses TLR-9 shows a significant difference ($p < 0.05$) between control and various booster (repetitions), but between booster 3 (day 42) and booster 4 (day 56), there is no significant difference by statistic despite the increasing amount of fluorescence T cells, while the amount of T cells which expresses CD4 shows a significant difference ($p < 0.05$) between control and various booster (repetitions), as well as the amount of T cells which expresses CD8 shows a significant difference ($p < 0.05$) between control and various booster as mentioned in Figure-3.

Discussion

Antibody titer (IgG) enhancement is caused by the memory cells' activation which happened on the 30th day after initial immunization, and adaptive immune system is like nerve system, capable of memorizing, particularly in experiencing antigen exposure. The immunization result indicated that *S. scabiei* var. *caprae* protein has an ability to induce humoral immune response to produce antibodies (IgG) in immunized rabbits. In accordance with the principles of immunization method, which is increasing the degree of immunity, providing protective immunity by inducing memory response toward specific pathogens with non-virulent or non-toxic antigen [20-23]. In the initial immunization process, there was a recognition by naive T cells and naive B cells; then there was an active phase which was expansive clonal from T cells and differentiated B cells that occurred on day 7 post-immunization [23]. The next process was an antigen elimination or also known as effector phase, this process occurred on day 14, after that apoptosis process occurred and this was followed by the memory cells' activation in day 30 [20,22]. Adaptive immune

system is like the nervous system, capable of memorizing particularly in experiencing antigen exposure; this is the reason why immunity is developed through immunization like the phenomena showed in experimental animal. When the experimental animal immunized by one of antigen, immune response (antibody) or cell-mediated will be appeared after several days, increased rapidly and exponentially (can reach the peak on day 50), then decreased gradually, this is the characteristic of primer immune response [24].

S. scabiei var. *caprae* protein has a signal character mediated by TLR9, as well as an immune response induced through CD4 activation related to humoral and CD8 immune responses related to cellular immune responses. This finding is an immune cell communication that plays a role in the dependent pathogenesis of the type of protein expressed in each animal. Based on the exploratory results of the *S. scabiei* var. *caprae* mite protein, it has been demonstrated that the mite contains a protein capable of inducing humoral and cellular immune responses that can be used further research for the development of subunit vaccines for goat scabies.

Conclusion

Antigenic protein of *S. scabiei* var. *caprae* has the ability to induce humoral and cellular immune response through the enhancement of IgG titer, expression of TLR-9, CD4, and CD8 in rabbits' T cells. TLR-9 signal is capable not only plays a role in innate immunity but also in adaptive immunity. Antigenic protein of *S. scabiei* var. *caprae* may contain a ligand which acts as a receptor that is involved in the pathogen-associated molecular pattern, and as preliminary research to explore the immunogenic protein which plays a role in the immune system activation




Figure-3: The amount T cells of rabbits which express toll-like receptor9 (a), CD4 (b), CD8 (c), resulted from immunization with soluble mite protein.

for the future studies in vaccine development to overcome scabies on goats.

Authors' Contributions

NDRL, WMY, LTS, PH, and DC were carried out the main research works, PH and DC performed the statistical analysis of the main data in the experiments, and all of the authors were approved the final manuscript. All authors read and approved the final manuscript.

Acknowledgments

We would like to thank the Ministry of Research, Technology, and Higher Education, Indonesia. This study was supported by research Grant, No. 004/SP2H/LT/DRPM/IV/2017. We also thank the Rector of Universitas Airlangga and Director of Research and Innovation Department, Universitas Airlangga.

Competing Interests

The authors declare that they have no competing interests.

References

1. Tarigan, S. (2004) Antibody responses in naïve and sensitised goats infested by *Sarcoptes scabiei*. *Ind. J. Anim. Vet. Sci.*, 9(4): 258-265.
2. Casais, R., Grandá, V., Balseiro, A., Cerro, A.D., Dalt, N., González, E., Bra, P., Prieto, J.M. and Montoya, M. (2016) Vaccination of rabbits with immunodominant antigens from *Sarcoptes scabiei* induced high levels of humoral responses and pro-inflammatory cytokines but confers limited protection. *Parasit. Vectors*, 9: 435-447.
3. Alasaad, S., Rossi, L., Heukelbach, J., Perez, J.M., Hamarshah, O., Otiende, M. and Zhu, X.Q. (2013) The neglected navigating web of the incomprehensibly emerging and re-emerging sarcoptic mite. *Infect. Infect Genet. Evol.*, 17: 253-259.
4. Gu, X., Xie, Y., Wang, S., Peng, X., Lai, S. and Yang, G. (2014) Immune response induced by candidate *Sarcoptes scabiei* var. *cuniculi* DNA vaccine encoding paramyosin in mice. *Exp. Appl. Acarol.*, 63: 401-412.
5. Lastuti, N.D.R. (2009) Specific antigenic protein 57.3 kDa of *Sarcoptes scabiei* var. *Caprae* as Material Candidate of Scabies Diagnostic Kit for Goat and Toll-like Receptor-Mediated Immune Responses. Doctoral Dissertation. Postgraduate Program. Universitas Airlangga.
6. Lastuti, N.D.R., Rantam, F.A., Hastutiek, P. and Chrismanto, D. (2017) Toll-like receptor (TLRs) play a role in adaptive immunity in rabbits immunized by *Sarcoptes scabiei* proteins. *VMIC Conf. Proc. KnE Life Sci.*, 1: 1-9.
7. Kammanadiminti, S.J., Mann, B.J., Dutil, L. and Chadee, K. (2003) Regulation of toll-like receptor 2 expression by gal-lectin of *Entamoeba histolytica*. *Faseb. J.*, 18: 155-157.
8. Arlian, L.G., Morgan, M.S. and Neal, J.S. (2003) Modulation of cytokine expression in human keratinocytes and fibroblasts by extracts of scabies mites. *Am. J. Trop. Med. Hyg.*, 69: 652-656.

9. Clement, M., Ladell, K., Makinde, J.E., Miles, J.J., Edwards, E.S., Dolton, G., Williams, T., Schauenburg, A.J.A., Cole, D.K., Lauder, S.N., Gallimore, A.M., Godkin, A.J., Burrows, S.R., Price, D.A., Sewell, A.K. and Wooldridge, L. (2011) Anti-CD8 antibodies Can trigger CD8+T cell effector function in the absence of TCR engagement and improve peptide MHC-I tetramer staining. *J. Immunol.*, 187: 654-663.
10. Pasare, C. and Medzhitov, R. (2005) Control of B cell responses by toll-like receptor. *Nature*, 438(7066): 364-368.
11. Blander, J.M. and Medzhitov, R. (2006) Toll-dependent selection of microbial antigens for presentation by dendritic cells. *Nature*, 440(7085): 808-812.
12. Xiao, H., Li, X. and Abbot, D.W. (2008) In: Gregory, W.K., editor. Analysis of TLR Expression, Regulation and Signalling. Chapter 3: Signalling by Toll-Like Receptor. CRC Press, Taylor & Francis Group, Boca. p39-55.
13. McCluskie, M.J. and Krieg, A.M. (2006) Enhancement of infectious disease vaccines through TLR9 dependent recognition of CpG DNA. *Curr. Top. Microbiol.*, 311: 155-178.
14. Schetter, C. and Vollmer, J. (2004) Toll-like receptors involved in response to microbial pathogens: Development of agonists for toll-like receptor 9. *Curr. Opin. Drug. Discov. Dev.*, 7: 204-210.
15. Bhardwaj, N., Gnjatic, S.S. and Sawhney, N.B. (2010) TLR agonists: Are they good adjuvants? *Cancer J.*, 16(4): 382-391.
16. Toussi, D.N. and Massari, P.P. (2014) Review: Immune adjuvant effect of molecularly-defined toll-like receptor ligands. *Vaccines*, 2: 323-353.
17. Soulsby, E.J.L. (1986) Helminths, Arthropods and Protozoa of Domesticated Animal. 7th ed. The English and Protozoa of Society and Baillire, Tindall, London. p504-506.
18. Reed, R., Holmes, D., Weyers, J. and Jones, A. (2003) Practical Skills in Biomolecular Sciences. 2nd Ed. Pearson, Prentice Hall. Ashford Colour Press Ltd., Gosport. p237-319.
19. Rantam, F.A. (2003) The Method of Immunology. Airlangga University Press, Surabaya. p145-155.
20. Abbas, A.K. and Lichtman, A.H. (2005) Cellular and Molecular Immunology. 5th ed. International Edition. Elsevier Saunders Inc., Philadelphia, PA, Pennsylvania. p41-105, 411-432.
21. Bomstein, S., Frossling, J.K., Zakrisson, G. and Morner, T. (2006) Evaluation of a serological test (indirect ELISA) for the diagnosis of sarcoptic mange in red foxes. *Vet. Dermatol.*, 17(6): 411-416.
22. Rodriguez-Cadenas, F., Carbajal-Gonzales, M.T., Fregeneda-Grandes, J.M., Aller-Gancedo, J.M. and Rojo-Vazquez, F.A. (2010) Clinical evaluation and antibody responses in sheep after primary and secondary experimental challenges with the mange mites *Sarcoptes scabiei* var. *ovis*. *Vet. Immunol. Immunopathol.*, 133: 109-116.
23. Tarigan, S. and Huntley, J.E. (2005) Failure to protect goats following vaccination with soluble proteins of *Sarcoptes scabiei*: Evidence for a role for IgE antibody in protection. *Vet. Parasitol.*, 133(1): 101-109.
24. Casais, R., Millan, J., Rosell, J.M., Dalton, K.P. and Prieto, J.M. (2015) Evaluation of an ELISA using recombinant Ssλ20ΔB3 antigen for the serological diagnosis of *Sarcoptes scabiei* infestation in domestic and wild rabbits. *Vet. Parasitol.*, 214(3-4): 315-321.

Humoral and cellular immune response induced by antigenic protein of *Sarcoptes scabiei* var. *caprae*

ORIGINALITY REPORT

18%

SIMILARITY INDEX

16%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	link.springer.com Internet Source	3%
2	www.e-sciencecentral.org Internet Source	2%
3	parasitesandvectors.biomedcentral.com Internet Source	2%
4	mafiadoc.com Internet Source	1%
5	W.P. Lokapirnasari, A. R. Dewi, A. Fathinah, S. Hidanah et al. " Effect of Probiotic Containing Supplementation on The Laying Performance and Egg Cholesterol of Quail ", IOP Conference Series: Earth and Environmental Science, 2018 Publication	1%
6	krishikosh.egranth.ac.in Internet Source	1%
7	www.scribd.com Internet Source	1%

8	download.atlantis-press.com Internet Source	1%
9	theses.gla.ac.uk Internet Source	1%
10	ijpa.tums.ac.ir Internet Source	1%
11	eprints.brighton.ac.uk Internet Source	<1%
12	Singh, Shanker K., Umesh Dimri, Bhaskar Sharma, Meeta Saxena, and Priyambada Kumari. "Assessment of the cytokine profile in peripheral blood mononuclear cells of naturally <i>Sarcoptes scabiei</i> var. <i>canis</i> infested dogs", <i>Veterinary Parasitology</i> , 2014. Publication	<1%
13	Jonathan Fior. "An Initial In Vitro Investigation into the Potential Therapeutic Use of SupT1 Cells to Prevent AIDS in HIV-Seropositive Individuals", <i>PLoS ONE</i> , 2012 Publication	<1%
14	mcb.asm.org Internet Source	<1%
15	www.biomedcentral.com Internet Source	<1%
16	hdl.handle.net	

<1%

17

Tarigan, S.. "Failure to protect goats following vaccination with soluble proteins of *Sarcoptes scabiei*: Evidence for a role for IgE antibody in protection", *Veterinary Parasitology*, 20051010

Publication

<1%

18

www.mdpi.com

Internet Source

<1%

19

digital.library.unt.edu

Internet Source

<1%

20

www.oncotarget.com

Internet Source

<1%

21

Jihyun Yang, Sang-Mu Shim, Thi Quyen Nguyen, Eun-Ha Kim, Kwang Kim, Yong Taik Lim, Moon-Hee Sung, Richard Webby, Haryoung Poo. "Poly- γ -glutamic acid/chitosan nanogel greatly enhances the efficacy and heterosubtypic cross-reactivity of H1N1 pandemic influenza vaccine", *Scientific Reports*, 2017

Publication

<1%

22

Y. Mulugeta, Hailu T. Yacob, Hagos Ashenafi. "Ectoparasites of small ruminants in three selected agro-ecological sites of Tigray Region, Ethiopia", *Tropical Animal Health and*

<1%

23

Zhang, Runhui, Quwu Jise, Wanpeng Zheng, Yongjun Ren, Xiang Nong, Xuhang Wu, Xiaobin Gu, Shuxian Wang, Xuerong Peng, Songjia Lai, and Guangyou Yang. "Characterization and evaluation of a *Sarcoptes scabiei* allergen as a candidate vaccine", *Parasites & Vectors*, 2012.

Publication

<1%

24

M. Nova Raditya, A.M. Made Bagus, Ulayatul Kustiati, Hevi Wihadmadyatami, Dwi Liliek Kusindarta. "Data of The Expression of Serotonin in Alzheimer's Disease (AD) Rat Model Under Treatment of Ethanolic Extract *Ocimum sanctum* Linn", *Data in Brief*, 2020

Publication

<1%

25

Lukiswanto, Bambang, Muhammad Arifin, and Ira Yudaniayanti. "Expression of TGF-beta1 during fracture repair in Ovariectomized Rats with therapy of *Cissus quadrangularis* extract", *Journal of Physiology and Pharmacology Advances*, 2015.

Publication

<1%

26

Norihisa Noguchi, Mitsuhiro Hase, Masami Kitta, Masanori Sasatsu, Koichi Deguchi, Megumi Kono. " Antiseptic susceptibility and distribution of antiseptic-resistance genes in methicillin-

<1%

27

Subhashisa Swain, Minakshi Bhatt, Sanghamitra Pati, Ricardo J. Soares Magalhaes. "Distribution of and associated factors for dengue burden in the state of Odisha, India during 2010–2016", *Infectious Diseases of Poverty*, 2019

Publication

<1%

28

Xiaobin Gu, Yue Xie, Shuxian Wang, Xuerong Peng, Songjia Lai, Guangyou Yang. "Immune response induced by candidate *Sarcoptes scabiei* var. *cuniculi* DNA vaccine encoding paramyosin in mice", *Experimental and Applied Acarology*, 2014

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Humoral and cellular immune response induced by antigenic protein of *Sarcoptes scabiei* var. *caprae*

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
