

Volume - 12

No. 2

February-2019

ISSN: 0972-8988

EISSN: 2231-0916



NAAS Rating : 5.71

Indexed in ESCI-Thomson Reuters, PubMed,
PubMed Central, DOAJ, Scopus, CABI, CAS etc.

Veterinary World

Open access and peer reviewed journal



Editorial office

Veterinary World,
Star, Gulshan Park,
NH-8A, Chandrapur Road,
Wankaner - 363621,
Dist. Morbi, Gujarat, India
Website: www.veterinaryworld.org
E-mail: editorveterinaryworld@gmail.com

Veterinary World

Editorial Office: Veterinary World, Star, Gulshan Park, NH-8A, Chandrapur Road, Wankaner - 363621, Dist. Morbi, Gujarat, India

ISSN: 0972-8988, EISSN: 2231-0916, www.veterinaryworld.org

Editor-in-Chief

Anjum V. Sherasiya - Ex-Veterinary Officer, Department of Animal Husbandry, Gujarat State, India

Associate Editors

Shambhunath Choudhary - Department of Biomedical & Diagnostic Sciences, College of Veterinary Medicine, The University of Tennessee, 2407 River Drive, Room A 201, Knoxville, TN 37996, U.S.A.

Suresh H. Basagoudanavar - FMD Vaccine Research Laboratory, IVRI, Bangalore, Bangalore- 560024, Karnataka, India

Editorial board

R. G. Jani - Ex-Coordinator Wildlife Health, Western Region Centre, Indo-US Project, Department of Veterinary Medicine, Veterinary College, Anand - 388001, Gujarat, India

G. N. Gongal - Technical Officer, WHO South-East Asia Regional Office, New Delhi -110002, India

Ranganath Mamidi - Dr. Julian E Stelzer's Lab, Department of Physiology & Biophysics, Medical School, Case Western Reserve University, Cleveland, OH - 44106, U.S.A.

Md. Tanvir Rahman - Department of Microbiology and Hygiene, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

Deepmala Agarwal - Cancer Prevention Laboratory, Pennington Biomedical Research Center, Baton Rouge, LA, U.S.A.

Foud Kasim Mohammad - Professor, Department of Pharmacology & Toxicology, Vice President for Administrative & Financial Affairs, University of Mosul, P.O. Box 11136, Mosul, Iraq

Abdel-Baset Nasr Sayed Ahmed - Professor and Head, Department of Animal Nutrition and Clinical Nutrition, Faculty of Veterinary Medicine, Assiut University, Assiut, Egypt

Nicole Borel - Department of Pathology, Vetsuisse Faculty, University of Zurich, CH-8057 Zürich, Switzerland

B. A. Lubisi - Virology, MED Programme, ARC - Onderstepoort Veterinary Institute, No. 100 Old Soutpan Road, Onderstepoort, Tshwane, 0110, South Africa

Kumar Venkitanarayan - Associate Professor, Graduate Programs Chair, Honors and Pre-Vet Programs Advisor, Department of Animal Science, University of Connecticut, Storrs, CT 06269, U.S.A.

Kemin Xu - Department of Veterinary Medicine, University of Maryland, College Park College Park, MD, 20742, U.S.A.

Vassilis Papatziros - Faculty of Veterinary Medicine, Department of Medicine (Porcine Medicine), University of Thessaly, Thessaly, Greece

Mathias Devreese - Laboratory of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Ghent University, Belgium

Sumeet Sharma - Edmonton North Animal Hospital, Edmonton, Alberta, T5X 3Y7, Canada

K. P. Singh - School of Medicine and Dentistry, University of Rochester, Department of Environmental Medicine, Room: 4- 6820, 601 Elmwood Avenue, Box-EHSC, Rochester, New York-14620, U.S.A.

Raj Mohan Raja Muthiah - Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, U.S.A.

Ashok K. Chockalingam - Division of Applied Regulatory Science, U.S. Food and Drug Administration, 10903, New Hampshire Avenue, Silver Spring, Maryland 20993, U.S.A.

Ashutosh Wadhwa - Poxvirus and Rabies Branch, Division of High-Consequence Pathogens and Pathology, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention, 1600 Clifton Road, NE, Mail Stop G33, Atlanta, GA 30 333, U.S.A.

Luiz Otavio Pereira Carvalho - Laboratory of Immunomodulation and Protozoology, Oswaldo Cruz Institute, Ministry of health (Brazil), Pavilhão "108" - Sala: 09, Av. Brasil, 4365 - Manguinhos, Rio de Janeiro - RJ, CEP: 21040-360, Brazil.

Mallikarjun Bidarimath - Cornell Stem Cell Program, Department of Biomedical Sciences, T2-012 Veterinary Research Tower, Cornell University, College of Veterinary Medicine, Ithaca, NY 14853-6401, USA.

Ratings of Veterinary World: NAAS - National Academy of Agricultural Sciences - 5.71, Scimago Journal Rank -0.284, Citescore - 0.57, SNIP - Source Normalized Impact per Paper - 0.570

Indexing and abstracting

Academic Journals Database, AGORA, AGRICOLA, AGRIS, CABI, CAS, DOAJ, EBSCO, EMBASE, ESCI - Thomson Reuters, Gale, Google Scholar, HINARI, Indian Animal Science Abstracts, Indian Science Abstracts, JournalSeek, Open J-gate, ProQuest, PubMed, PubMed Central, SCOPUS, TEEAL

Publisher: Veterinary World

Veterinary World is an open access journal, each issue available free of cost at www.veterinaryworld.org.

We accept online submission only. For more information regarding submission and publication charges, please visit www.veterinaryworld.org

Rs. 8000 for Indian/USD 250 for Abroad per copy

Printed and Published by Dr. Anjum V. Sherasiya on behalf of Veterinary World. Printed and Published at Star, Gulshan Park, N.H. 8A, Chandrapur Road, Wankaner-363621, Dist. Morbi, Gujarat, India.
Editor: Dr. Anjum V. Sherasiya

Veterinary World

ISSN: 0972-8988, EISSN: 2231-0916, www.veterinaryworld.org

Volume-12

No.2

February-2019

The articles in Veterinary World are open access articles licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Research (Published online: 05-02-2019)

1. Foot-and-mouth disease outbreaks in Egypt during 2013-2014: Molecular characterization of serotypes A, O, and SAT2
Emad Diab, Abdel-Hamid I. Bazid, Mohamed Fawzy, Wagdy R. El-Ashmawy, Adel A. Fayed and Magdy M. El-Sayed
Veterinary World, 12(2): 190-197

Research (Published online: 08-02-2019)

2. Prevalence and diversity of gastrointestinal protozoa in Madura cattle at Bangkalan Regency, East Java, Indonesia
Poedji Hastutiek, Wiwik Misaco Yuniarti, Mufasirin Djaeri, Nunuk Dyah Retno Lastuti, Endang Suprihati and Lucia Tri Suwanti
Veterinary World, 12(2): 198-204

Research (Published online: 08-02-2019)

3. Staphylococcus aureus isolated from selected dairies of Algeria: Prevalence and susceptibility to antibiotics
Asmaa Manel Matallah, Leila Bouayad, Sofiane Boudjellaba, Faiza Mebkhout, Taha Mossadak Hamdi and Nadjia Ramdani-Bougussa
Veterinary World, 12(2): 205-210

Research (Published online: 09-02-2019)

4. Reduction of proteolysis of high protein silage from Moringa and Indigofera leaves by addition of tannin extract
Anuraga Jayanegara, Aldi Yaman and Lilis Khotijah
Veterinary World, 12(2): 211-217

Research (Published online: 09-02-2019)

5. Investigation of Mycobacterium paratuberculosis in Arabian dromedary camels (Camelus dromedarius)
Mohamed A. Salem, Wael M. El-Deeb, Ahmed A. Zaghawa, Fadel M. Housawi and Ahmed M. Alluwaimi
Veterinary World, 12(2): 218-223

Research (Published online: 09-02-2019)

6. Coinfection of diarrheagenic bacterial and viral pathogens in piglets of Northeast region of India
Hosterson Kylla, Tapan K. Dutta, Parimal Roychoudhury and Prasant K. Subudhi
Veterinary World, 12(2): 224-230

Research (Published online: 11-02-2019)

7. Genetic characterization of S1 gene of infectious bronchitis virus isolated from commercial poultry flocks in West Java, Indonesia
Rahajeng Setiawaty, Retno Damajanti Soejoedono and Okti Nadia Poetri
Veterinary World, 12(2): 231-235

Research (Published online: 12-02-2019)

8. Antibacterial and cytotoxic activities of the Syzygium polyanthum leaf extract from Malaysia
Muhammad Luqman Nordin, Abdul Aziz Othman, Arifah Abdul Kadir, Rumaizi Shaari, Abdinasir Yusuf Osman and Maizan Mohamed
Veterinary World, 12(2): 236-242

Research (Published online: 12-02-2019)

9. Analysis of bacterial contamination and antibiotic residue of beef meat from city slaughterhouses in East Java Province, Indonesia
Koesnoto Soepranianondo, Dhandy Koesoemo Wardhana, Budiarto and Diyantoro
Veterinary World, 12(2): 243-248

Research (Published online: 13-02-2019)

10. Morphology and morphometry of adult nematodes on Sumatran elephants (Elephas maximus sumatranus) in Way Kambas National Park area, Indonesia
Rahmania Prahardani, Lintang Winantya Firdausy, Yanuartono and Wisnu Nurcahyo
Veterinary World, 12(2): 249-253

Research (Published online: 13-02-2019)

11. Alteration in behavior of rat after chronic exposure to acetamiprid
Samiran Mondal, Saktipada Pradhan and Sunit K. Mukhopadhyay
Veterinary World, 12(2): 254-257

Research (Published online: 14-02-2019)

12. Antibiotic resistance: A cross-sectional study on knowledge, attitude, and practices among veterinarians of Haryana state in India

Thulasiraman Parkunan, Manju Ashutosh, Bharathy Sukumar, Jatinder Singh Chera, Sendhil Ramadas, B. Chandrasekhar, S. Ashok Kumar, Rachana Sharma, M. Santhosh Kumar and Sachinandan De
Veterinary World, 12(2): 258-265

Research (Published online: 15-02-2019)

13. A preliminary molecular survey of Babesia divergens and first evidence of Theileria annulata in cattle from Saudi Arabia
Mohamed W. Ghafar and Sayed A. M. Amer
Veterinary World, 12(2): 266-270

Research (Published online: 16-02-2019)

14. Stimulation of non-specific immunity, gene expression, and disease resistance in Nile Tilapia, *Oreochromis niloticus* (Linnaeus, 1758), by the methanolic extract of the marine macroalga, *Caulerpa scalpelliformis*
Omita Yengkhom, Konda Subramanian Shalini, P. A. Subramani and R. Dinakaran Michael
Veterinary World, 12(2): 271-276

Research (Published online: 20-02-2019)

15. Molecular characterization of pathogenic 4/91-like and QX-like infectious bronchitis virus infecting commercial poultry farms in Indonesia
Michael H. Wibowo, Teridah E. Ginting and Widya Asmara
Veterinary World, 12(2): 277-287

Research (Published online: 20-02-2019)

16. *Toxocara vitulorum* cuticle glycoproteins in the diagnosis of calves' toxocariasis
Eman E. El Shanawany, Soad E. Hassan, Adel A.- H. Abdel-Rahman and Eman H. Abdel-Rahman
Veterinary World, 12(2): 288-294

Research (Published online: 22-02-2019)

17. Assessment of safety and quality of fermented milk of camels, cows, and goats sold and consumed in five localities of Burkina Faso
Hama Cissé, Jean Ulrich Muandze-Nzambe, Namwin Siourime Somda, Adama Sawadogo, Soungalo Moustapha Drabo, Francois Tapsoba, Cheikna Zongo, Yves Traore and Aly Savadogo
Veterinary World, 12(2): 295-304

Research (Published online: 22-02-2019)

18. Reproductive performance of pigs raised by intensive management system in Abuja, Nigeria
Kenneth Owoicho Abah, Joy Iyojo Itodo, Simon Azubuike Ubah and Ibrahim Shettima
Veterinary World, 12(2): 305-308

Research (Published online: 23-02-2019)

19. Antiretroviral activity of *Pterois volitans* (red lionfish) venom in the early development of human immunodeficiency virus/acquired immunodeficiency syndrome antiretroviral alternative source
Andy Noorsaman Sommeng, R. Muhammad Yusuf Arya, Mikael Januardi Ginting, Diah Kartika Pratami, Heri Hermansyah, Muhamad Sahlan and Anondho Wijanarko
Veterinary World, 12(2): 309-315

Research (Published online: 23-02-2019)

20. Chemotherapeutic control of Gram-positive infection in white sea bream (*Diplodus sargus*, Linnaeus 1758) broodstock
Nadia G. M. Ali, Ibrahim M. Aboyadak and Heba S. El-Sayed
Veterinary World, 12(2): 316-324

Research (Published online: 23-02-2019)

21. Comparative efficacy of two-dimensional mode and color Doppler sonography in predicting gender of the equine fetus
M. Mebarki, R. Kaidi, A. Azizi and M. Basbaci
Veterinary World, 12(2): 325-330

Research (Published online: 24-02-2019)

22. Isolation and molecular characterization of *Mannheimia haemolytica* and *Pasteurella multocida* associated with pneumonia of goats in Chhattisgarh
Nidhi Rawat, Varsha Rani Gilhare, Krishna Kumar Kushwaha, Deeksha Dipak Hattimare, Foziya Farzeen Khan, Rajesh Kumar Shende and Dhananjay Kumar Jolhe
Veterinary World, 12(2): 331-336

Research (Published online: 27-02-2019)

23. Resveratrol mitigates hypercholesterolemia exacerbated hyperthermia in chronically heat-stressed rats
Hosam Al-Tamimi, Amani Al-Dawood, Saddam Awaishesh and Tony Abdalla
Veterinary World, 12(2): 337-344

Research (Published online: 28-02-2019)

24. Comparative effects of amiodarone and dronedarone treatments on cardiac function in a rabbit model
Worakan Boonhoh, Anusak Kijtawornrat and Suwanakiet Sawangkoon
Veterinary World, 12(2): 345-351

Prevalence and diversity of gastrointestinal protozoa in Madura cattle at Bangkalan Regency, East Java, Indonesia

Poedji Hastutiek¹, Wiwik Misaco Yuniarti², Mufasirin Djaeri¹, Nunuk Dyah Retno Lastuti¹, Endang Suprihati¹ and Lucia Tri Suwanti^{1,3}

1. Department of Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga, Jl. Mulyorejo, Kampus C Unair, Surabaya, 60111; 2. Department of Clinical Science, Faculty of Veterinary Medicine, Universitas Airlangga, Jl. Mulyorejo, Kampus C Unair, Surabaya, 60111; 3. Department of Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga, Institute of Tropical Diseases, Universitas Airlangga, Jl. Mulyorejo, Kampus C Unair, Surabaya, 60111.

Corresponding author: Poedji Hastutiek, e-mail: poedjihastutiek@gmail.com

Co-authors: WMY: wiwikmisaco@yahoo.com, MD: mufasirin@fkh.unair.ac.id, NDRL: nunukdyah53@gmail.com, ES: esuprihati@yahoo.co.id, LTS: tswant@gmail.com

Received: 21-09-2018, **Accepted:** 20-12-2018, **Published online:** ***

doi: 10.14202/vetworld.2019.198-204 **How to cite this article:** Hastutiek P, Yuniarti WM, Djaeri M, Lastuti NDR, Suprihati E, Suwanti LT (2019) Prevalence and diversity of gastrointestinal protozoa in Madura cattle at Bangkalan Regency, East Java, Indonesia, *Veterinary World*, 12(2): 198-204.

Abstract

Aim: This study aimed to describe the gastrointestinal protozoa in Madura cattle at Bangkalan Regency, East Java, Indonesia.

Materials and Methods: A total of 500 samples of Madura cattle feces were collected from 10 districts at Bangkalan Regency. Those ten districts represent the lowland and upland areas, and each district was represented by one village. The collected feces were examined using native, sedimentation, and floating methods. The species identification was determined by their morphology.

Results: There were 357 (71.4%) samples positively infected with protozoan. The highest rate of sample with protozoan infection was at Kamal District (88.23%), and Bangkalan District (52.83%) was the lowest one. There were six species of protozoa that infected gastrointestinal tract; those are *Eimeria* spp., *Balantidium* spp., *Isospora* spp., *Blastocystis* spp., *Entamoeba* spp., and *Cryptosporidium* spp. The highest number of protozoa found in this research was *Eimeria* (53.42%) followed by *Blastocystis* (14.43%). In this study, we found that 295 samples (58.76%) infected by one kind of protozoa, 53 samples (10.56%) infected by two kinds of protozoa, and 11 samples (2.19%) infected by three kinds of protozoa. In addition, there were 65.54% of bulls infected with protozoa, considerably lower than cows (72.97%). Cattle aged 6 months-2 years old (73.39%) and >2 years old (71.25%) are known more prone to protozoan infections than cattle aged <6 months (66.15%).

Conclusion: The present study revealed that protozoan infection of cattle is common in Bangkalan Regency. Studies focused on determining that the prevalence of protozoan, risk factors for the parasitism, and the geographic distribution are needed and will be effective guide for prevention and control measures.

Keywords: Bangkalan, Madura cattle, mapping, protozoa.

Introduction

Madura cattle is one of the Indonesian local cattle that are widely developed in East Java, especially in Madura Island. Madura cattle have great potentials to be developed because they are genetically tolerant to hot climates and marginal environments, resistant to tick infestations, and highly adaptable to low feed quality, as well as require less food compared to imported cattle. Besides, Madura cattle are easy to maintain, easy to breed, and resistant to various diseases [1]. The population of Madura cattle in Bangkalan Regency in 2016 was 200,279 and predicted to increase every year parallel to the promotion

of crossbred artificial insemination program between Madura cattle and Limousine cattle (Madrasin) [2].

Beef cattle as a potential commodity in the development of rural farming must be supported by an adequate maintenance system. The business of animal husbandry is faced with problems of reproductive disorders and chronic parasitic diseases, especially protozoan infections. Gastrointestinal disease needs special attention because it can be an obstacle that affects the acceleration of livestock development in the countryside so that it can cause economic losses due to decreased livestock productivity, decreased weight, quality of meat, skin, and internal organs, growth retardation in young animals, and danger of zoonoses. Delay weight gain in cattle with protozoan infection can reach >40% compared with healthy cows [3,4].

Based on the examination results of fecal samples gathered from Madura cattle butchered at slaughterhouse at Surabaya, it was found that those cattle were infected by several kinds of protozoa species, such as *Eimeria* spp., *Balantidium* spp., and *Entamoeba* spp.

Copyright: Hastutiek, et al. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

(unpublished data). However, there has been no research published about protozoan infections in the gastrointestinal tract of Madura cattle in Bangkalan Regency, Madura, Indonesia.

Materials and Methods

Ethical approval

The present study was based on the laboratory examination of cattle feces without treatment. The samples were collected as per standard sample collection procedure, directly from the rectum without disturbing the animals, and were accompanied by a responsible veterinarian.

Study area

This research was conducted at 10 districts located in coastal areas (each district was represented by one village) in Bangkalan Regency with high live-stock population. Fecal sampling was conducted at sites with different altitudes. Bangkalan, Socah, and Kamal districts are located on lowland with altitude <25 m above sea level, while Tanjung Bumi, Sepuluh, Klampis, Arosbaya, Burneh, Tragah, and Labang districts are located on a plateau with an altitude of 25-200 m above sea level.

Fecal sample collection and analysis

A total of 500 fecal samples were collected for this research. Fecal samples were taken randomly from bulls and cows and then divided based on age such as <6 months, 6 months-2 years, and >2 years. Fecal sampling was conducted from April to May 2017. During fecal sampling process, questionnaire and interview with the farmer were also conducted to obtain certain information about farmer (name, age, gender, education, and breeding experience) and cattle (breed and quantity, gender, age, type of maintenance, treatment that has ever given, the other reared cattle, type or material of the enclosure, environmental condition, or livestock care).

The samples were collected directly from the rectum and brought to the laboratory in mini zip locked polythene bags and added with 2.5% potassium bichromate for examination. Each plastic bag was labeled with registered sample number and then stored in a container with ice. Afterward, the fecal samples were examined using native methods, as well as sedimentation and modified Fulleborn's floating methods [5]. To determine the existence of protozoa, identification key methods were carried out [6]. The positive result of protozoan-infected cattle could be determined when protozoa were found during examination using one of those methods. The prevalence of the protozoan infections was expressed in percentage value using the following formula:

$$P = (\text{Positive results: Number of samples}) \times 100\%.$$

Statistical analysis

The data obtained will be analyzed descriptively and presented in the form of the prevalence of

protozoan infections based on district, kind of protozoa, sex, and age.

Results and Discussion

Based on this research, it was found that there were 357 from 500 samples (71.4%) positively infected by protozoa. The highest prevalence of protozoan infections was found in Kamal district, in which 88.23% (45/51) samples were positive, followed by Sepuluh district 86.8% (46/53), Klampis district 80% (32/40), Tanjung Bumi district 75% (36/48), Arosbaya district 74.51% (38/51), Burneh district 68% (34/50), Labang district 67.31% (35/52), Socah district 67.28% (37/55), Tragah district 57.14% (28/49), and Bangkalan district 52.83% (28/53) (Table-1).

The high prevalence in this study was in accordance with the study of Volpato *et al.* [7], which reported the prevalence of intestinal protozoan infection in dairy calf in Brazil. Those ten districts are located in coastal areas and are well known as livestock meeting points from several locations as well as temporary shelters for cattle that will be traded out of the island or for Idul Adha. On the other hand, the results of this research indicate that the prevalence of protozoan infections in Madura cattle from those 10 districts was high. Therefore, routine monitoring for protozoan infections should be preceded by performing fecal examination so that the infections can be completely controlled to improve the health and productivity of the cattle.

Research result showed that, from 359 positive samples, there were 295 (58.76%) samples infected by one kind of protozoan, 53 (10.56%) samples by two kinds of protozoan, and 11 (2.19%) samples infected with three kinds of protozoan.

The highest of single species protozoan infection was at Klampis district 80% (32/40), while the lowest was at Bangkalan district 49.06% (26/53). District with the highest infection of two species protozoan was Sepuluh district with 26.42% (14/53), whereas the lowest one at Bangkalan district with only 3.77% (2/53). At Kamal district, there were 9.8% (5/51) cattle infected with three kinds of protozoan species. At Labang district, Arosbaya district and Socah district

Table-1: The prevalence of protozoan infection in the gastrointestinal of Madura cattle in each district in Bangkalan Regency.

District	Number of samples	Number of positive samples (%)
Mangkong Arosbaya	51	38 (74.51)
Bator Ma'adan Klampis	40	32 (80)
Keleyan Socah	53	35 (66.04)
Gili Anyar Kamal	51	45 (88.24)
Maneron Sepuluh	53	46 (86.79)
Bumi Anyar Tanjung Bumi	48	36 (75)
Bringen Labang	52	35 (67.31)
Kemoneng Tragah	49	28 (57.14)
Ketengan Burneh	50	34 (68)
Kramin Bangkalan	53	28 (52.83)
Total	500	357 (71.4)

were 7.69% (4/52), 1.96% (1/51), and 1.82% (1/55) respectively (Table-2).

The results showed that there were six species of protozoa infecting the gastrointestinal tract of Madura cattle, i.e., *Eimeria* spp., *Balantidium* spp., *Isospora* spp., *Blastocystis* spp., *Entamoeba* spp., and *Cryptosporidium* spp. The result also indicated that one cattle could be infected by one, two, or even more three kinds of protozoa species at once (Table-2) (Figures-1-6).

Eimeria is the most common species of protozoan-infected Madura cattle in 10 districts, whether it was single or mixed infections. The second common species of protozoan-infected Madura cattle in Bangkalan district, both single and mixed infections, was *Blastocystis*. However, *Blastocystis* was

not found infecting cattle in Arosbaya, Klampis, and Burneh districts.

Some types of protozoa found in this research showed to have zoonotic potentials, such as *Balantidium* spp., *Entamoeba* spp., *Blastocystis* spp., and *Cryptosporidium* spp. *Cryptosporidium parvum* oocyte is usually found in exoskeleton and gastrointestinal tract of *Musca domestica* on dairy farms. Flies as mechanical vectors have the ability to spread diarrheal diseases caused by protozoa as they can travel up to 20 miles [8]. *Cryptosporidium bovis* oocyte can infect six species of livestock and poultry in Tunisia [9]. The prevalence of *Cryptosporidium* infection in cattle in Northern Nigeria reported to be 22.3% [10,11], whereas the prevalence of *Blastocystis* spp., *Giardia* spp., and *Entamoeba* spp. was 14.6%, 12.45%, and 7.45%, respectively [12]. The

Table-2: The prevalence of protozoan species in the gastrointestinal tract of Madura cattle in Bangkalan Regency.

Districts	Samples positively infected with protozoa						Number of positive samples (%)
	One type of protozoan species	Total	Two types of protozoan species	Total	Three types of protozoan species	Total	
Mangkon	<i>Eimeria</i>	31	<i>Eimeria</i>	5	<i>Eimeria</i>	1	38 (74.51)
Arosbaya	<i>Balantidium</i>	1	<i>Balantidium</i>		<i>Balantidium Isospora</i>		
Bator	<i>Eimeria</i>	32	-		-		32 (80)
Ma'adan							
Klampis							
Keleyan	<i>Eimeria</i>	33	<i>Eimeria</i>	1	<i>Eimeria</i>	1	35 (66.04)
Socah			<i>Blastocystis</i>		<i>Blastocystis</i>		
					<i>Balantidium</i>		
Gili Anyar	<i>Eimeria</i>	12	<i>Eimeria</i>	2	<i>Eimeria</i>	3	45 (88.23)
Kamal			<i>Balantidium</i>		<i>Blastocystis</i>		
					<i>Balantidium</i>		
	<i>Blastocystis</i>	14	<i>Eimeria</i>	9	<i>Eimeria</i>	2	
			<i>Blastocystis</i>		<i>Blastocystis</i>		
	<i>Balantidium</i>	2	<i>Blastocystis</i>	1	<i>Entamoeba</i>		
			<i>Balantidium</i>				
Maneron	<i>Eimeria</i>	22	<i>Eimeria</i>	2	-	-	46 (86.8)
Sepuluh			<i>Balantidium</i>				
	<i>Blastocystis</i>	10	<i>Eimeria</i>	11			
			<i>Blastocystis</i>				
			<i>Blastocystis</i>	1			
			<i>Balantidium</i>				
Bumi Anyar	<i>Eimeria</i>	23	<i>Eimeria</i>	5	-	-	36 (75)
Tanjung	<i>Blastocystis</i>	7	<i>Blastocystis</i>				
Bumi	<i>Balantidium</i>	1					
Bringen	<i>Eimeria</i>	10	<i>Eimeria</i>	3	<i>Eimeria</i>	1	35 (67.31)
Labang			<i>Blastocystis</i>		<i>Blastocystis</i>		
					<i>Entamoeba</i>		
	<i>Blastocystis</i>	14	<i>Blastocystis</i>	1	<i>Eimeria</i>	1	
			<i>Balantidium</i>		<i>Blastocystis</i>		
					<i>Balantidium</i>		
	<i>Entamoeba</i>	1	<i>Blastocystis</i>	1	<i>Eimeria</i>	1	
			<i>Entamoeba</i>		<i>Blastocystis</i>		
					<i>Cryptosporidium</i>		
			<i>Blastocystis</i>	1	<i>Eimeria</i>	1	
			<i>Cryptosporidium</i>		<i>Blastocystis</i>		
					<i>Isospora</i>		
Kemoneng	<i>Eimeria</i>	12	<i>Eimeria</i>	5	-	-	28 (57.14)
Tragah	<i>Blastocystis</i>	11	<i>Blastocystis</i>				
Ketengan	<i>Eimeria</i>	13	<i>Eimeria</i>	2	-	-	34 (68)
Burneh			<i>Balantidium</i>				
	<i>Balantidium</i>	18	<i>Eimeria</i>	1			
			<i>Entamoeba</i>				
Kramat	<i>Eimeria</i>	21	<i>Eimeria</i>	2	-	-	28 (52.83)
Bangkalan	<i>Balantidium</i>	4	<i>Balantidium</i>				
	<i>Blastocystis</i>	1					



Figure-1: *Blastocystis* spp. (400×).



Figure-4: *Eimeria* (400×).



Figure-2: *Cryptosporidium* spp. (400×).



Figure-5: *Isospora* (400×).



Figure-3: *Balantidium* (400×).




Figure-6: *Entamoeba* spp. (400×)

high prevalence of *Blastocystis* spp. infection can lead to death in livestock [13,14]. The prevalence of protozoan infections in the gastrointestinal tract of dogs in France was 42.2%, and caused by *Blastocystis* spp. and *Cryptosporidium* spp. Both can potentially trigger infections to their hosts [15]. Meanwhile, the prevalence of *Entamoeba bovis* infection was 36% and mostly found in cattle suffering from diarrhea [16].

The number of samples with single infection caused by *Eimeria* spp. was 211, in which the highest was found in Socah district with 35 samples. Single infection caused by *Blastocystis* spp. was found in 57 samples and found at seven districts, in which the highest was found in Kamal and Labang districts, with 14 samples of each. Samples with single infection caused by *Balantidium* spp. was 26, in which the highest one was found in Burneh district with 18 samples.

Single infection caused by *Entamoeba* spp. only found in one cattle in Labang district.

Samples infected by two species of protozoan, such as *Eimeria* and *Blastocystis*, were found in 34 at six districts. Sepuluh district with 11 samples was the highest. There were five samples infected with three species of protozoa (*Eimeria* spp., *Blastocystis* spp., and *Balantidium* spp.), which were found in Kamal district.

Based on sex, the prevalence of protozoan infections in bulls 65.54% (116/169) was lower than cows 72.97% (243/333). The comparison of the prevalence of protozoan infections based on sex in each district is shown in Table-3.

The prevalence of protozoan infections in cattle <6 months of age was 68.18% (45/66), with the highest case found in Klampis, Tanjung Bumi, and Burneh districts (100%), and the lowest was found in Sepuluh district (25%). Furthermore, the prevalence of protozoan infections in cattle aged from 6 months

to 2 years was 73.39% (91/124). The highest prevalence was found in Arosbaya district (87.1%), and the lowest was found in Bangkalan district (23.08%). For the cattle >2 years, the prevalence of protozoan infections was 71.47% (223/312), the highest was found in Sepuluh district (94.12%), and the lowest was in Arosbaya district (52.63%) (Table-4).

Research conducted in Iran revealed that there were four types of protozoa species found in pigs with a prevalence of 64% [17], while the prevalence of protozoan infections in the gastrointestinal tract of carnivores in Iran was 80.4% [18]. Another research in India showed that 83.08% of cattle were infected by endoparasites, with the higher risk factor found in cows (85.97%) than bulls (69.23%) and in the adult ones that aged >6 months (85.97%) compared to those that aged <6 months (61.17%) [19]. Similarly, the prevalence of protozoan infections in cows in this research was also higher than in bulls due to predisposing genetic factor, hormonal factor, stress-reducing

Table-3: The prevalence of protozoan infections in the gastrointestinal tract of Madura cattle in Bangkalan Regency based on sex.

Districts	Bulls		Cows		Total of samples	
	Number of samples	Positive samples (%)	Number of samples	Positive samples (%)	Number of samples	Positive samples (%)
Mangkon Arosbaya	13	6 (46.15)	38	32 (84.21)	51	38 (74.51)
Bator Ma'adan Klampis	10	8 (80)	30	24 (80)	40	32 (80)
Keleyan Socah	14	6 (42.86)	39	29 (74.44)	53	37 (67.28)
Gili Anyar Kamal	19	15 (78.95)	32	30 (93.75)	51	45 (88.23)
Maneron Sepuluh	23	19 (82.61)	30	27 (90)	53	46 (86.8)
Bumi Anyar Tanjung Bumi	22	17 (77.27)	26	19 (73.08)	48	36 (75)
Bringen Labang	4	3 (75)	48	32 (66.67)	52	35 (67.31)
Kemoneng Tragah	8	3 (37.5)	41	25 (60.98)	49	28 (57.14)
Ketengan Burneh	50	34 (68)	-	-	50	34 (68)
Kramin Bangkalan	4	3 (28)	49	25 (50.02)	53	28 (52.83)
Total	167	114 (68.26)	333	243 (72.97)	500	357 (71.4)

Table-4: The prevalence of protozoan infections in the gastrointestinal tract of Madura cattle in Bangkalan Regency based on age.

Districts	Age of samples				Number of samples	Positive samples		
	<6 months	Positive samples (%)	>6 months-2 years	Positive samples (%)			>2 years	Positive samples (%)
Mangkon Arosbaya	2	1 (50)	31	27 (87.1)	19	10 (52.63)	51	38
Bator Ma'adan Klampis	6	6 (100)	9	7 (77.78)	25	19 (76)	40	32
Keleyan Socah	17	12 (70.59)	10	8 (80)	26	15 (57.69)	55	37
Gili Anyar Kamal	2	1 (50)	18	15 (83.33)	31	29 (93.55)	51	45
Maneron Sepuluh	4	1 (25)	14	13 (86.67)	35	32 (94.12)	53	46
Bumi Anyar Tanjung Bumi	3	3 (100)	8	6 (75)	37	27 (72.97)	48	36
Bringen Labang	7	5 (71.43)	2	1 (50)	43	29 (67.44)	52	35
Kemoneng Tragah	10	8 (80)	2	-	37	20 (54.05)	49	28
Ketengan Burneh	1	1 (100)	16	11 (68.75)	33	22 (66.67)	50	34
Kramin Bangkalan	13	5 (38.46)	13	3 (23.08)	27	20 (74.07)	53	28
Total	65	43 (65.15)	122	91 (74.59)	313	223 (71.25)	500	357 (71.40)

immune factor to infection, parturition factor, and lactation factor, resulting in weakness and malnutrition.

The prevalence of protozoan infections in Madura cattle in this research that aged from 6 months to 2 years was also higher than those aged <6 months due to their grazing habit in larger areas that were contaminated with protozoa. Hence, the prevalence of the infections in calf is always lower, since they were less exposed to larger areas contaminated with protozoa and it is also because the calf is still suckling.

Madura cattle are reared in cages or not, vulnerable to protozoan infection that contaminate the grass they eat. If one of the cattle was infected by protozoa, the area then would be considered as a good place for various types of protozoa to develop. Forages that were given to Madura cattle actually have been through the process of forging; however, the prevalence of protozoan infection in the gastrointestinal tract is still high because their feed and water have been contaminated by protozoa.

Rainy season that increased the humidity is also suspected to be the cause of the high protozoan infections in the gastrointestinal tract of Madura cattle. In other words, rainy season with high humidity and low temperatures is a condition favored by protozoa to thrive. Parasitic infections in livestock were also affected by various factors, including geographical location, environmental conditions, cage quality, sanitation and hygiene, cage density, temperature, humidity, as well as vegetation [6,20].

In this study, management of cage and sanitation is also known to be unfit since feces that were cleared out from the cages were dumped right around the cages, resulting in moisten up the cage which can elevate the risk of reinfection. The construction of the cages is also known to be very traditional and still not equipped with the feces and urine disposal line. The infections caused by one, two, and three types of protozoa species in this research may be caused by decreased immune system condition to resist against protozoan infection [3,20].

Poor ventilation system that can reduce sunlight to enter the cage will make the ground floor of the cage moist, so that it can be a good medium for parasitic growth. Pigs and cows in Korea are also known to be infected with *Balantidium coli* and *Entamoeba* spp., in which cysts and oocysts potentially pollute and contaminate their environment due to poor sanitation [21].

In addition, the data collected from the questionnaires indicated that most farmers were over 50 years old with low education (never attended school) or up to junior high school. Therefore, they had low understanding in receiving information as well as new ideas and technology to improve their cattle performance. High incidence of protozoan infection in this study indicates that gastrointestinal tract infections in Madura cattle were chronic and required treatment.

Conclusion

The prevalence of protozoan infections on the gastrointestinal tract of Madura cattle was 71.51%. They were *Eimeria* spp., *Balantidium* spp., *Isoospora* spp., *Blastocystis* spp., *Entamoeba* spp., and *Cryptosporidium* spp. About 58.76% samples were infected by one, 53.56% samples were infected by two types, and 2.19% were infected with three types of protozoan species. The prevalence of protozoan infections in cows and bulls was 72.97% and 65.54%, respectively. Based on age, the prevalence of protozoan infections in Madura cattle was 68.18%, 73.39%, and 71.47%, for 6 months, 6 month-2 years, and >2 years, respectively.

Authors' Contributions

PH, MD, and NDRL planned and designed the whole study. ES, LTS, and WMY collected sample, carried out the whole work, and wrote the manuscript. PH helped in identification of parasites and microscopic examination. MD, ES, and WMY helped during manuscript writing, cross-checking, and revision. All authors read and approved the final manuscript.

Acknowledgments

The authors are highly thankful to Dean and Colleague of the Faculty of Veterinary Medicine, Universitas Airlangga for their cooperation and provision of research facilities. The fund for the study was provided by Faculty of Veterinary Medicine, Universitas Airlangga, Indonesia (Dean of Faculty of Veterinary Medicine Decision Letter No. 56/UN3.1.6/2017).

Competing Interests

The authors declare that they have no competing interests.

Publisher's Note

Veterinary World remains neutral with regard to jurisdictional claims in published institutional affiliation.

References

1. Nurgartiningsih, V.M.A. (2016) Evaluation of performance in female Madura cattle in Madura Island, Indonesia. *Anim. Prod.*, 18(3): 125-130.
2. Widi, T.S.M., Udo, H.M.J., Oldenbroek, K., Budi Satria, I.G.S., Baliarti, E. and van der Zijpp, A.J. (2013) Unique cultural values of Madura cattle: Is cross-breeding a threat? *Anim. Genet. Resour.*, 54(1): 141-152.
3. Thompson, R.C.A. and Smith, A. (2011) Zoonotic enteric *Protozoa*. *Vet. Parasitol.*, 182(1): 70-78.
4. Maharana, B.R., Kumar, B., Sudhakar, N.R., Behera, S.K. and Patbandha, T.K. (2016) Prevalence of gastrointestinal parasites in bovines in and around Junagadh (Gujarat). *J. Parasitol. Dis.*, 40(4): 1174-1178.
5. Heidari, H., Dehkordi, Z.S., Moayedi, R. and Gharekhani, J. (2014) Occurrence and diversity of *Eimeria* species in cattle in Hamedan Province, Iran. *Vet. Med.*, 59(6): 271-275.
6. Soulsby, E.J.L. (1986) *Helminths, Arthropods and Protozoa of Domesticated Animals*. 7th ed. Bailliere Tindall, London.
7. Volpato, A., Tonin, A.A., Machado, G., Stefan, L.M.,

- Campigotto, G., Glombowsky, P., Galli, G.M., Favero, J.F. and da Silva, A.S. (2017) Gastrointestinal *Protozoa* in dairy calves: Identification of risk factors for infection. *Rev. MVZ Cordoba*, 22(2): 5910-5924.
8. Szostakowska, B., Lozowska, W.K., Racewicz, M., Knight, R., Tamang, L., Myjak, P. and Graczyk, T.K. (2004) *Cryptosporidium parvum* and *Giardia lamblia* recovered from flies on a cattle farm and in a landfill. *Appl. Environ. Microbiol.*, 70(6): 3742-3744.
 9. Soltane, R., Guyot, K., Dei-Cas, E. and Ayadi, A. (2007) Prevalence of *Cryptosporidium* Spp. (*Eucoccidiorida: Cryptosporidiidae*) in seven species of farm animals in Tunisia. *Parasite*, 14(4): 335-338.
 10. Adamu, S.G., Adamu, N.B., Aliyu, A.U., Atsanda, N.N., Mustapha, F.B., Muhammad, Y.A. and Umaru G.A. (2015) Prevalence of *Cryptosporidium* infection in cattle in Maiduguri, North Eastern Nigeria. *Bangl. J. Vet. Med.*, 13(1): 25-28.
 11. Ayinmode, A.B. and Fagbemi, B.O. (2010). Prevalence of *Cryptosporidium* infection in cattle from South Western infection in cattle from South Western Nigeria. *Vet. Arch.*, 80(6): 723-731.
 12. Badparva, E., Fallahi, S. and Aarab-Mazar, Z. (2015) *Blastocystis*: Emerging protozoan with high prevalence in Iran. *Nov. Biomed.*, 3(4): 214-221.
 13. Stensvold, C.R., Suresh, G.K., Tan, K.S., Thompson, R.C., Traub, R.J., Viscogliosi, E., Yoshikawa, H. and Clark, C.G. (2007). Terminology for *Blastocystis* of subtype a consensus. *Trends Parasitol.*, 23(3): 93-96.
 14. Yoshikawa H., Wu, Z., Pandey, K., Pandey, B.D., Sherchand, J.B. and Yanagi T. (2009) Molecular characteristic of *Blastocystis* isolates from children and rhesus monkey in Kathmandu, Nepal. *Vet. Parasitol.*, 160(3-4): 295-300.
 15. Orman M., Bories, J., El Safadi, D., Poirel, M.T., Gantois, N., Benamrouz-Vanneste, S., Delhaes, L., Hugonnard, M., Certad, G., Zenner, L. and Viscogliosi, E. (2015) Prevalence and genetic diversity of the intestinal parasite *Blastocystis* spp. and *Cryptosporidium* Spp, in house-hold dogs in France and evaluation of zoonotic transmission risk. *Vet. Parasitol.*, 214(1-2): 167-170.
 16. Al-shabbani, A.H. (2016) Direct detection of *Entamoeba bovis* in calves infected by diarrhea by using polymerase chain reaction technique. *J. Vet. Med. Sci.*, 7(1): 132-137.
 17. Yaghobii, K., Sarkari, B., Mansouri, M. and Motazedian, M.H. (2016) Zoonotic intestinal *Protozoa* of the wild boars, *Sus scrofa*, in Persian Gulf's Coastal Area (Bushehr Province), Southwestern Iran. *Vet. World*, 9(2): 1047-1050.
 18. Sarvi, S., Daryani, A., Syarif, M., Rahimi, M.T., Kohansal, M.H., Mirshafiee, S., Siyatpanah, S., Hossein, A.A. and Gholami, S. (2018) Zoonotic intestinal parasites of Carnivora: A systemic review in Iran. *Vet. World*, 11(11): 58-65.
 19. Singh, E., Kaur, P., Singla, L.D. and Bal, S. (2017) Prevalence of gastrointestinal parasitism in small ruminant in Western Punjab, India. *Vet. World*, 10(1): 61-66.
 20. Marskole, P., Verma, Y., Dixit, A.K. and Swamy, M. (2016) Prevalence and burden of gastrointestinal parasites in cattle and buffaloes in Jabalpur, India. *Vet. World*, 9(11): 1214-1217.
 21. Ismail, H.A., Jeon, H.K., Yu, Y.M., Do, C. and Lee, Y.H. (2010) Intestinal parasite infections in pigs and beef cattle in rural areas of Chungcheongnam-do, Korea. *Korean J. Parasitol.*, 48(4): 347-349.
