

THE SOUTHEAST ASIAN JOURNAL OF TROPICAL MEDICINE AND PUBLIC HEALTH



Vol 48 No. 6

November 2017



Official Publication of the SEAMEO Regional Tropical Medicine
and Public Health Network (TROPMED)



The SEAMEO Regional Tropical Medicine and Public Health Project was established in 1967 to help improve the health and standard of living of the peoples of Southeast Asia by pooling manpower resources of the participating SEAMEO member countries in a cooperative endeavor to develop and upgrade the research and training capabilities of the existing facilities in these countries. By promoting effective regional cooperation among the participating national centers, it is hoped to minimize waste in duplication of programs and activities. In 1992 the Project was renamed the SEAMEO Regional Tropical Medicine and Public Health Network.*

**SEAMEO=Southeast Asian Ministers of Education Organization*

THE SOUTHEAST ASIAN JOURNAL OF TROPICAL MEDICINE AND PUBLIC HEALTH

Editorial Panel

Honorary Editor	Tan Chongsuphajsiddhi (Thailand)
Editor	Vicente Y Belizario (Philippines)
Executive Editor	Suvanee Supavej (Thailand)
Consulting Editor	Peter D Echeverria (Thailand)
Editorial Office	Vimolpatana Indradat (Administrator) The Southeast Asian Journal of Tropical Medicine and Public Health SEAMEO TROPMED Network 420/6 Ratchawithi Road Bangkok 10400, Thailand Tel: +66 (0) 23549197, 23549145, 23549146, 26444331, 26445135 Fax: +66 (0) 23544337, 23549144 E-mail: tmseameo@mahidol.ac.th Website: http://www.tm.mahidol.ac.th/seameo/publication.htm http://www.seameotropmednetwork.org

Editorial Board

Alejandro Herrin (Philippines), Chansuda Wongsrichanalai (Cambodia), David Evans (Switzerland), Dwip Kitayaporn (Thailand), Eduardo Gotuzzo (Peru), J Kevin Baird (Indonesia), Jong Yil Chai (Korea), May Ho (Canada), Mark Robson (USA), Mediadora Sanie (Philippines), Michael Good (Australia), Myint-Lwin (Myanmar), Nicholas J White (UK), Nick Walters (USA), Pirom Kamolratanakul (Thailand), Pornthep Chanthavanich (Thailand), Prapon Wilairat (Thailand), Srisin Khusmith (Thailand), Stephen L Hoffman (USA), Stephen W King (Thailand), Virasakdi Chongsuvivatwong (Thailand), Yong Poovorawan (Thailand)

Advisory Board

Caridad A Ancheta (Philippines), CP Ramachandran (Malaysia), David Molyneux (UK), David Warrell (UK), HM Gilles (UK), Jane Baltazar (Philippines), Lye Munn Sann (Malaysia), Louis H Miller (USA), M Jegathesan (Malaysia), Santasiri Sornmani (Thailand), Soemilah Sastroamidjojo (Indonesia), Somei Kojima (Japan), Tsutomu Takeuchi (Japan), Walther H Wernsdorfer (Austria), Tang Lin Hua (China), Nina Gloriana (Philippines)

SEAMEO REGIONAL TROPICAL MEDICINE AND PUBLIC HEALTH NETWORK

TROPMED GOVERNING BOARD MEMBERS

BRUNEI	Dr HAJAH RAFIDAH BINTI HAJI GHARIFF
CAMBODIA	HE Prof Dr ENG HUOT
INDONESIA	Prof Dr TJANDRA YOGA ADITAMA
LAO PDR	Dr SOMCHANH XAYSIDA
MALAYSIA	Datuk Dr NOOR HISHAM ABDULLAH
MYANMAR	Dr KYAW KHAING
PHILIPPINES	Dr PAULYN JEAN B ROSELL-UBIAL
SINGAPORE	Prof JOHN EU-LI WONG
THAILAND	Dr SUWANNACHAI WATTANAYINGCHAROENCHAI
TIMOR LESTE	Dr JOAO MARTINS
VIETNAM	Dr TRAN THI GIANG HUONG

TROPMED CENTRAL OFFICE

Dr PRATAP SINGHASIVANON	<i>Secretary General/Coordinator</i>
Dr MA SANDRA B TEMPONGKO	<i>Deputy Coordinator</i>
Mrs VIMOLPATANA INDRADAT	<i>Publications</i>

The Southeast Asian Journal of Tropical Medicine and Public Health is published by the SEAMEO Regional Tropical Medicine and Public Health Network (SEAMEO TROPMED Network).

One annual volume comprises of six bi-monthly issues (January, March, May, July, September, and November). Annual subscription for 6 issues is USD 310.00 (including postage). Single copies USD 60.00. Subscription payment in US dollars must be made by credit card (VISA or MASTER). Use the order form on the last page of our journal and send to SEAMEO TROPMED Network, 420/6 Ratchawithi Road, Bangkok 10400, Thailand.

The Southeast Asian Journal of Tropical Medicine and Public Health has been selected for coverage in Thomson Reuters information services, beginning with Vol 40 (1) 2009: Science Citation Index (SciSearch®), Journal Citation Reports/Science Edition, and Current Contents®/Clinical Medicine, in addition to the existing inclusion in Biological Abstracts, BIOSIS Previews, and Zoological Record.

No part of this Journal may be reproduced in any form by photostat, microfilm or any other means.
ISSN: 0125-1562

THE SOUTHEAST ASIAN JOURNAL OF TROPICAL MEDICINE AND PUBLIC HEALTH

Volume 48

Number 6

November 2017

CONTENTS

Page

Molecular epidemiology of <i>Blastocystis</i> sp in animals reared by the aborigines during wet and dry seasons in rural communities, Pahang, Malaysia	1151
SA Noradilah, TS Anuar, N Moktar, IL Lee, F Salleh, SNAAManap, NSHM Mohtar, SM Azrul, WO Abdullah, SA Nordin, SR Abdullah	
Review. Chloroquine as a second line treatment for malaria in Lao PDR: risks and benefits	1161
D Duangdany, X Xu, W Yao	
Permot (<i>Passiflora foetida</i> Linn.) leaf extracts as bioinsecticide against <i>Aedes aegypti</i> larvae	1169
P Hastutiek, A Sunarso, RH Prasetyo	
Diversity, seasonal abundance and biting activity of <i>Anopheles</i> species in relation to climatic factors in northeastern Thailand	1175
P Poolphol, RE Harbach, P Sriwichai, W Srisuka, K Aupalee, K Taai, S Thongsahuan, S Uttamangkapong, R Phuackchantuck, N Morakote, P Somboon, A Saeung, U Chaithong	
Ovicidal and larvicidal properties of <i>Pergularia extensa</i> and <i>Spermacoce hispida</i> ethanol root extracts on <i>Anopheles stephensi</i> Liston	1188
D Abirami, K Kovendan, B Chandramohan	
Molecular epidemiological study on norovirus infection in two distinct hospitals in north-eastern Thailand, 2013-2015	1200
R Guntapong, K Ruchusatsawat, B Suwannakan, N Panthasri, W Kittiwongsunthorn, V Chaichitwanitkul, K Chumpon, R Tacharoenmuang, P Singchai, S Upachai, M Boonchan, N Takeda, S Sangkitporn, K Motomura	
Monoclonal antibody-based latex agglutination for rapid identification of leptospire isolated from patient blood and rat kidney in Thailand	1210
P Ekpo, U Rungpanich, D Suwancharoen, D Phulsuksombath, V Wuthiekanun, Y Suputtamongkol	
Seroprevalence of leptospirosis among town service workers in Kelantan, Malaysia	1222
MA Zainuddin, N Ismail, SA Hassan, MN Shafei, MR Abdullah, ZW Mohamad, NA Yaacob, AM Nikman, A Daud, O Maizurah	
Factors associated with treatment outcomes of patients with extensively drug-resistant gram-negative infections treated with colistin combination therapy: a pilot study	1230
P Rotjanapan, P Kongprasom, P Bunupuradah, S Auparakkitanon, J Krongvorakul, J Sueajai	
Research note. <i>Salmonella</i> prevalence, antibiograms and serotypes from chicken meat and egg in Khon Kaen city, Thailand	1241
P Neunchat, K Yangkao, N Earm-sa-ard, S Chatuphonprasert, J Wiwattrakul, J Manaijsin, N Polruang, A Sopa, P Sukon, K Kanistranon, S Khaeng-air, B Noppon	
Prevalence, virulence profiles, and genetic relationship of atypical enteropathogenic <i>Escherichia coli</i> O145 from beef, southern Thailand	1248
A Sae-lim, P Jearanai, P Rattanachuay, P Sukhumungoon	
Antibacterial effect of the water extract of <i>Houttuynia cordata</i> water extract against multi-drug resistant <i>Escherichia coli</i>	1260
J Li, M Rehman, H Zhang, MK Iqbal, K Mehmood, S Huang, F Nabi	
Epidemiology of <i>Moraxella catarrhalis</i> infections among Thai patients treated at Siriraj Hospital, Thailand during 2012-2015	1267
C Tribuddharat, S Srifuengfung	

Using clinical profile and initial laboratory results to differentiate leptospirosis, scrub typhus and dengue viral infections among children with an acute febrile illness	1274
..... S Assawawiroonhakarn, A Geater, K Laoprasopwattana	
Pneumococcal meningitis at a Thai hospital over a 10-year period: clinical outcomes, serotypes, and antimicrobial susceptibility patterns	1281
..... J Suphanklang, W Santimaleeworagun, S Thunyaharn, J Traipattanakul	
A survey of antibiotic self-medication and over the counter drug use among undergraduate medical students in Yogyakarta, Indonesia	1290
..... S Sandhu, Y Suryani, I Dwiprahasto, J Atthobari	
Risk factors and etiologies of clean and clean contaminated surgical site infections at a tertiary care center in Malaysia	1299
..... WJ Leong, H Hasan, Z Zakaria, MZH Ghazali, SAA Hamid, SA Hassan	
Prevalence of glucose 6-phosphate dehydrogenase deficiency and genetic mutations among Karen and Lao populations in Thailand	1308
..... A Kanchanavithayakul, K Prasittisa, P Kiat-Amornrak, M Chanda, P Kittiwatanasarn, I Nuchprayoon, CL Cheepsunthorn	
Relation between sitting time and sociodemographic factors of Japanese living overseas	1318
..... KP Izawa, K Oka	
Effect of fluoride varnishes enhanced with different calcium phosphate on microhardness of primary teeth	1325
..... P Rirattanapong, K Vongsavan, C Saengsirinavin, N Jantarakam	
Author index	1330
Subject index	1335
Errata	1338

PERMOT (*PASSIFLORA FOETIDA* LINN.) LEAF EXTRACTS AS BIOINSECTICIDE AGAINST *Aedes aegypti* LARVAE

Poedji Hastutiek¹, Agus Sunarso¹ and R Heru Prasetyo²

¹Department of Parasitology, Faculty of Veterinary Medicine;

²Department of Parasitology, Faculty of Medicine, Universitas Airlangga, Indonesia

Abstract. The aim of this study was to investigate the effectiveness of *Passiflora foetida* Linn. extracted by various organic solvents (n-hexane, ethyl acetate and ethanol) against IV instar *Aedes aegypti* larvae. The components fractionated by thin layer chromatography of n-hexane extract were alkaloids and terpenoids, that of ethyl acetate fraction was terpenoid, and that of ethanol fraction was phenol. Gas chromatography-mass spectrometry analysis of n-hexane fraction indicated that the presence of isophytol, neophytadiene, 9,12,15-octadecatrienoic acid, 13-octadecenal, and phytol; and of the n-hexane fraction as isophytol and phytol. Larvae ($n = 25$) were exposed to each organic solvent extract (0, 500, 1,000, 1,500, and 2,000 mg/l) for 24 hours, each experiment conducted four times. Mortality data were analyzed using probit analysis at 95% significance level. The 50% lethal concentration of n-hexane extract, 440 mg/l, was the lowest, but too high to be used effectively against *Aedes aegypti* larvae. Further purification will be required to identify the active larvicidal compound(s).

Keywords: *Aedes aegypti*, *Passiflora foetida*, fourth instar larva, larvicidal activity

INTRODUCTION

Surabaya city, Indonesia is prone to dengue hemorrhagic fever (DHF), with the incidence increasing yearly (Suroso, 1997). The highest incidence of DHF in Surabaya was 58 per 100,000 in 1968, with 24 deaths (a case fatality rate of 41,3% compared to a mean of <1%) (Surabaya City Health Department, 2005). The disease has spread to numerous cities and almost all provinces in Indonesia are affected by 2004 (Soegianto *et al*, 2004).

Dengue is caused by dengue virus transmitted by mosquito vectors *Aedes*

aegypti and *Aedes albopictus* (Sumarmo, 2004). The increasing number of dengue virus infections each year is closely related to environmental sanitation, social-economic conditions and human behavior (Arsin and Wahiduddin, 2004). The most appropriate way to cope with this disease is to control the vectors, a key strategy employed around the world (Okumo *et al*, 2007).

The use of insecticides directed at the larval stages of the mosquito is accepted as the most general way to control these insects. The use of Abate SG (1% temephos sand granules) has been employed in Indonesia since 1976 (Suroso, 1997). Four years later, it was established as part of a mass eradication program of *Ae. aegypti* (Ministry of Health Indonesia, 1999). But the danger of resistance and side effects caused

Correspondence: R Heru Prasetyo, Department of Parasitology, Faculty of Medicine, Universitas Airlangga, Jl. MayJen Prof Dr Mustopo 47, Surabaya 60131, East Java, Indonesia.
E-mail: rheru_prasetyo@yahoo.co.id

by insecticides is inevitable, and resistance of *Ae. aegypti* against temephos was found in Surabaya (Rahardjo, 2006). Given the rapid development of insecticide resistance in vector mosquitoes, the development of alternative sources of insecticides, such as those derived from plants (bioinsecticide) is a sound strategy. Replacement of chemical insecticides with more environmentally friendly bioinsecticides is essential to anticipate the negative health impact of DDT. One of the alleged medicinal plants containing active insecticidal compounds, which can be utilized as a bioinsecticide against *Ae. aegypti* is the plant Permot (*Passiflora foetida* Linn.), which grows wild and can be easily obtained in Indonesia (Wijayakusuma *et al*, 1995).

The purpose of this study was to extract, isolate, and analyze chemical components from Permot leaves and to investigate the potential of the extracts as bioinsecticides against *Ae. aegypti* larvae.

MATERIALS AND METHODS

Collection of Permot leaves

Permot leaves were collected from several locations in Surabaya. Identification was performed at the LIPI Purwodadi Botanical Garden, Pasuruan, East Java. Leaves were cleaned and aerated for 5-7 days by placing them in an area sheltered from the sun, and then ground to powder (simplicia).

Rearing of mosquito larvae

Mosquitoes utilized in this study were fourth instar *Ae. aegypti* larvae. Rearing of *Ae. aegypti* eggs were performed by placing them in plastic trays filled with water at 28.5°C and 72.5 % humidity. Eggs hatched after 1-2 days and the first instar larvae were moved to another tray containing 1 liter of water and fed with

four catfish pellets (Comfeed, Pandaan-Pasuruan, East Java, Indonesia). After 1-2 days the first instar larvae developed into second instar larvae, characterized by exfoliation, which then moved to another tray also contains 1 liter of water and catfish pellets. Within 2-3 days third instar larvae appeared and by the following 2-3 days fourth instar larvae were generated.

Extraction of Permot leaves

One kg of Permot leaf was macerated in 1 liter of n-hexane and left to stand for 24 hours. The solution was then filtered and the process was repeated three times. The residue was re-extracted with 1 liter of ethyl acetate and processed as described above. The residue was again re-extracted with 1 liter of ethanol solvent as described above. The filtrate from each solvent was collected and evaporated using a rotary evaporator under reduced pressure to obtain a viscous extract.

Thin layer chromatography (TLC) analysis of n-hexane, ethyl acetate and ethanol extracts

The presence of alkaloids, flavonoids, phenols, and terpenoids in ethanol, ethyl acetate and n-hexane extracts were tested by TLC using aluminium foil-backed silica gel and hexane as mobile phase (followed by treatment with Dragendorff reagent), or a mixture of n-hexane, ethyl acetate (4:1) (treatment with anisaldehyde sulfuric acid), or a mixture of chloroform:ethyl acetate:formic acid (0.5:9:0.5) (treatment with FeCl₃).

Identification of chemical compounds by gas chromatography-mass spectrometry (GC-MS)

Fraction containing the lowest LC₅₀ value (concentration producing 50% lethality) was analyzed using GC-MS. The sample solution inserted to GC-MS 2010 S

(Shimadzu, Portland, OR). Identification of peaks was performed based on database from LIBRARY WILEY7-LIB.

Determination of LC₅₀ value

Ethanol, ethyl acetate and n-hexane extracts were emulsified in distilled water using Tween 80 and dimethyl sulf-oxide (4:1 v/v). Fourth instar larvae of *Ae. aegypti* ($n = 25$) were soaked for 24 hours with each solution of 0, 500, 1,000, 1,500, and 2,000 mg/l. Larvae are scored as killed when they did not move or respond to mechanical stimuli. Each experiment was repeated four times.

Data analysis

Relationship between concentration test solution and mortality of fourth instar *Ae. aegypti* larvae was analyzed using Probit for Windows SPSS 10 (SPSS, Chicago, IL) to obtain lethal LC₅₀ values at a significance level of 95%.

RESULTS

TLC identification of Permot leaf ethanol, ethyl acetate and n-hexane extracts

One kg of Permot leaf of *Simplicia* yielded 126, 27 and 34 g of ethanol, ethyl acetate and n-hexane extract, respectively, which were separated by TLC. N-hexane extract was positive with Dragendorff reagent (orange red) indicating the presence of alkaloids (Fig 1A). A mobile phase of a mixture of n-hexane:ethyl acetate (4:1) and anisaldehyde sulfuric acid treatment resulted in red purple or




Fig 1—Thin layer chromatograms of Permot leaf ethanol (EOH), ethyl acetate (EA) and n-hexane (H) extracts. A) Separation with hexane and treatment with Dragendorff reagent. B) Separation with a mobile phase of mixture of n-hexane, ethyl acetate (4:1), and treatment with anisaldehyde sulfuric acid. C) Separation with a mobile phase of a mixture of chloroform:ethyl acetate:formic acid (0.5:9:0.5) and treatment with FeCl₃.

violet color indicative of the presence of terpenoids in n-hexane and ethyl acetate fractions (Fig 1B). A mobile phase of a mixture of butanol:glacial acetic acid:water (4:1:5) and treatment with chloroform:acetone:formic acid (6:6:1) under ammonia vapor failed to produce a yellow color in any of the three extracts indicating the absence of flavonoids (data not shown). A mobile phase of a mixture of chloroform:ethyl acetate:formic acid (0.5:9:0.5) and FeCl₃ treatment generated a black stain with the ethanol extract indicating the presence of phenols (Fig 1C).




Fig 2—Gas chromatography-mass spectrometry profile of Permot leaf n-hexane extract displayed 28 peaks GC-MS indicated that the n-hexane extract contained alkaloids and terpenoids (isophytol and phytol compound). Five peaks of which were subsequently shown to be isophytol, neophytadiene, 9,12,15-octadecatrienoic acid. GC-MS used for analyzing fraction containing the lowest LC₅₀ value. The sample solution inserted to GC-MS. Based on chromatogram of GC can be known the profile of peak of sample components. Result of quality analysis shown with a lot of peak that represents a lot of compounds. In quantitative analysis the height or peak area represents the relative percentage of the compound. Numbers above peaks shows time retention (in minute).

GC-MS identification of Permot leaf hexane extract

GC-MS analysis of Permot leaf n-hexane extract displayed 28 peaks (Fig 2), five peaks of which were subsequently shown to be isophytol, neophytadiene, 9,12,15-octadecatrienoic acid, 13-octadec-

nal, and phytol (data not shown).

Toxicity tests of Permot leaf extracts against fourth instar *Ae. aegypti* larvae

Probit analysis of the toxicity of Permot leaf extracts (concentrations ranging from 500 to 2,000 mg/l) against fourth instar *Ae. aegypti* larvae revealed among the three extracts that of n-hexane was the most potent, with an IC₅₀ of 440 mg/l (Table 1).

DISCUSSION

Of the three organic solvent extract, namely, ethanol, ethyl acetate and n-hexane, of Permot leaf the latter extract possessed the lowest LC₅₀ against fourth instar *Ae. aegypti* larvae. TLC of Permot leaf n-hexane extract indicated that it contained alkaloids and terpenoids. Wijayakusuma *et al* (1995) found alkaloid compounds in Permot

plant, which possess the ability to work as a poison against mosquito larvae and are more abundant in leaf than in other parts of the plant. GC-MS further indicated that the n-hexane extract in addition contained isophytol and phytol compounds, both of which are terpenoids, in line with the

Table 1
Probit analysis of LC₅₀ and LC₉₅ of Permot leaf extracts against fourth instar *Aedes aegypti* larvae after 24 hours exposure.

Extract	LC50 (mg/l)	LC95 (mg/l)	Regression equation
n-hexane	440	1,796	Y = 2.692 X - 7.116
Ethyl acetate	2,122	8,668	Y = 2.692 X - 7.116
Ethanol	3,076	12,560	Y = 2.692 X - 7.116

LC_n = concentration producing n% lethality.

findings of Estrada *et al* (2013) suggesting phytol is the main component of *P. alliaceae* leaf. Mathew and Thoppil (2011) reported that the important components in essential oil of *Salvia splendens* acting as larvicidal against *Ae. albopictus* larvae are phytol and cyclooctasulfur.

Other studies have shown that plant-based insecticide has the power to kill larvae and mosquitoes. For example extract of papaya seeds and leaves have LC50 442,311 mg/l against *Anopheles* sp larvae (Hastuty, 2014). Another study showed larvicidal activity of eucaplitol and camphor from Rosemary leaf extract against *Culex quinquefasciatus* (Yu *et al*, 2013).

Alkaloids possess the ability to work as a good contact poison because of their ability to penetrate the insect cuticle. They are very effective against a variety of insects, especially those with soft-bodies. In larvae, alkaloids work well as a contact poison and stomach poison. These compounds work at the the insect central nervous system ganglia (Soparat, 2010). Alkaloids can also cause gastrointestinal poison by entering through the larvae mouth (Soparat, 2010). However, alkaloids function slowly in insects as they inhibit feeding (a stop-feeding action).

In summary, the study shows that n-hexane extract of Permot leaf was a better biolarvacide against *Ae. aegypti*

compared to ethanol or ethyl acetate extract. The n-hexane containing alkaloids and terpenoids (isophytol and phytol). However, the LC₅₀ value against *Aedes aegypti* (fourth instar larvae) was considered too low to be used effectively, and future studies should be directed towards isolating and identifying the active components, including toxicological data and effects on non-target organisms and the environment.

ACKNOWLEDGEMENTS

The research (PUPT) was funded by DIPA BOPTN Universitas Airlangga, fiscal year 2013. The authors express their gratitude to the Chairman of LPPM Universitas Airlangga, the Director of the Institute of Tropical Disease, Universitas Airlangga, and the Chairman of Department of Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga.

REFERENCES

- Arsin AA, Wahiddudin. Factors that influence dengue fever in Makassar city. *J Kedokteran Yarsi* 2004; 12: 23-33 (in Indonesian).
- Estrada AC, Angulo MG, Argáez RB, Sánchez ER. Insecticidal effects of plant extracts on immature whitefly *Bemisia tabaci* Genn. (Hemiptera: Aleyroideae). *E J Biotechnol Plant Biotechnol* 2013; 16 (1).

- Hastuty, Henny Sesanti Budi. Test the potential of *Carica papaya* leaf and seed ekstrak as larvaside to *Anopheles* larvae in Jayapura Regency Papua (Script). Makassar: Universitas Hassanuddin. 2014 (in Indonesian).
- Mathew J, Thoppil JE. Chemical composition and mosquito larvicidal activities of *Salvia* essential oils. *Pharmaceut Biol* 2011; 49: 456-63.
- Ministry of Health Indonesia (Department of Kesehatan Republik Indonesia, DepKes RI): Guidance book for DHF prevention and Control in Indonesia Packchage B. Jakarta. DepKes RI, 1999: 1-5.
- Okumu FO, Knols BGJ, Fillinger U. Larvasidal effect of neem (*Azadiracht indica*) oil formulation on the malaria vector *Anopheles gambiae*. *Malaria J* 2007; 6: 63.
- Raharjo B. Susceptibility of *Aedes aegypti* (Linnaeus) larvae from Surabaya, Palembang and in some area in Bandung to Temephos (Abate 1 SG). Bandung: School of Biological Science, Institut Teknologi Bandung, 2016 (in Indonesian).
- Soegianto S, Sustini F, Wirahjanto A. Epidemiology of dengue haemorrhagic fever. In: Dengue haemorrhagic fever, review and the new findings in 2003 era. 1st ed. Surabaya: Airlangga University Press Surabaya, 2004: 1-10 (in Indonesian).
- Soparat S. Chemical ecology and function of alkaloids, Bangkok: Kasetsart University, 2010. [Cited 2011 Sept 10]. Available from: <http://pirun.ku.ac.th/g4686045/media/alkaloid.pdf>
- Sumarmo PS. The problem of dengue haemorrhagic fever in Indonesia. Jakarta: Universitas Indonesia, 2004: 1-13 (in Indonesian).
- Surabaya Health City Office (Dinas Kesehatan Kota Surabaya). Incidence rate and case fatality rate of dengue haemorrhagic fever in Surabaya city in last 5 years, 2005. Surabaya: Surabaya Health Office, 2005.
- Suroso T. A review of dengue hemorrhagic fever and its control in Indonesia. Yogyakarta: Seminar on Recent Advance in Molecular Diagnostic, 1997.
- Wijayakusuma H, Dalimarta S, Wirian AS. Medicinal plants in Indonesia. Pustaka Jakarta 1995: 106-7 (in Indonesian).
- Yu J, Liu XY, Yang B, et al. Larvicidal activity of essential extract of *Rosmarinus officinalis* against *Culex quinquefasciatus*. *J Am Mosq Control Assoc* 2013; 29: 44-8.