

Earphone Swab-Derived 126bp and 143bp mtDNA at D-loop Region - Electrophoretic Band Reaction on Room Temperature

by Ahmad Yudianto

Submission date: 03-Sep-2020 06:27PM (UTC+0800)

Submission ID: 1378817475

File name: p_Region_-_Electrophoretic_Band_Reaction_on_Room_Temperature.pdf (574.36K)

Word count: 2555

Character count: 13791

1 Earphone Swab-Derived 126bp and 143bp mtDNA at D-loop Region: Electrophoretic Band Reaction on Room Temperature

Ahmad Yudianto^{1,2} Simon Martin Manyanza Nzilibili^{1,3}

¹Forensic Science Program, Post Graduate School, Universitas Airlangga, 4-6 Airlangga Rd., 60286 Surabaya – Indonesia.

²Department of Forensic and Medico-legal, Faculty of Medicine, Universitas Airlangga, Surabaya – Indonesia

³Ministry of Health, Community Development, Gender Elderly and Children, Dodoma – Tanzania.

Keywords: DNA quality, Earphone swabs, Forensic evidence, Mitochondrial DNA, Room temperature.

Abstract: Exploration of crime scene devices, leftover, and its surrounding is of significance in forensic identification due to its potential harbor of biological evidences. Such harboring of evidences proposes traceability under Locard's principle. By body use-contact, devices such as earphones potentiate revelation of the unknown by analysis of cerumen or traces logged from ear. Despite of this usability, environmental challenge is suggested to impact the forensic vitality of the logged evidences. To understand the phenomena, this study applied 30 used earphones at 0, 1, 7, 14 and 20 days room temperature exposure. From the study; amplified electrophoretic reaction showed that, 143bp (HVS I, nt 16268-16410) mtDNA D-loop region reacted with a positive (+) detection only for day 1 (4 [66.67%] samples) and day 7 (3 [50%] samples) of room temperature exposure. Meanwhile, visualization of the results of 126bp (HVS II, nt 34-159) mtDNA D-loop region showed a positive (+) detection only for day 1 (2 [33.37%] of samples) and day 7 (6 [100%] of samples) of room temperature exposure. Data interpretation conclude the effect of room temperature on the trend and quality of earphone swab-derived DNA from day 1 to day 20.

1 BACKGROUND

Forensic identification by means of Deoxyribonucleic Acid (DNA) analysis is modern and referred as among of the three primary independent preferred methods in practice (Dumache et al. 2016; Ruitberg et al. 2001). Its accuracy is by its stability inside biological evidences (International Committee of the Red Cross 2009). Traces of these biological evidences are either freely dispersed at criminal environment or found logged and harbored in objects/devices lastly used and left at the scene (Ah Van Oorschot et al. 2010; Song et al. 2015; Shaler 2011). Such devices harboring samples include earphones under the use-contact theory by Locard's principle (Byard et al. 2016). Logging of the biological samples into earphone attached to the outer ear skin is by expelled cerumen and degrading body cells (Shokry & Filho 2017). In forensic, these devices are becoming useful and crucial in investigative tasks as reported by Seo et al. (2002) though challenged by exposing condition.

Among the challenging condition aggregated by environmental factors, temperature is potentially studied to impact successful profiling hence hindered identification. (Caputo et al. 2011; Toothman et al. 2008). Theory to improve analysis have opted for mini-primer (Butler 2007; Hwan et al. 2008). The method profile reduced size of Short Tandem Repeat (STR) locus (Coble & Butler 2005) also hyper variable segment [HVS] 1 or 2 of mitochondrial DNA (mtDNA), as amplification products (Gabriel et al. 2001). This study, therefore, present the association of room temperature on electrophoretic visualisation as part of the project analysed the effect of room temperature on the quality of earphone swab-derived DNA at 126bp and 143bp.

2 MATERIALS AND METHODS

The laboratory experimental study designed a time serial of exposure of 0, 1, 7, 14 and 20 days interval. Finally, analysis presented the electrophoretic visualization of earphone swab-derived

mitochondrial DNA resulted from room temperature effect on logged samples.

2.1 Volunteer Consent and Sample Preparation

A total of 30 samples of earphone swab derived DNA was taken from respondents who have agreed and signed a written informed consent. The materials of study were earphone swabs.

2.2 Experimentation protocol

The materials for DNA extraction were DNAzol reagent (Invitrogen, ThermoFisher Scientific, Waltham, MA, USA) and 70% ethanol (EMSURE®, Merck KGaA 64271 Darmstadt, Germany). The material for Polymerase Chain Reactions (PCR) were PCR mix - 12.5 µl (Promega Corporation, Madison, USA) comprising of dNTPs (ATP, CTP, TTP GTP), MgCl₂, Taq polymerase and Nuclease-free water; mtDNA primers - 143bp (HVS I, nt 16268-16410) AFDIL 5'CACTAGGATACCAACAAACC3' and 5'GAGGATGGTCAAGGGAC3' according to Edson et al. (2004), 126bp (HVS II, nt 34-159) AFDIL 5'TCTCCAAGCGGGTGGTGCATTTA3' and 5'AAATAATAGGCAGGAGAGGATATC3' according to Gabriel et al. (2001).

3 RESULTS AND DISCUSSION

Polymerase Chain Reaction Amplification and Electrophoresis Detection

Cell lysed and isolated during extraction produced DNA template for PCR processing. The template under this study was polymerized and amplified using two hyper variable segments: (HVS) II (126bp, nt 34-159) and HVS I (143bp, nt 16268-16410) of D-Loop region of the mtDNA as shown in Figure 1 to Figure 4.

143bp and 126bp mtDNA amplicon profiling at day 1 and 7

Profiling of 143bp and 126bp resulted into a band contrast formation. Samples amplification became positively detected on both primers (143bp and 126bp) from day one and seven, though two samples under 126 appeared fainter after 7 days exposure (Figure 1).




Figure 1: Electrophoretic visualization of 143-bp and 126-bp PCR products of earphone swab samples (M = marker, H1& H7 = samples at day 1 and 7)

143bp mtDNA amplicon electrophoresis of day 1, 7, 14 and 20

Under 2% agarose gel electrophoresis 143bp DNA amplicon, day 1 present positive [+] detection of two samples with faint bands and other 2 samples demonstrated a negative [-] detection. Day 7, 3 samples detected negatively [-] (no band displayed) and there was only one sample that showed a positive [+] detection with a faint band too. Amplification and electrophoresis of samples by 143bp (HVS I, nt 16268-16410) D-loop mitochondrial DNA amplicon on day 14 and 20 using 2% agarose gel displayed no band or negative detection as shown in Figure 3.




Figure 2: Electrophoretic visualization of 143-bp PCR products of earphone swab samples (M = marker, H1& H7 = samples at day 1 and 7)




Figure 3: Electrophoretic visualization of 143-bp PCR products of earphone swab samples (M = marker, H14 & H20 = samples at day 14 and 20)

126bp mtDNA amplicon electrophoresis of day 7, 14 and 20

Amplification and electrophoretic visualization of 126bp mitochondrial DNA amplicon using 2% agarose gel on PCR swabbed earphone products exposed at 7, 14, and 20 intervals displayed variably. Day 7 exposed sample amplification gave a positive [+] contrast detection, bands were visible for the mtDNA as shown in presented in Figure 4. Contrary to day seven, amplification and electrophoresis of samples exposed for 14 and 20 days resulted into a no display of band contrast hence a negative [-] detection.




Figure 4: Visualization of 126-bp PCR products of earphone swab samples (M = marker, H7, H14 & H20 = samples at day 7, 14 and 20)

Relative extent effect of exposition on day 1, 7, 14 and 20

A representation of six samples for general percentage effect according to analysis is summarized in Table 1. The results show that, detection of the effects of room temperature exposure on earphone swab-derived 143bp (HVS I, nt 16268–16410) and 126bp (HVS II, nt 34–159) mtDNA D-loop region depend also on the exposure interval and the D-loop specific region between the two (143bp and 126bp). Day one exposition analysis, 66.67% (4) of samples were positively detected at 143bp (HVS I, nt 16268–16410) D-loop region while 33.37% (2) were negatively detected

similar to those positively detected at 126bp (HVS II, nt 34–159) D-Loop region. On day 7: 50% (3) of samples were detected at 143bp (HVS I, nt 16268–16410) D-loop region and 100% (6) of samples detected at 126bp (HVS II, nt 34–159) D-loop region. Day 14 and 20, visualized a zero percentage of DNA positive detection due to exposure effect to room temperature on all earphone swab samples (143-bp (HVS I, nt 16268–16410) and 126bp (HVS II, nt 34–159) D-loop region).

Table 1: Results of detection of the effect of room temperature exposure on earphone swab-derived 143-bp (HVS I, nt 16268–16410) and 126-bp (HVS II, nt 34–159) mtDNA D-loop region.

Exposure Interval	HVS I 143bp nt: 16268 – 16410		HVS II 126bp nt: 34 – 159	
	Detection (+)	Detection (-)	Detection (+)	Detection (-)
Days 1	4(66.7%)	2(33.3%)	2(33.3%)	4(66.7%)
Days 7	3(50%)	3(50%)	6(100%)	0(0%)
Days 14	0(0%)	6(100%)	0(0%)	6(100%)
Days 20	0(0%)	6(100%)	0(0%)	6(100%)

4 DISCUSSION

126bp and 143bp mtDNA Electrophoretic Band Reaction on Room Temperature

According to bands formation, analysis descriptively derived information from detection and contrasting. Detection of effects of room temperature is related to DNA quality contained. DNA quality includes DNA levels, DNA purity and DNA condition state (degradation). DNA degradation is suggested to be the causes of DNA detection failure in PCR-based DNA analysis, this is well portrayed in Figure 3 and some bands in Figure 2 and 4. This is in consistent with (Bartlett & Stirling 2003) with regard to several possible causes for failure detection of DNA, including a minimal amount of target DNA, DNA degradation that prevent primer from annealing, lack of DNA polymerase and PCR cycle also presence of PCR inhibitors. The failure of amplification is characterized by the absence of bands on the

electrophoresis results due to lack of DNA polymerase and PCR cycle. This can be controlled by the use of PCR master mix (Lorenz 2012). From the article lack of cycle in a PCR reaction is controlled by optimizing the primers used for PCR while adhering to Mg^{2+} recommended concentration (should be 1.5 to 4 mM). An excessive amount of taq polymerase in PCR mix causes a smear formation (non-specific reaction) on electrophoresis, whereas inadequacy of it causes a reduced efficiency of amplification and non-appearance of bands.

According (Hall et al. 2014; Schuch et al. 2013), DNA damage caused by external factors such as extreme temperature will result in many types of damage, like damage to DNA strands (both double and single strand), base damage (damage to DNA bases), sugar damage and even presence of DNA crosslink and DNA-protein crosslink – hence DNA degradation. This degradation can be fast or slow, depending on the affecting factors and length of exposure. Findings of this study shows that earphone swabs can still be used as a material for DNA analysis of mtDNA with regard to the effects of exposure to room temperature (28.5°C–30°C) up to day 7. This is in potential consideration when applied to expedite forensic identification, especially in a mass disaster or criminal cases on the crime scene with little trace evidence.

5 CONCLUSION

This study concludes that; the length of exposure to room temperature has effect on the quality of earphone swab-derived DNA. Thereby visualization of the results of PCR on 143bp (HVS I, nt 16268-16410) mtDNA D-loop region showed a positive (+) detection only for day 1 (4 [66.67%] samples) and day 7 (3 [50%] samples) of room temperature exposure. Meanwhile, visualization of the results of PCR on 126bp (HVS II, nt 34-159) mtDNA D-loop region showed a positive (+) detection on similar exposure interval - day 1 (2 [33.37%] samples) and day 7 (6 [100%] samples) of room temperature.

ACKNOWLEDGEMENTS

We appreciate participation of volunteers and all who made this research possible.

REFERENCES

- Ah Van Oorschot, R., Ballantyne, K.N. & Mitchell, R.J., 2010. Forensic trace DNA: a review. *Investigative Genetics*, 1, p.14.
- Anon, PCR Troubleshooting | LSR | Bio-Rad. Available at: <https://www.bio-rad.com/en-id/applications-technologies/pcr-troubleshooting?ID=LUSO3HC4S> [Accessed May 14, 2018].
- Bartlett, J.M.S. & Stirling, D., 2003. A short history of the Polymerase Chain Reaction. *Humana Press*, 226(PCR Protocols Volumen 226 de Methods in molecular biology), p.545. Available at: <http://www.springerprotocols.com/Abstract/doi/10.1385/1-59259-384-4:3#>.
- Butler, J.M., 2007. Short tandem repeat typing technologies used in human identity testing. *BioTechniques*, 43(4).
- Byard, R.W. et al., 2016. Locard's Principle of Exchange, Dental Examination and Fragments of Skin. *Journal of Forensic Sciences*, 61(2), pp.545–547.
- Caputo, M., Bosio, L.A. & Corach, D., 2011. Long-term room temperature preservation of corpse soft tissue: an approach for tissue sample storage. *Investigative genetics*, 2(1), p.17. Available at: <http://www.investigativegenetics.com/content/2/1/17>.
- Coble, M.D. & Butler, J.M., 2005. Characterization of New MiniSTR Loci to Aid Analysis of Degraded DNA. *Journal of Forensic Sciences*, 50(1), pp.1–11. Available at: <http://www.astm.org/doiLink.cgi?JFS2004216>.
- Dumache, R. et al., 2016. Molecular DNA analysis in forensic identification. *Clinical Laboratory*, 62(1–2), pp.245–248. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27012057> [Accessed May 11, 2018].
- Edson, S.M. et al., 2004. *Naming the Dead - Confronting the Realities of Rapid Identification of Degraded Skeletal Remains*.
- Gabriel, M.N. et al., 2001. *Improved MtDNA Sequence Analysis of Forensic Remains Using a "Mini-Primer Set" Amplification Strategy*.
- Hall, A., Sims, L.M. & Ballantyne, J., 2014. Assessment of DNA damage induced by terrestrial UV irradiation of dried bloodstains: Forensic implications. *Forensic Science International: Genetics*, 8(1).
- Hwan, Y.L. et al., 2008. A modified mini-primer set for analyzing mitochondrial DNA control region sequences from highly degraded forensic samples. *BioTechniques*, 44(4), pp.555–558. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18476821> [Accessed May 12, 2018].
- International Committee of the Red Cross, 2009. *Missing People, DNA Analysis and Identification of Human Remains*. Available at: www.icrc.org/eng/resources/documents/publication/p4010.htm.
- Lorenz, T.C., 2012. Polymerase Chain Reaction: Basic

- Protocol Plus Troubleshooting and Optimization Strategies. *Journal of Visualized Experiments*, (63), pp.1–15. Available at: <http://www.jove.com/video/3998/>.
- Ruitberg, C.M., Reeder, D.J. & Butler, J.M., 2001. STRBase: a short tandem repeat DNA database for the human identity testing community. *Nucleic Acids Research*, 29(1), pp.320–322. Available at: <https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/29.1.320>.
- Schuch, A.P. et al., 2013. DNA damage as a biological sensor for environmental sunlight. *Photochemical & Photobiological Sciences*, 12(8), p.1259. Available at: <http://xlink.rsc.org/?DOI=c3pp00004d>.
- Seo, Y. et al., 2002. Mitochondrial DNA and STR typing of matter adhering to an earphone. *Journal of forensic sciences*, 47(3), pp.605–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12051345> [Accessed May 12, 2018].
- Shaler, R.C., 2011. Chapter 13: The Biological Crime Scene. In *Crime Scene Forensics: A Scientific Method Approach*. pp. 338–373.
- Shokry, E. & Filho, N.R.A., 2017. Insights into cerumen and application in diagnostics: Past, present and future prospective. *Biochemia Medica*, 27(3).
- Song, G.H. et al., 2015. Effects of fluoride on DNA damage and caspase-mediated apoptosis in the liver of rats. *Biological Trace Element Research*, 166(2), pp.173–182.
- Toothman, M.H. et al., 2008. Characterization of human DNA in environmental samples. *Forensic science international*, 178(1), pp.7–15. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18420364> [Accessed April 4, 2017].

Earphone Swab-Derived 126bp and 143bp mtDNA at D-loop Region - Electrophoretic Band Reaction on Room Temperature

ORIGINALITY REPORT

7%

SIMILARITY INDEX

6%

INTERNET SOURCES

3%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** unair.ac.id Internet Source 4%
- 2** ejfs.springeropen.com Internet Source 1%
- 3** Agung Sosiawan, Ahmad Yudianto, Abdul Hadi Furqoni, Simon Martin Manyanza Nzilibili, Indah Nuraini. "Full-sibling allelic frequency and sharing among Madurese: STR technique by 12 locus and the sex-typing amelogenin gene", Egyptian Journal of Forensic Sciences, 2019
Publication 1%
- 4** Sandrine Perrier, Jörg Hau, Didier Gasparutto, Jean Cadet, Alain Favier, Jean-Luc Ravanat. "Characterization of Lysine–Guanine Cross-Links upon One-Electron Oxidation of a Guanine-Containing Oligonucleotide in the Presence of a Trilysine Peptide", Journal of the American Chemical Society, 2006
Publication 1%

5

e-journal.unair.ac.id

Internet Source

1%

6

Simon Martin Manyanza Nzilibili, Moh. Kurniadi
Hendry Ekodiyanto, Pudji Hardjanto, Ahmad
Yudianto. "Concentration and Purity DNA
Spectrophotometer: Sodium
Monofluorophosphate forensic impended effect",
Egyptian Journal of Forensic Sciences, 2018

Publication

<1%

7

Margaret Cox, Peter Jones. "Ethical
considerations in the use of DNA as a
contribution toward the determination of
identification in historic cases: considerations
from the Western front", New Genetics and
Society, 2014

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Earphone Swab-Derived 126bp and 143bp mtDNA at D-loop Region - Electrophoretic Band Reaction on Room Temperature

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
